

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 August 2002 (22.08.2002)

PCT

(10) International Publication Number
WO 02/063975 A2

(51) International Patent Classification⁷: **A23K**

(21) International Application Number: PCT/US02/04919

(22) International Filing Date: 14 February 2002 (14.02.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/269,188 14 February 2001 (14.02.2001) US
09/847,232 2 May 2001 (02.05.2001) US

(71) Applicant: **VENTRIA BIOSCIENCE** [US/US]; 4110 North Freeway Boulevard, Sacramento, CA 95834-1219 (US).

(72) Inventors: **HUANG, Ning**; 1226 Caricia Drive, Davis, CA 95616 (US). **RODRIGUEZ, Raymond, L.**; 3017 Anza Avenue, Davis, CA 95616 (US). **HAGIE, Frank, E.**; 4400 Truxel Road, No. 191, Sacramento, CA 95834 (US).

(74) Agents: **ARNETT, Stephen, E.** et al.; Perkins Coie LLP, P.O. Box 2168, Menlo Park, CA 94026 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FEED ADDITIVE COMPOSITIONS AND METHODS

(57) Abstract: The invention is directed to improved animal feed compositions comprising one or more milk proteins produced in the seeds of a magnetic plant and methods of making and using the same.

WO 02/063975 A2

FEED ADDITIVE COMPOSITIONS AND METHODS

References

The following references are cited herein, and to the extent they may be
5 pertinent to the practice of the invention, are incorporated herein by reference.

Alber and Kawasaki, *Mol. and Appl. Genet.* 1:419-434, 1982.

Aniansson, G, et al. (1990). *Anti-Adhesive Activity Of Human Casein Against Streptococcus Pneumoniae And Haemophilus Influenzae*. Micro Pathog 8(5):315-323.

Arnold R.R. et al., *Infect Immun.* 28, 893-898, 1980.

10 Ausubel F.M. et al., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, N.Y., 1993.

Bhan, M.K. et al., *J Pediatr Gastroenterol Nutr* 7:208-213, 1988.

Beatty, K., et al., Kinetics of association of serine proteinases with native and oxidized alpha-1-proteinase inhibitor and alpha-1-antichymotrypsin. *J. Biol. Chem.*
15 255:3931-3934, 1980.

Boesman-Finkelstein M. and Finkelstein R.A. *FEBS Letters*, 144, 1-5, 1982.

Bradford M. *Analytical Biochem.* 72, 248-254, 1976.

Brandt, et al., *Carlsberg Res. Commun.* 50, 333-345, 1985.

Briggs, D, et al. (1981). Malting And Brewing Science - Volume 1, Malt And Sweet Wort (Chapman & Hall, New York, 2nd edition).
20

Briggs, D (1998). Malts And Malting (Blackie Academic And Professional, New York).

Bullen J.J. et al., *Br. Med. J.* 1, 69-75, 1972.

Carrell, R.W. et al. Structure and variation of human alpha-1-antitrypsin. *Nature*
25 298:329-334, 1983.

Castañón M.J. et al., *Gene*, 66, 223-234, 1988.

Chandan R.C., *J Dairy Sci*, 51: 606-607, 1968.

Chong, D.K. and Langridge, W.H., *Transgenic Res* 9: 71-8, 2000.

Chowanadisai, W. and Lönnerdal, B. Alpha-1-antitrypsin and antichymotrypsin in
30 human milk: origin, concentrations, and stability. *Am. J. Clin. Nutr.* 2001:(in press)

Chrispeels K., *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53, 1991.

Clark et al., *J. Biol. Chem.* 264:17544-17550, 1989.

Davidson LA, et al. (1987). *Persistence Of Human Milk Proteins In The Breast-Fed Infant*. Acta Pediatrica Scand 76(5):733-740.

Davidson, L.A., and Lönnerdal, B. Fecal alpha-1-antitrypsin in breast-fed infants is derived from human milk and is not indicative of enteric protein loss. *Acta Paediatr. Scand.* 79:137-141, 1990.

della-Cioppa *et al.*, *Plant Physiol.* 84:965-968, 1987.

5 Dellaporta S.L. *et al.*, *Plant Mol. Biol. Rep.* 1, 19-21, 1983.

Depicker *et al.*, *Mol. Appl. Genet.* 1:561-573, 1982.

Dewey K.G. *et al.*, *J Pediatrics* 126: 696-702, 1995.

Dewey K.G. *et al.*, *Pediatrics* 89: 1035-1041, 1992.

Dewey K.G. *et al.*, *Am J Clin Nutr* 57: 140-145, 1993.

10 Ditta *et al.*, *Proc. Nat. Acad. Sci., U.S.A.* 77:7347-7351, 1980.

Doncheck, J (1995). *Malts And Malting*. In: Kirk-Othmer Encyclopedia Of Chemical Technology (John Wiley & Sons, New York, 4th edition), volume 15, pp. 947-962.

Eley, E (1990). *A Wandful World*. Food Processing August 1990, pp. 17-18.

15 Faure A. and Jollès P., *et al.*, *Comptes Rendus Hebdomadaires des Seances de L Academie des Sciences. D: Sciences Naturelles*, 271, 1916-1918, 1970.

Fujihara T. and Hayashi K., *Archives of Virology* 140:1469-1472, 1995.

Gastañaduy, A. *et al.*, *J Pediatr Gastroenterol Nutr* 11:240-6, 1990.

Gelvin, S.B. *et al.*, eds. *PLANT MOLECULAR BIOLOGY MANUAL*, 1990.

20 Gielen, *et al.*, *EMBO J.* 3:835-846, 1984

Goto, F *et al.*, *Nature Biotech.* 17, 282-286, 1999.

Grover M. *et al.*, *Acta Paediatrica* 86:315-316, 1997.

Hamosh, M, *et al.* (1999). *Protective Function Of Human Milk: The Fat Globule*. Semin Perinatol 23(3):242-9.

25 Harmsen M.C. *et al.*, *Journal of Infectious Diseases* 172:380-388, 1995.

Hickenbottom, JW (1977). *Sweeteners In Biscuits And Crackers*. The Bakers Digest, December 1977, pp. 18-22.

Hickenbottom, JW (1983). AIB Research Department Technical Bulletin, V(3), March 1983, pp. 1-8.

30 Hickenbottom, JW (1996). *Processing, Types, And Uses Of Barley Malt Extracts And Syrups*. Cereal Foods World 41(10): 778-790.

Hickenbottom JW (1997). *Malts In Baking*. American Society Of Bakery Engineers, Technical Bulletin #238, August 1997, pp. 1010-1013.

Huang N. *et al.*, *Plant Mol. Biol.*, 23, 737-747, 1993.

Hough, JS, et al. (1982).). Malting And Brewing Science - Volume 2, Hopped Wort And Beer (Chapman & Hall, New York, 2nd edition).

Huang N. et al., *J CAASS*, 1, 73-86, 1990a.

Horvath H. et al., *Proc. Natl. Acad. Sci. USA*, 97, 1914-1919, 2000.

5 Humphreys, D.P. and Glover, D.J. Therapeutic antibody production technologies: molecules, applications, expression and purification. *Curr Opin Drug Discov Devel* Mar;4(2):172-85, 2001.

Huang N. et al., *Plant Mol. Biol.* 14, 655-668, 1990b.

10 Huang, J., et al., Expression and purification of functional human alpha-1-antitrypsin from cultured plant cells. *Biotechnol. Prog.* 17:126-133, 2001.

Jensen L.G. et al., *Proc. Natl. Acad. Sci. USA*, 93, 3487-3491, 1996.

Jigami Y. et al., *Gene*, 43, 273-279, 1986.

Johnson, T. Human alpha-1-proteinase inhibitor. *Methods Enzymol.* 80:756-757, 1981.

15 Jollès P., *LYSOZYMES--MODEL ENZYMES IN BIOCHEMISTRY AND BIOLOGY*. Birkhäuser Verlag, Basel; Boston, 1996.

Jouanin et al., *Mol. Gen. Genet.* 201:370-374, 1985.

Kortt A.A. et al., Dimeric and trimeric antibodies: high avidity scFvs for cancer targeting. *Biomol Eng* 18:95-108, 2001.

20 Kovar M.G. et al., *Pediatrics* 74: 615-638, 1984.

Kunz C. et al., *Clin. Perinatol.* 26(2):307-33, 1999.

Lake, C. (1988). *From Strength To Strength. Flood Flavorings, Ingredients, Packaging And Processing*, February 1988, pp. 55 -57.

Langridge et al., *Planta* 156,166-170, 1982.

25 Larrick, J.W. et al., Production of antibodies in transgenic plants. *Res Immunol* 149:603-8, 1998.

Lásztity, R. The chemistry of cereal proteins. CRC Press, Boca Raton, 1996.

Lee-Huang S. et al., *Proc. Natl. Acad. Sci. USA*, 96, 2678-2681, 1999.

Lindberg, T. Protease inhibitors in human milk. *Pediatr. Res.* 13:969-972, 1979.

30 Lollike K. et al., *Leukemia*, 9, 206-209, 1995a.

Lönnerdal B., *Am L Clin Nutr*, 42: 1299-1317, 1985.

Lönnerdal, B. Recombinant milk proteins- an opportunity and a challenge. *Am. J. Clin. Nutr.* 63:622S-626S, 1996.

Maga E. et al., *J. of Food Protection*, 61, 52-56, 1998.

35 Maga E. et al., *Transgenic Research*, 3, 36-42, 1994.

- Maga E. *et al.*, *Journal of Dairy Science*, 78, 2645-2652, 1995.
- Maniatis, *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL*, 2nd Edition, 1989.
- Matsumoto, A. *et al.*, *Plant Mol Bio* 27:1163-72, 1995.
- 5 Maynard, J. and Georgiou, G. Antibody engineering *Annu Rev Biomed Eng* 2:339-76, 2000.
- McBride and Summerfelt, *Plant Mol. Biol.* 14:269-276, 1990.
- Mcgilligan, KM, *et al.* (1987). *Alpha-1-Antitrypsin Concentration In Human Milk. Pediatr Res* 22(3):268-270. Mitra A. and Zhang Z., *Plant Physiol.* 106, 977-981, 1994.
- 10 McGilligan, K.M. *et al.*, Alpha-1-antitrypsin concentration in human milk. *Pediatr. Res.* 22:268-270, 1987.
- Mitra, A. and Zhang, Z., *Plant Physiol* 106:977-81, 1994.
- Moe, T, *et al.* (1995). *Mashing Of Rice With Barley Malt Under Non-Conventional Process Conditions For Use In Food Processes. International Journal Of*
- 15 *food Science And Technology* 29:635-649.
- Moore, K (1978). *Rice-Barley Malt Combo Improves Functionality And Nutrition Of Meats, Breads, And Baked Goods. Food Product Development* 12(5): 74.
- Moreira-Ludewig, R. and Healy, C., *J. Pharm. and Tox. Meth.*, 27, 75-100, 1992.
- Motil K.J., *Curr Opin Pediatr* 12(5):469-76, 2000.
- 20 Murphy MS (1998). *Growth Factors And The Gastrointestinal Tract. Nutrition* 14(10):771-4.
- Nakajima H. *et al.*, *Plant Cell Reports*, 16, 674-679, 1997.
- Newburg, DS (1999). *Human Milk Glycoconjugates That Inhibit Pathogens. Curr Med Chem* 6(2):117-127.
- 25 Newburg, DS, *et al.* (1998). *Role Of Human-Milk Lactadherin In Protection Against Symptomatic Rotavirus Infection. Lancet* 351(1910):1160-4.
- NIH publication, American Academy of Allergy and Immunology Committee on adverse reaction to food and National Institute of Allergy and Infectious Diseases, NIH, Bethesda, 1984
- 30 Peeters, K., *et al.*, Production of antibodies and antibody fragments in plants. *Vaccine* 9:2756-61, 2001.
- Peterson JA, *et al.* (1998). *Glycoproteins Of The Human Milk Fat Globule In The Protection Of The Breast-Fed Infant Against Infections. Bio Neonate* 74(2):143-162.
- Piper, D.W. and Fenton, B.H. pH stability and activity curves of pepsin with
- 35 special reference to their clinical importance. *Gut* 6:506-508, 1965.

- Prosser, CG (1996). *Insulin-Like Growth Factors In Milk And Mammary Gland. J Mammary Gland Bio Neoplasia* (3):297-306.
- Raikhel N., *Plant Phys.* 100:1627-1632, 1992.
- Rey, M.W., *Nucleic Acid Res.* 18, 5288, 1990.
- 5 Romer *et al.*, *Biochem. Biophys. Res Commun.* 196:1414-1421, 1993.
- Rudloff, S. and Lönnerdal, B. Solubility and digestibility of milk proteins in infant formulas exposed to different heat treatments. *J. Pediatr. Gastroenterol. Nutr.* 15:25-33, 1992.
- Saariinen K.M. *et al.*, *Adv Exp Med Biol* 478:121-30, 2000.
- 10 Salmon V. *et al.*, *Protein Expression and purification* 9, 203-210, 1997.
- Salmon, V. *et al.*, *Protein Expr Purif* 13:127-135, 1998.
- Samaranayake Y.H. *et al.*, *Apmis*, 105, 875-883, 1997.
- Sambrook J. *et al.*, *MOLECULAR CLONING: A LABORATORY MANUAL* (Second Edition), Cold Spring Harbor Press, Plainview, N.Y, 1989.
- 15 Satue-Gracia M.T. *et al.*, *J Agric Food Chem* 48(10):4984-90, 2000.
- Schütte H. and Kula M.R., *Biotechnology and Applied Biochemistry*, 12, 599-620, 1990.
- Sfat, M, *et al* (1981). *Malts And Malting*. In: Kirk-Othmer Encyclopedia Of Chemical Technology (John Wiley & Sons, New York, 3rd edition), volume 14, pp. 810-20 823.
- Shah *et al.*, *Science* 233:478-481, 1986.
- Shin , K, *et al.* (2000). *PCR Cloning And Baculovirus Expression Of Human Lactoperoxidase And Myeloperoxidase*. Biochem Biophys Res Commun 27(13):831-836.
- 25 Shugar D., *Biochim. Biophys. Acta*, 8, 680-68, 1952.
- Symbicon AB (1995). *Human Milk Kappa-Casein And Inhibition Of Helicobacter pylori Adhesion To Human Gastric Mucosa*. J Pediatr Gastroenterol Nutr 21(3):288-296.
- Takai I. *et al.*, *J. Chrom. B, Biomedical Applications*, 685, 21-25, 1996.
- 30 Tsuchiya K. *et al.*, *Applied Microbiology and Biotechnology*, 38, 109-114, 1992.
- Von Heijne *et al.*, *Plant Mol. Biol. Rep.* 9:104-126, 1991.
- Wang C.S. and Kloer H.U., *Anal. Biochem.*, 139, 224-227, 1984.
- Wang C. *et al.*, *Comp. Biochem. Physiol.* 78B, 575-580, 1984.
- Ward P.P. *et al.*, *Bio/technology* 10, 784-789, 1992.
- 35 Ye X. *et al.*, *Science* 287, 303-305, 2000.

Yoshimura K. *et al.*, *Biochem. Biophys Res Com*, 150, 794-801, 1988.

Background of the Invention

The profitability of livestock and poultry producers is dependent upon the efficiency and rate that nutrients in feed are converted into salable products such as meat, milk and eggs. An animal's maximum growth rate and efficiency of conversion of feed into products (feed conversion efficiency) are set by its genetic potential.

Environmental factors grouped under the term "stress" prevent the expression of an animal's genetic potential. A common form of stress encountered by animals is that of infectious challenge. Animals are exposed to millions of potential pathogens daily, through contact, ingestion, and inhalation. Infectious challenge may or may not result in clinical disease, depending upon the pathogenicity of the challenging microorganism and the immunocompetence of the animal. Regardless of the outcome, a stress response is indicated by decreased growth performance and feed conversion efficiency when the immune system is stimulated.

Veterinarians, like their medical doctor counterparts, occasionally prescribe the therapeutic use of antibiotics in order to protect the health of animals. The animal industries also rely on the chronic, low level (sub-therapeutic) feeding of antibiotics to prevent disease and mitigate the economic impact of infection. Approximately 100% of chicken and turkeys, 90% of swine and veal calves, and 60% of cattle raised in the United States receive diets that continuously contain antibiotics. These antibiotics are fed prophylactically with the goal of preventing infectious disease before it can be established. The use of antibiotics at sub-therapeutic levels (ST antibiotics) permits the large-scale husbandry of animals at very high population densities, with minimal labor costs and results in cost effective and wholesome production of foods. Modern animal farming in the United States is virtually dependent upon the use of antibiotics throughout the production cycle.

In 1995, ST antibiotics used by the animal production industries accounted for \$3.3 billion in sales. One study estimated that the use of sub-therapeutic antibiotics alone, saved the US swine industry \$2 billion in production costs annually. Sub-therapeutic levels of antibiotics in the diet or water allow an increased rate of growth and feed conversion efficiency, and increase the general state of health of fish, chickens, pigs, sheep, and cattle. Regulatory statutes restrict ST antibiotics to those that are poorly absorbed from the digestive tract and consequently do not contaminate meat, milk, and eggs. For this reason, the mode of action of ST antibiotics is limited to effects

within the digestive tract. By definition, ST antibiotics exert their beneficial effects by suppressing the growth of microorganisms. Antibiotics lack efficacy in germ-free or highly sanitized environments and do not require the presence of clinically identifiable disease or pathogenic agents.

5 The behavioral and metabolic changes that occur during an immune response to pathogen challenge include anorexia, fever, decreased accretion of skeletal muscle, synthesis of acute phase proteins, increased use of amino acids as an energy source, and decreased use of fat as an energy source. These changes form the basis for impaired growth, poor feed utilization, and altered nutrition requirements in growing
10 chickens and pigs challenged by non-infectious pathogens.

 According to a recent National Academy of Sciences Report, antibiotics "enhance growth and production performance because an animal can reduce that portion of the nutrition requirement associated with fighting subclinical disease and bolstering health defense processes, thereby enhancing the portion of nutrients
15 available for growth and production."

 However, the sub-therapeutic use of antibiotics in animals has also been shown to have a negative impact on human health. This is due to the emergence in food animals of zoonotic microorganisms that are resistant to antibiotics. Furthermore, the enlarged pool of resistance genes created by ST-antibiotics is thought to result in
20 decreased therapeutic efficacy of antibiotics used for treatment of a variety of human infections. For this reason, the use of antibiotics by animal production industries has come under intense scrutiny by the Food and Drug Administration, the Institute of Medicine, and the World Health Organization. All of these regulatory bodies have called for decreased use of ST-antibiotics by the animal production industries. Further, the
25 potential threat to human health due to ST antibiotic use in animal production has prompted several European countries (Sweden, Denmark, Switzerland) to completely ban their use. In July of 1999, the European Union banned the sub-therapeutic use of most antibiotics by all of its member countries. Australia is considering a similar ban. A recent report by the National Academy of Sciences in the United States examining this
30 issue does not call for a total ban but argues for the development of alternatives so that ST-antibiotic use in animals can be markedly reduced.

Summary of the Invention

 Accordingly, it is an object of the invention to provide an animal feed or feed
35 supplement containing a transgenic plant-produced heterologous, anti-microbial protein

which when fed to production animals results in improved feed efficiency, and methods for using and producing the improved animal feed.

In one aspect, the invention includes an improved feed for production animals (poultry and hoofed farm animals), comprising one or more plant-derived feed

5 ingredients, substantially unsupplemented with small-molecule antibiotics, and, as an additive, a seed composition containing a flour, extract, or malt obtained from mature monocot seeds and one or more seed-produced heterologous, anti-microbial proteins in substantially unpurified form.

In one embodiment, the one or more seed-produced anti-microbial protein(s) present in
10 the seed composition are an anti-microbial proteins, including the anti-microbial milk proteins such as lactoferrin, lysozyme, lactoferricin, lactohedrin, kappa-casein, haptocorrin, or lactoperoxidase, a milk protein like alpha-1-antitrypsin that may function as an antimicrobial by inhibiting proteolysis of other anti-microbial proteins, non-milk anti-microbial proteins and acute-phase proteins, e.g., proteins that are produced
15 normally in production animals in response to infection, and small anti-microbial proteins. Exemplary anti-microbial proteins are lysozyme and lactoferrin, where lysozyme is preferably present in an amount between about 0.05 and 0.5 grams protein/kg feed, and lactoferrin, in an amount between 0.2 to 2 grams/protein/kg feed.
17.

20 In another embodiment, the one or more seed-produced anti-microbial protein(s) present in the food are acute-phase, non-milk proteins selected from the group consisting of C-reactive protein, serum amyloid A; ferritin, haptoglobin, seromucoids, ceruloplasmin, 15-keto-13,14-dihydro-prostaglandin F2 alpha, fibrinogen, alpha-1-acid glycoprotein, mannose binding protein, lipopolysaccharide binding protein, alpha-2
25 macroglobulin and defensins.

In still another embodiment, the the one or more seed-produced anti-microbial protein(s) present in the food are antimicrobial peptides selected from the group consisting of cecropin, magainin, defensins, tachyplesin, parasin I, buforin I, PMAP-23, moronecidin, anoplin, gambicin, and SAMP-29.

30 In another embodiment, the one or more seed-produced anti-microbial protein(s) present in the food are antimicrobial pproteins selected from the group consisting: CAP37, granulysin, secretory leukocyte protease inhibitor, CAP18, ubiquickidin, bovine antimicrobial protein-1, Ace-AMP1, tachyplesin, big defensin, Ac-AMP2, Ah-AMP1, and CAP18.

35 The seed composition may be prepared as follows:

(a) the flour is prepared by milling mature monocot seeds,

(b) the extract is prepared by suspending milled flour in a buffered aqueous medium; and

(c) the malt is prepared by (i) steeping barley seeds to a desired water content,

5 (ii) germinating the stepped barley, (iii) drying the germinated seeds, under conditions effective to stop germination, (iv) crushing the dried seeds, (v) optionally, adding crushed seeds from a non-barley monocot plant, (vi) forming a mixture of crushed seeds in water, and (vii) malting the crushed seed mixture until a desired malt is achieved, where at least one of the barley or non-barley monocot seeds contain such anti-
10 microbial protein(s). Step (v) in the malt-producing step may include adding to the crushed dried barley seeds, mature rice transgenic seeds that produce an anti-microbial protein.

In another aspect, the invention includes an improvement over existing methods for achieving high growth rates in production animals, particularly methods in which a
15 production animal is fed a feed supplemented with subclinical (sub-therapeutic or ST) levels of one or more small-molecule antibiotics. The improvement comprises replacing the small-molecule antibiotic(s) in the feed with a seed composition containing a flour, extract, or malt obtained from mature monocot seeds and one or more seed-produced heterologous, anti-microbial proteins in substantially unpurified form.

20 In a related aspect, the invention includes a method of producing a feed for production animals. The method includes the steps of first obtaining a monocot plant that has been stably transformed with a first chimeric gene having (i) a transcriptional regulatory region from a monocot gene having a seed maturation-specific promoter, (ii) operably linked to said transcriptional regulatory region, a leader/targeting DNA
25 sequence encoding a monocot seed-specific transit sequence capable of targeting a linked polypeptide to an endosperm-cell organelle, and (iii) a protein-coding sequence encoding a heterologous, anti-microbial normally.

The transformed plant is cultivated under seed-maturation conditions, and the mature seeds are harvested, then extracted to yield a flour, extract, or malt composition
30 containing the anti-microbial protein in substantially unpurified form. The seed composition is added to an animal feed that is substantially free of small-molecule antibiotics, i.e., has no or significantly reduced amounts of small-molecule antibiotics added.

In preparing the seed composition:

35 (a) the flour is prepared by milling mature monocot seeds,

(b) the extract is prepared by suspending milled flour in a buffered aqueous medium; and

(c) the malt is prepared by (i) steeping barley seeds to a desired water content, (ii) germinating the stepped barley, (iii) drying the germinated seeds, under conditions effective to stop germination, (iv) crushing the dried seeds, and (v) after mixing the crushed seeds with water, malting the crushed seed mixture until a desired malt is achieved.

Exemplary transcriptional regulatory regions in the chimeric gene are from the promoter of the group of genes: rice glutelins, rice globulins, oryzins, and prolamines, barley hordeins, wheat gliadins and glutenins, maize zeins and glutelins, oat glutelins, and sorghum kafirins, millet pennisetins, and rye secalins genes. Exemplary leader/targeting sequences are likewise from the group of genes selected from the group of rice glutelins, rice globulins oryzins, and prolamines, barley hordeins, wheat gliadins and glutenins, maize zeins and glutelins, oat glutelins, and sorghum kafirins, millet pennisetins, and rye secalins genes.

In one preferred embodiment, the transcriptional regulatory region in the chimeric gene is a rice glutelin Gt1 promoter, and the leader DNA sequence is a rice glutelin Gt1 signal sequence capable of targeting a linked polypeptide to a protein storage body. An exemplary glutelin Gt1 promoter and glutelin Gt1 signal sequence are included within the sequence identified by SEQ ID NO:15. In another preferred embodiment, the transcriptional regulatory region in the chimeric gene is a rice globulin Glb promoter, and the leader DNA sequence is a rice glutelin Gt1 signal sequence capable of targeting a linked polypeptide to a protein storage body. An exemplary globulin Glb promoter and glutelin Gt1 signal sequence are included within the sequence identified by SEQ ID NO:16.

The transformed monocot seed may further encode at least one transcription factors O2, PBF, and Reb, as exemplified by SEQ ID NOS: 31, 32, and 33, respectively, and preferably O2 and/or PBF.

The protein-coding sequence may be the coding sequence for a milk protein selected from the group consisting of lactoferrin, lysozyme, lactoferricin, lactohedrin, kappa-casein, haptocorrin, lactoperoxidase, alpha-1-antitrypsin, and immunoglobulins, preferably a sequence which has been codon-optimized for expression in monocots. Exemplary codon-optimized sequences for human milk proteins are represented by SEQ ID NOS: 1, 3, and 7-14. Exemplary coding sequences for acute-phase, non-milk proteins are identified by SEQ ID NOS: 36, and 46-56. Exemplary coding sequences for

antimicrobial peptides are identified by SEQ ID NOS: 34-68, 40-41, and 43. Exemplary sequences for other anti-microbial proteins are identified by SEQ ID NOS: 37, 45, and 57-59.

The plant may be further stably transformed with a second chimeric gene having
5 (i) a transcriptional regulatory region from a monocot gene having a seed maturation-specific promoter, (ii) operably linked to said transcriptional regulatory region, a transit DNA sequence encoding a monocot seed-specific transit sequence capable of targeting a linked polypeptide to an endosperm-cell organelle, and (iii) a protein-coding sequence encoding a second heterologous, anti-microbial protein.

10 In still another aspect, the invention includes an improved feed for production animals (poultry and hoofed farm animals), comprising one or more plant-derived feed ingredients, substantially unsupplemented with small-molecule antibiotics, and, as an additive, a seed composition containing one or more seed-produced heterologous, anti-microbial proteins in substantially purified form.

15 These and other objects and features of the invention will become more fully apparent when the following detailed description of the invention is read in conjunction with the accompanying drawings.

Brief Description of the Figures

20 Figure 1 is a map of the pAPI159 expression construct that contains the human lysozyme coding sequence under the control of a Gt1 promoter and Gt1 signal sequence.

Figure 2 shows the results of Western blot analysis for the expression of recombinant human lysozyme in various tissues of rice plants, where lanes 1 and 15 are a human milk lysozyme standard; lane 2 is a broad range molecular weight marker from Sigma; lanes 3 and
25 4 represent mature seed tissue extracts; lanes 5 and 6 represent germinated seed extracts; lanes 7 and 8 represent root tissue extracts; lanes 9 and 10 represent extracts from young root tissue; lanes 11 and 12 represent leaf extracts; and lanes 13 and 14 represent extracts from young leaf; from untransformed ("U") or transgenic ("T") plants, respectively. The total loading protein amount was 40 µg per lane.

30 Figure 3 shows the effect of incubating recombinant human lysozyme from transgenic rice seed, a human lysozyme standard (30 µg/ml), a control (20 mM sodium phosphate, pH 7.0, 5 mM EDTA) or an untransformed rice extract on the growth of *E.coli* strain JM109. At the end of the incubation (for the time indicated), an aliquot of the mixture was plated on LB plates and colony forming units per ml (CFU/ml) was calculated.

Figure 4 is a graph showing the specific activity of lysozyme, as determined by incubating an identical concentration of a human lysozyme standard, human lysozyme from transgenic rice (plant) and lysozyme from chicken egg white with a standard amount of *M. luteus*, followed by evaluation of the reduction in the turbidity due to the activity of lysozyme over five minutes.

Figure 5A: Thermal stability of human lysozyme ("Hlys") and recombinant human lysozyme from transgenic rice ("rHLys"). Lysozyme was dissolved at 100 µg/ml in PBS. The mixtures were subjected to different temperatures for different lengths of time. At the end of each heat treatment, the remaining lysozyme activity was assessed by activity assay. Figure 5B: pH stability of Hlys and rHlys. Lysozyme was dissolved in different buffers at 100 µg/ml. The mixture was incubated at 37°C for 30 min. The lysozyme activity was determined by activity assay.

Figure 6 presents the results of an analysis of lysozyme expression in transgenic rice grains over several generations. Proteins from 1 g of brown rice flour were extracted with 40 ml of extraction buffer containing 0.35 M NaCl in PBS. Extraction was conducted at room temperature for 1 h with shaking. Homogenate was centrifuged at 14,000 rpm for 15 min at 4°C. Protein supernatant was removed and diluted as needed for lysozyme turbidimetric activity assay. Extraction was repeated three times and standard deviation was shown as an error bar. Lysozyme yield was expressed as percentage of total soluble protein (%TSP).

Figure 7 is a restriction map of the pAPI164 plasmid that contains the human lactoferrin coding sequence under the control of a rice glutelin (Gt1) promoter, aGt1 signal peptide, and a nopaline synthase (NOS) terminator/polyadenylation site.

Figure 8 shows the results of a SDS-PAGE analysis for human lactoferrin stained with Coomassie blue, where lane 1 is the molecular weight marker; lanes 2 - 5 are purified human derived lactoferrin (Sigma, USA); lanes 6 - 10 are single seed extracts from homozygous transgenic lines and lane 11 is a seed extract from non-transformed TP-309.

Figure 9 shows the results of a Western blot analysis of various tissues of the transgenic rice plants, demonstrating the tissue specificity of rLF expression. Lane 1 is the molecular weight marker; lane 2 is human lactoferrin (Sigma, USA); lane 3 is an extract from leaf; lane 4 is an extract from sheath; lane 5 is an extract from root; lane 6 is an extract from seed and lane 7 is an extract from 5-day germinated seeds.

Figure 10 is a bar diagram illustrating the bactericidal effect of native human lactoferrin ("nHLF") and purified recombinant human lactoferrin produced by transgenic rice ("rHLF") on growth of *E. coli* (EPEC) after pepsin/pancreatic treatment.

Figure 11 is a graph illustrating pH-dependent iron release by native human
5 lactoferrin ("nHLF") and purified recombinant human lactoferrin produced by transgenic rice seeds ("rHLF").

Figure 12 shows the binding and uptake of HLF to Caco-2 cells after *in vitro* digestion. Figure 12 A shows the determination of Dissociation constant. Figure 12B shows the number of binding sites for HLF on Caco-2 cells. Figure 12C shows the total uptake of HLF and Fe to
10 Caco-2 cells within 24 h. Figure 12D shows degradation of HLF after uptake into Caco cells determined by the amount of free ^{125}I in the cell fractions.

Figure 13 shows three AAT plasmids: pAPI255 containing Glb promoter, Glb signal peptide, codon-optimized AAT gene, Nos terminator and ampicillin resistance gene; pAPI250 containing Gt1 promoter, Gt1 signal peptide, codon-optimized AAT gene,
15 Nos terminator and ampicillin resistance gene; and pAPI282 containing Bx7 promoter, Bx7 signal peptide, codon-optimized AAT gene, Nos terminator and ampicillin resistance gene.

Figure 14 shows Coomassie brilliant blue staining of aqueous phase extraction of transgenic rice cells expressing human AAT. Both untransformed and transgenic rice
20 grains were ground with PBS. The resulting extract was spun at 14,000 rpm at 4° C for 10 min. Supernatant was collected and loaded onto a precast SDS-PAGE gel.

Figure 15 shows Western blot analysis of recombinant human AAT from transgenic rice grains. The extract from transgenic rice grain was separated by SDS-PAGE gel and then blotted onto a filter. The identification of AAT in rice grain was
25 carried out by anti-AAT antibody by Western analysis.

Figures 16A-B shows Coomassie staining (Figure 16A) and western blot analysis (Figure 16B) of protein from transgenic rice grains expressing AAT. The activity of rAAT was demonstrated by a band shift assay. AAT samples from different sources were incubated with equal moles of porcine pancreatic elastase (PPE) at 37°C
30 for 15 min. Negative control for band shift assay was prepared with the AAT samples incubated with equal volume of PPE added. Lane M is molecular weight markers. Lane 1a is purified AAT from human plasma. Lane 1b is purified AAT from human plasma + PPE. Lane 2a is protein extract containing AAT from transgenic rice seed; Lane 2b is protein extract containing AAT from transgenic rice seed + PPE. Lane 3a is
35 untransformed seed extract. Lane 3b is untransformed seed extract + PPE. A shifted

band was shown in lane 1b, 2b and 3b in Figure 16A. The shifted band was confirmed to contain AAT entity by Western blot in Figure 16B.

Figure 17A-C are schematic representations of 3 plasmids containing the Reb coding sequence under the control of 3 different promoters. Figure 17A shows the globulin promoter (Glb), with the Reb gene and the Reb terminator. Figure 17B shows the actin promoter (Act), with the Reb gene and the Reb terminator. Figure 17C shows the native Reb promoter, with the Reb gene and the Reb terminator.

Figures 18A-B are schematic depictions of 2 plasmids which contain different transcription factor coding sequences under the control of the rice endosperm-specific glutelin promoter (Gt-1). Figure 18A shows plasmid pGT1-BPBF (API286) containing the Gt1 promoter, barley prolamins box binding factor (BPBF), Nos terminator and kanamycin resistance gene. Figure 18B shows pGT1-PBF (API285) containing the Gt1 promoter, the maize prolamins box binding factor (PBF), Nos terminator and kanamycin resistance gene.

Figure 19 illustrates the results of an analysis for the expression of recombinant human lysozyme in mature seed of T₀ transgenic plants derived from progenitor cells transformed with constructs containing the human lysozyme gene expressed under the control of the Glb promoter and the Reb gene expressed under the control of its own promoter ("Native-Reb"). Seeds of 30 plants containing the Reb and lysozyme genes and seeds from 17 plants containing only the lysozyme gene were analyzed for lysozyme, with twenty individual seeds of each plant analyzed.

Figure 20 is a comparison of the codon-optimized epidermal growth factor sequence ("Egfactor") with a native epidermal growth factor sequence ("Native Gene"), aligned to show 53 codons in the mature sequences, with 27 (51%) codon changes and 30 (19%) nucleotides changes.

Figure 21 is a restriction map of the 4,143 bp plasmid, API270 (pGlb-EFG v2.1), showing an expression cassette for epidermal growth factor ("EGF"), and containing a Glb promoter, a Glb signal peptide, codon optimized EGF, a Nos terminator and an ampicillin resistance selectable marker.

Figure 22 is a restriction map of the 3877 bp plasmid, API303 (pGt1-EGF v2.1), showing an expression cassette for epidermal growth factor (EGF), and containing a rice Gt1 promoter, a Gt1 signal peptide, codon optimized EGF, a Nos terminator and an ampicillin resistance selectable marker.

Figure 23 is a Western blot analysis of recombinant human EFG ("rhEGF") in transgenic rice seed. Lane 1 shows a broad range of molecular weight markers. Lane

2 shows rhEGF expressed in yeast, loaded at 125 ng. Lanes 2 to 6 show rhEGF expressed from different transgenic rice seeds. Lane 7 is from seeds of control untransformed TP 309.

5 Figure 24 is a comparison of the codon-optimized insulin-like growth factor I sequence ("Insgfact") with a native human insulin-like growth factor I sequence ("native Gene"), aligned to show 70 codons in the mature sequences, with 40 (57%) codon changes and 47 (22%) nucleotides changes.

10 Figure 25 is a restriction map of the 3928 bp plasmid, API304 (pGt1-IGF v2.1), showing an expression cassette for insulin-like growth factor I ("IGF"), and containing a rice Gt1 promoter, a Gt1 signal peptide, codon optimized IGF, a Nos terminator and an ampicillin resistance selectable marker.

15 Figure 26 is a restriction map of the 4194 bp plasmid, API271 (pGlb-IGF v2.1), showing an expression cassette for insulin-like growth factor I ("IGF"), and containing a Glb promoter, a Glb signal peptide, codon optimized IGF, a Nos terminator and an ampicillin resistance selectable marker.

20 Figure 27 is a Western blot analysis of recombinant human IGF-I ("rhIGF") expressed in transgenic rice seeds. Lane 1 shows a broad range of molecular weight markers. Lane 2 shows rhIGF expressed in yeast, loaded at 1 μ g. Lanes 3-9 show rhIGF from different transgenic seeds. Lane 10 is from seeds of control untransformed TP 309.

Figure 28 is a restriction map of the 5250 bp plasmid, API321 (pGlb-gt1sig-Haptocorrin v 2.1), showing an expression cassette for haptocorrin, and containing a Glb promoter, a Gt1 signal peptide, codon optimized haptocorrin, a Nos terminator and an ampicillin resistance selectable marker.

25 Figure 29 is a restriction map of the 4948 bp plasmid, API320 (pGt1-Haptocorrin v 2.1), showing an expression cassette for haptocorrin, and containing a Gt1 promoter, a Gt1 signal peptide, codon optimized haptocorrin, a Nos terminator and an ampicillin resistance selectable marker.

30 Figure 30 is a restriction map of the 4468 bp plasmid, API292 (pGlb-kcasein v2.1), showing an expression cassette for kappa-casein ("k-casein"), and containing a Glb promoter, a Glb signal peptide, a k-casein gene, a Nos terminator and an ampicillin resistance selectable marker.

35 Figure 31 is a restriction map of the 4204 bp plasmid, API297 (pGT1-kaapa-Casein v2.1), showing an expression cassette for kappa-casein, and containing a Gt1 promoter, a Gt1 signal peptide, mature kappa-casein polypeptide encoding gene, a Nos

terminator and an ampicillin resistance selectable marker.

Figure 32 is a restriction map of the 4834 bp plasmid, API420 (pGt1-LAD), showing an expression cassette for lactahedrin, and containing a Gt1 signal peptide, lactohedrin gene, a Nos terminator and a kanamycin resistance selectable marker.

Figure 33 is a restriction map of the 5638 bp plasmid, API418 (pGT1-LPO-S), showing an expression cassette for lactoperoxidase (minus the propeptide), and containing a Gt1 promoter, a Gt1 signal peptide, lactoperoxidase gene without the propeptide, a Nos terminator and a kanamycin resistance selectable marker.

Figure 34 is a restriction map of the 5801 bp plasmid, API416 (pGt1-lactoperoxidase), showing an expression cassette for codon optimized human lactoperoxidase, and containing a rice Gt1 promoter, a Gt1 signal peptide, codon optimized lactoperoxidase, a Nos terminator and a kanamycin resistance selectable marker.

Figure 35 is a restriction map of the 4408 bp plasmid, API230 (pBX7-Lysozyme v2.1.1), showing an expression cassette for codon optimized lysozyme, and containing a BX-7 promoter, a Gt1 signal peptide, codon optimized lysozyme gene, a Nos terminator and an ampicillin resistance selectable marker.

Figures 36A-B represent schematic diagrams of the map of 2 plasmids, API254 (Figure 36A) and API264 (Figure 36B) containing heterologous protein coding sequences under the control of the rice endosperm-specific globulin promoter (Glb), the Glb signal peptide, and Nos terminator. API254 contains the lactoferrin coding sequence, and API264 contains the human lysozyme coding sequence.

Figure 37 is a restriction map of the 4271 bp plasmid, API225, showing an expression cassette for codon optimized lysozyme, and containing a GT-3 promoter, a Gt1 signal peptide, codon optimized lysozyme, a Nos terminator and an ampicillin resistance selectable marker.

Figure 38 is a restriction map of the 4106 bp plasmid, API229, showing an expression cassette for codon optimized lysozyme, and containing a RP-6 promoter, a Gt1 signal peptide, codon optimized lysosyme, a Nos terminator and an ampicillin resistance selectable marker.

Figure 39 is a comparison of the expression of lysozyme under Gt1 or Glb promoter with Gt1 signal peptide or Glb signal peptide. Figure 50A is a schematic representation of plasmid API159 that contains Gt1 promoter, Gt1 signal peptide, a lysozyme gene and Nos terminator; plasmid API 228 that contains Glb promoter, Gt1

signal peptide, a lysozyme gene and Nos terminator; and plasmid API264 that contains Glb promoter, Glb signal peptide, a lysozyme gene and Nos terminator. Figure 39B shows the activities of lysozyme in lysozyme-positive seeds produced in transgenic rice plants transformed with API159, API228 and API264. The seeds from multiple lines of each construct were analyzed by the lysozyme activity assay. Individual seeds from each plant were analyzed. Seeds lacking detectable amounts of lysozyme were excluded. The activities of 20-lysozyme-positive seeds per plant, including both hemizygous and homozygous seeds were averaged. The average activities were plotted on the chart.

Figure 40 shows the expression time course of human lysozyme during endosperm development in transgenic line. Ten spikelets were harvested at 7, 14, 21, 28, 35, 42 and 49 days after pollination ("DAP") and analyzed by the lysozyme activity assay. The dark bars were from 159-1-53-16-1. The light bars were from 264-1-92-6-1.

Figure 41 is a bar graph comparing the level of lysozyme expression in transgenic T1 rice seeds under 7 different promoters: Gt1, Glb, Glub-2, Bx7, Gt3, Glub-1 and Rp6. All constructs contained a Gt1 signal peptide.

Detailed Description of the Invention

I. Definitions

Unless otherwise indicated, all terms used herein have the meanings given below, and are generally consistent with same meaning that the terms have to those skilled in the art of the present invention. Practitioners are particularly directed to Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (Second Edition), Cold Spring Harbor Press, Plainview, N.Y. and Ausubel FM *et al.* (1993) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, N.Y., for definitions and terms of the art. It is to be understood that this invention is not limited to the particular methodology, protocols, and reagents described, as these may vary.

All publications cited herein are expressly incorporated herein by reference for the purpose of describing and disclosing compositions and methodologies that might be used in connection with the invention.

The term "polypeptide" refers to a biopolymer compound made up of a single chain of amino acid residues linked by peptide bonds. The term "protein" as used herein may be synonymous with the term "polypeptide" or "peptide" or may refer, in addition, to a complex of two or more polypeptides.

The term "anti-bacterial protein" refers to a polypeptide that is has the ability to limit or diminish the intensity or duration of infection by a micro-organism, e.g., bacteria, virus, or fungal organism, when administered to the gut of a production animal.

Included within this group of proteins are those that are (i) "bacteriostatic protein,"

- 5 meaning the protein is capable of inhibiting the growth of, but not capable of killing bacteria, (ii) "bactericidal protein," meaning a protein capable of killing bacteria, (iii) anti-viral proteins, (iii) agents, such as alpha-antitrypsin, that act, at least in part, by reducing proteolysis of other anti-microbial proteins, (iv) acute phase proteins which are induced in production animals in response to infection, (v) probiotic proteins, and (vi) cationic
10 antimicrobial proteins, such as bactericidal permeability increasing protein, lactoferrin, transferrin, cathepsin G, cystatin, CAP18, and pepsinogen C. (See, e.g., Robert E. W. Hancock *et al.*, 2000.), recognizing that some anti-microbial proteins will be in two or more of these classes.

"Anti-microbial proteins" include, without limitation,

- 15 (i) anti-microbial milk proteins (either human or non-human) lactoferrin, lysozyme, lactoferricin, lactohedrin, kappa-casein, haptocorrin, lactoperoxidase, alpha-1-antitrypsin, and immunoglobulins, e.g., IgA,

- (ii) acute-phase proteins, such as C-reactive protein (CRP); lactoferrin; lysozyme; serum amyloid A (SAA); ferritin; haptoglobin (Hp); complements 2-9, in
20 particular complement-3; seromucoid; ceruloplasmin (Cp); 15-keto-13,14-dihydro-prostaglandin F2 alpha (PGFM); fibrinogen (Fb); alpha(1)-acid glycoprotein (AGP); alpha(1)-antitrypsin; mannose binding protein; lipopolysaccharide binding protein; alpha-2 macroglobulin and various defensins,

- (iii) antimicrobial peptides, such as cecropin, magainin, defensins, tachyplesin, parasin
25 I, buforin I, PMAP-23, moronecidin, anoplin, gambicin, and SAMP-29, and

- (iv) other anti-microbial protein(s), including CAP37, granulysin, secretory leukocyte protease inhibitor, CAP18, ubiquicidin, bovine antimicrobial protein-1, Ace-AMP1, tachyplesin, big defensin, Ac-AMP2, Ah-AMP1, and CAP18.

- The term "milk protein" or "proteins normally present in milk" refers to a protein or
30 biologically active fragments thereof, found in normal mammalian milk, e.g., human milk, including, without limitation, of lactoferrin, lysozyme, lactoferricin, EGF, IGF-I, lactohedrin, kappa-casein, haptocorrin, lactoperoxidase, alpha-1-antitrypsin, and immunoglobulins, and biologically active fragments thereof. The milk protein may be human milk protein, i.e., having the protein sequence of a normal human milk protein, or

active fragment thereof, or of another mammalian source, e.g., bovine or ovine, or of a non-mammalian source, e.g., avian source.

The term "active fragments of an anti-microbial protein" refers to a modified anti-microbial protein containing either amino acid substitutions, deletions or additions, or
5 fragment substitutions or deletions, but retain the anti-microbial activity of the native anti-microbial protein.

A "small-molecule antibiotic" refers to one a non-peptide antibiotic having a molecular weight typically less than 2 Kdaltons, typically of fungal origin, meaning produced by a strain of fungus, or being an analog of such a compound. Exemplary
10 small-molecule antibiotics that have been used in animal feed, at subclinical levels, include, as examples, bacitracin, roxarsone; penicillin G, amoxicillin, semduramicin, chlortetracycline, tetracycline, neomycin, salinomycin, and virginiamycin.

The term "vector" refers to a nucleic acid construct designed for transfer between different host cells. An "expression vector" refers to a vector that has the
15 ability to incorporate and express heterologous DNA fragments in a foreign cell. Many prokaryotic and eukaryotic expression vectors are commercially available. Selection of appropriate expression vectors is within the knowledge of those having skill in the art. Accordingly, an "expression cassette" or "expression vector" is a nucleic acid construct generated recombinantly or synthetically, with a series of specified nucleic acid
20 elements that permit transcription of a particular nucleic acid in a target cell. The recombinant expression cassette can be incorporated into a plasmid, chromosome, mitochondrial DNA, plastid DNA, virus, or nucleic acid fragment. Typically, the recombinant expression cassette portion of an expression vector includes, among other sequences, a nucleic acid sequence to be transcribed and a promoter.

25 ——— The term "plasmid" refers to a circular double-stranded (ds) DNA construct used as a cloning vector, and which forms an extrachromosomal self-replicating genetic element in many bacteria and some eukaryotes.

The term "selectable marker-encoding nucleotide sequence" refers to a nucleotide sequence capable of expression in plant cells and where expression of the
30 selectable marker confers to plant cells containing the expressed gene the ability to grow in the presence of a selective agent. As used herein, the term "Bar gene" refers to a nucleotide sequence encoding a phosphinothricin acetyltransferase enzyme that upon expression confers resistance to the herbicide glufosinate-ammonium ("Basta").

A "transcription regulatory region" or "promoter" refers to nucleic acid sequences
35 that influence and/or promote initiation of transcription. Promoters are typically

considered to include regulatory regions, such as enhancer or inducer elements. The promoter will generally be appropriate to the host cell in which the target gene is being expressed. The promoter, together with other transcriptional and translational regulatory nucleic acid sequences (also termed "control sequences"), is necessary to
5 express any given gene. In general, the transcriptional and translational regulatory sequences include, but are not limited to, promoter sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, and enhancer or activator sequences.

"Chimeric gene" or "heterologous nucleic acid construct", as defined herein
10 refers to a construct which has been introduced into a host and may include parts of different genes of exogenous or autologous origin, including regulatory elements. A chimeric gene construct for plant/seed transformation is typically composed of a transcriptional regulatory region (promoter) operably linked to a heterologous protein coding sequence, or, in a selectable marker heterologous nucleic acid construct, to a
15 selectable marker gene encoding a protein conferring antibiotic resistance to transformed plant cells. A typical chimeric gene of the present invention, includes a transcriptional regulatory region inducible during seed development, a protein coding sequence, and a terminator sequence. A chimeric gene construct may also include a second DNA sequence encoding a signal peptide if secretion of the target protein is
20 desired.

A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably
25 linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, "operably linked" elements, e.g., enhancers, do not have to be contiguous.
30 Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The term "gene" means the segment of DNA involved in producing a polypeptide chain, which may or may not include regions preceding and following the coding region,

e.g. 5' untranslated (5' UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons).

The term "sequence identity" means nucleic acid or amino acid sequence identity in two or more aligned sequences, aligned using a sequence alignment
5 program.

The term "% homology" is used interchangeably herein with the term "% identity" and refers to the level of nucleic acid or amino acid sequence identity between two or more aligned sequences, when aligned using a sequence alignment program. For example, 70% homology means the same thing as 70% sequence identity determined by
10 a defined algorithm, and accordingly a homologue of a given sequence has greater than 80% sequence identity over a length of the given sequence. Exemplary levels of sequence identity include, but are not limited to, 80, 85, 90 or 95% or more sequence identity to a given sequence, e.g., the coding sequence for lactoferrin, as described herein.

15 Exemplary computer programs which can be used to determine identity between two sequences include, but are not limited to, the suite of BLAST programs, e.g., BLASTN, BLASTX, and TBLASTX, BLASTP and TBLASTN, publicly available on the Internet at "www.ncbi.nih.gov/BLAST/". See, also, Altschul, S.F. *et al.*, 1990 and Altschul, S.F. *et al.*, 1997.

20 Sequence searches are typically carried out using the BLASTN program when evaluating a given nucleic acid sequence relative to nucleic acid sequences in the GenBank DNA Sequences and other public databases. The BLASTX program is preferred for searching nucleic acid sequences which have been translated in all reading frames against amino acid sequences in the GenBank Protein Sequences and
25 other public databases. Both BLASTN and BLASTX are run using default parameters of an open gap penalty of 11.0, and an extended gap penalty of 1.0, and utilize the BLOSUM-62 matrix. [See, Altschul, *et al.*, 1997.]

A preferred alignment of selected sequences in order to determine "% identity" between two or more sequences, is performed using for example, the CLUSTAL-W
30 program in MacVector version 6.5, operated with default parameters, including an open gap penalty of 10.0, an extended gap penalty of 0.1, and a BLOSUM 30 similarity matrix.

A nucleic acid sequence is considered to be "selectively hybridizable" to a reference nucleic acid sequence if the two sequences specifically hybridize to one
35 another under moderate to high stringency hybridization and wash conditions.

Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex or probe. For example, "maximum stringency" typically occurs at about $T_m - 5^\circ\text{C}$ (5° below the T_m of the probe); "high stringency" at about $5-10^\circ$ below the T_m ; "intermediate stringency" at about $10-20^\circ$ below the T_m of the probe; and "low stringency" at about $20-25^\circ$ below the T_m . Functionally, maximum stringency conditions may be used to identify sequences having strict identity or near-strict identity with the hybridization probe; while high stringency conditions are used to identify sequences having about 80% or more sequence identity with the probe.

Moderate and high stringency hybridization conditions are well known in the art (see, for example, Sambrook *et al*, 1989, Chapters 9 and 11, and in Ausubel *et al.*, 1993, expressly incorporated by reference herein). An example of high stringency conditions includes hybridization at about 42°C in 50% formamide, 5X SSC, 5X Denhardt's solution, 0.5% SDS and 100 $\mu\text{g/ml}$ denatured carrier DNA followed by washing two times in 2X SSC and 0.5% SDS at room temperature and two additional times in 0.1X SSC and 0.5% SDS at 42°C .

As used herein, "recombinant" includes reference to a cell or vector, that has been modified by the introduction of a heterologous nucleic acid sequence or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found in identical form within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all as a result of deliberate human intervention.

A plant cell, tissue, organ, or plant into which a heterologous nucleic acid construct comprising the coding sequence for an anti-microbial protein or peptide has been introduced is considered transformed, transfected, or transgenic. A transgenic or transformed cell or plant also includes progeny of the cell or plant and progeny produced from a breeding program employing such a transgenic plant as a parent in a cross and exhibiting an altered phenotype resulting from the presence of the coding sequence for an anti-microbial protein. Hence, a plant of the invention will include any plant which has a cell containing introduced nucleic acid sequences, regardless of whether the sequence was introduced into the plant directly through transformation means or introduced by generational transfer from a progenitor cell which originally received the construct by direct transformation.

The term "transgenic plant" refers to a plant that has incorporated exogenous nucleic acid sequences, *i.e.*, nucleic acid sequences which are not present in the native ("untransformed") plant or plant cell. Thus a plant having within its cells a heterologous

polynucleotide is referred to herein as a "transgenic plant". The heterologous polynucleotide can be either stably integrated into the genome, or can be extra-chromosomal. Preferably, the polynucleotide of the present invention is stably integrated into the genome such that the polynucleotide is passed on to successive generations. The term "transgenic" as used herein does not encompass the alteration of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or spontaneous mutation. "Transgenic" is used herein to include any cell, cell line, callus, tissue, plant part or plant, the genotype of which has been altered by the presence of heterologous nucleic acids including those transgenics initially so altered as well as those created by sexual crosses or asexual reproduction of the initial transgenics.

Terms "transformed", "stably transformed" or "transgenic" with reference to a plant cell means the plant cell has a non-native (heterologous) nucleic acid sequence integrated into its genome which is maintained through two or more generations.

The term "expression" with respect to a protein or peptide refers to the process by which the protein or peptide is produced based on the nucleic acid sequence of a gene. The process includes both transcription and translation. The term "expression" may also be used with respect to the generation of RNA from a DNA sequence.

The term "introduced" in the context of inserting a nucleic acid sequence into a cell, means "transfection", or "transformation" or "transduction" and includes the incorporation of a nucleic acid sequence into a eukaryotic or prokaryotic cell where the nucleic acid sequence may be incorporated into the genome of the cell (for example, chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (for example, transfected mRNA).

By "host cell" is meant a cell which contains a vector and supports the replication, and/or transcription or transcription and translation (expression) of the expression construct. Host cells for use in the present invention can be prokaryotic cells, such as *E. coli*, or eukaryotic cells such as yeast, plant, insect, amphibian, or mammalian cells. In general, host cells are monocotyledenous or dicotyledenous plant cells.

A "plant cell" refers to any cell derived from a plant, including undifferentiated tissue (e.g., callus) as well as plant seeds, pollen, progagules and embryos.

The term "mature plant" refers to a fully differentiated plant.

The terms "native" and "wild-type" relative to a given plant trait or phenotype refers to the form in which that trait or phenotype is found in the same variety of plant in nature.

5 The term "plant" includes reference to whole plants, plant organs (for example, leaves, stems, roots, etc.), seeds, and plant cells and progeny of same. Plant cell, as used herein includes, without limitation, seeds, suspension cultures, embryos, meristematic regions, callus tissue, leaves roots shoots, gametophytes, sporophytes, pollen, and microspores. The class of plants that can be used in the methods of the
10 present invention is generally as broad as the class of higher plants amenable to transformation techniques, including both monocotyledenous and dicotyledenous plants.

 The term "seed" is meant to encompass all seed components, including, for example, the coleoptile and leaves, radicle and coleorhiza, scutulum, starchy endosperm, aleurone layer, pericarp and/or testa, either during seed maturation and
15 seed germination.

 The term "seed in a form for use as a food or food supplement" includes, but is not limited to, seed fractions such as de-hulled whole seed, flour (seed that has been de-hulled by milling and ground into a powder) a seed protein extract (where the protein fraction of the flour has been separated from the carbohydrate fraction) and/or a purified
20 protein fraction derived from the transgenic grain.

 The term "purifying" is used interchangeably with the term "isolating" and generally refers to the separation of a particular component from other components of the environment in which it was found or produced. For example, purifying a recombinant protein from plant cells in which it was produced typically means subjecting
25 transgenic protein containing plant material to biochemical purification and/or column chromatography.

 The term "animal feed" refers to feed, commercial or otherwise, used as part or all of the diet of a production animal. The feed is typically a mixture or grains or grain-derived components and one or more nutritional supplements, e.g., amino acids, fat,
30 vitamins, minerals, and the like.

 "Monocot seed components" refers to carbohydrate, protein, and lipid components extractable from monocot seeds, typically mature monocot seeds.

 "Malted-seed components" refers to seed-derived components, predominantly carbohydrate components, after conversion of complex carbohydrates to malt sugars by

malting, i.e., treating with malting enzymes such as a barley amylase and glucanases, under conditions effective to conversion seed-derived carbohydrates to malt sugars.

"Substantially unpurified form", as applied to anti-microbial proteins in a seed extract means that the protein or proteins present in the extract are present in an amount less than 50% by weight, typically between 0.25 and 2.5 percent by weight.

"Seed maturation" or "grain development" refers to the period starting with fertilization in which metabolizable reserves, e.g., sugars, oligosaccharides, starch, phenolics, amino acids, and proteins, are deposited, with and without vacuole targeting, to various tissues in the seed (grain), e.g., endosperm, testa, aleurone layer, and scutellar epithelium, leading to grain enlargement, grain filling, and ending with grain desiccation.

"Inducible during seed maturation" refers to promoters which are turned on substantially (greater than 25%) during seed maturation.

"Heterologous DNA" or "foreign DNA" refers to DNA which has been introduced into plant cells from another source, or which is from a plant source, including the same plant source, but which is under the control of a promoter or terminator that does not normally regulate expression of the heterologous DNA.

"Heterologous protein" is a protein, including a polypeptide, encoded by a heterologous DNA.

A "signal/targeting/transport sequence" is an N- or C-terminal polypeptide sequence which is effective to localize the polypeptide or protein to which it is attached to a selected intracellular or extracellular region, including an intracellular vacuole or other protein storage body, chloroplast, mitochondria, or endoplasmic reticulum, or extracellular space or seed region, such as the endosperm, following secretion from the cell.

A "product" encoded by a DNA molecule includes, for example, RNA molecules and polypeptides.

A DNA sequence is "derived from" a gene if it corresponds in sequence to a segment or region of that gene. Segments of genes which may be derived from a gene include the promoter region, the 5' untranslated region, and the 3' untranslated region of the gene.

"Alpha-amylase" as used herein refers to an enzyme which principally breaks starch into dextrins.

"Beta-amylase" as used herein refers to an enzyme which converts starch and dextrins into maltose.

"Cereal adjuncts" as used herein refers to cereal grains, principally barley, wheat, rye, oats, maize, sorghum and rice, or processed whole or portions thereof, especially the starch fraction, which are added to the barley mash, which allows the barley enzymes to hydrolyze both the barley starch and the starch derived from the cereal adjunct. "Transgenic cereal adjuncts" as used herein refers to transgenic cereal grains, principally barley, wheat, rye, oats, maize, sorghum and rice, and which is expressing a recombinant molecule in a grain part, principally the endosperm (starch) layer.

"Conversion" as used herein refers to the process of starch hydrolysis, usually catalyzed by acid or enzyme action, which produces dextrose, maltose, and higher polysaccharides from starch.

"Diastatic enzyme (amylolytic)" as used herein refers to an enzyme capable of causing the hydrolysis of starch.

"Diastatic malt flour" as used herein refers to enzyme active flour milled from germinated (malted) barley.

"Diastatic malt syrup" as used herein refers to enzyme active liquid malt syrup (barley and cereal adjuncts).

"Dry diastatic malt" as used herein refers to a blend of diastatic malted barley flour, wheat flour and dextrose with standardized enzyme levels at 20degrees and 60 degrees Lintner.

"Dry nondiastatic malt" as used herein refers to spray dried form of liquid nondiastatic malt extract or syrup.

"Lintner" as used herein refers to a laboratory measurement of enzyme activity strength. The higher the value, the higher activity.

"Dried malt" as used herein refers to the dried grain resulting from controlled germination of cereal grains, usually barley, but other cereals can be malted as well.

"Malt extract" as used herein refers to a viscous concentrate of the water extract of dried malt.

"Maltodextrin" as used herein refers to a purified, concentrated aqueous solution of nutritive saccharides, obtained from edible starch, or the dried product derived from the solution. Maltodextrins have a dextrose equivalent of less than 20 and are considered 'non-sweet soluble solids'. They are usually marketed dry, but may be obtained as a concentrated solution. Maltodextrins are usually offered as 10 to 14 D.E. products or as 15 to 19 D.E. versions. Another maltodextrin, with a D.E. of about 5, is sometimes manufactured, but currently not used widely. Composition of maltodextrins

is roughly 65 to 80% higher saccharides, 4 to 9% pentasaccharides, 4 to 7% tetrasaccharides, and 5 to 9% trisaccharides. Traces of mono and disaccharides are present. They are usually used as bulking agents or viscosity builders, without sweetness.

5 "Malt syrup" as used herein refers to viscious concentrate of the water extract of dried 'malt' and other cereal grains.

"Malt" refers to a malt extract or malt syrup.

"Nondiastatic malt syrup" as used herein refers to liquid malt syrup (barley and cereal adjuncts) without enzyme activity.

10 "Transgenic malt extract" as used herein refers to a vicious concentrate of the water extract of dried malt which includes a recombinant protein, polypeptide and/or metabolite.

"Transgenic malt syrup" as used herein refers to vicious concentrate of the water extract of dried 'malt' and other cereal grains which includes a recombinant protein, polypeptide and/or metabolite.

15 The term "production animals" includes poultry, such as chickens, turkeys, squab, and ducks, and hoofed farm animals, such as cattle, sheep, goats, and pigs that are grown and harvested for human food consumption.

20 II. Animal Feed-State of the Art

As described above, animal feed typically contains sub-therapeutic levels of antibiotics, which have been found to increase weight gain in animals by about 3-15% over a given feed period. The antibiotics are believed to suppress the microbial flora, and hence reduce the immune responses in the animal to low-level infection.

25 Untreated, such immune responses result in the animal eating less and therefore gaining less weight. In other words, the antibiotics help keep the animal in a "hungry" healthy state.

It has been reported that a loss of about 20% animal weight results from filthy animal-growing conditions, and a reclamation of about 4-15% of the "lost" weight takes place when antibiotics are incorporated into feed.

30 The reactions of an animal's immune system to infection is both specific against the causal agent and nonspecific, yielding predictable metabolic changes termed the "acute-phase response" (Johnson, 1997). The acute phase proteins (APP) produced during this response play a major role in the inflammatory process in mammals. In human and veterinary medicine, determination of the plasma concentration of these

proteins gives valuable clinical information on the infection inflammation process (See, e.g., Gruys *et al.*, 1994.)

In general, the inflammatory response begins with release of interleukin-1 (IL-1) and tumor necrosis factor-alpha (TNF-alpha) by monocytes and macrophages that subsequently activates a complex cascade of other inflammatory mediators including IL-2, IL-6 and IL-8 (Dinarelli, 1994; Jensen *et al.*, 1998).

The various acute phase proteins rise in response to activation by pro-inflammatory cytokines (such as IL-1 and TNF-alpha) that are secreted into the circulation from sites of infection or inflammatory lesions. The infection may be subclinical and of bacterial, viral or other origin and still effect both growth and appetite in animals.

Data in chicks, rats, and mice implicate interleukin-1 (IL-1), tumor necrosis factor-alpha (TNF), and interleukin-6 (IL-6), as the primary monokines involved in the acute-phase response. The metabolic changes orchestrated by these cytokines represent a homeostatic response that alter the partitioning of dietary nutrients away from the growth and skeletal muscle accretion in favor of metabolic processes that support their immune response and disease resistance.

The release of cytokines leads to both the stimulation and inhibition of protein synthesis. Exemplary positive acute phase proteins, include C-reactive protein (CRP); lactoferrin; lysozyme; serum amyloid A (SAA); ferritin; haptoglobin (Hp); complements 2-9, in particular complement-3; seromucoid; ceruloplasmin (Cp); 15-keto-13,14-dihydro-prostaglandin F2 alpha (PGFM); fibrinogen (Fb); alpha(1)-acid glycoprotein (AGP); alpha(1)-antitrypsin; mannose binding protein; lipopolysaccharide binding protein; alpha-2 macroglobulin and various defensins. (See, e.g., Laurell, 1985 and Thompson *et al.*, 1992.) The circulating concentration of these acute phase proteins are biochemical markers of inflammation. Cytokines can also inhibit the synthesis of proteins (negative acute phase proteins), e.g., serum albumin and transferrin. (See, e.g., P. D. Eckersall, Ed., IN: TEXTBOOK OF THE JAPANESE SOCIETY OF VETERINARY CLINICAL PATHOLOGY 10-21, 1999; Regassa *et al.*, 1999.)

Acute phase proteins that are reportedly produced by the intestinal epithelium include, but are not limited to, serum amyloid a, complement-3, mannose binding protein and various defensins.

In the past decade, the role of anti-microbial proteins has become increasingly apparent and there is a growing body of evidence that their role in defense against microbes may be as important to the host as antibodies, immune cells, and phagocytes.

Analysis of serum acute phase proteins such as SAA, Hp and AGP in bovine serum or plasma suggests a correlation to a number of disease states and may be used by meat inspectors to identify animals with current infection. (See, e.g., Conner *et al.*, 1989; Wittum *et al.*, 1996; Hofner *et al.*, 1994; Horadagoda *et al.*, 1994; Hirvonen *et al.*, 1996; Horadagoda *et al.*, 1999; Salonen *et al.*, 1996; and Saini *et al.*, 1998.)

Cationic anti-microbial peptides are also involved in the response to infection in all mammals and may be constitutively expressed or induced by infectious organisms or their products. A number of antimicrobial peptides show activity against a broad range of bacterial strains, including antibiotic-resistant isolates. Although the mechanism is not part of the invention, the antimicrobial peptides prevent cytokine induction by bacterial products in tissue culture and human blood, and they block the onset of sepsis in mouse models of endotoxemia. Exemplary antimicrobial peptides include, but are not limited to, animal cationic proteins such as bactericidal permeability increasing protein, lactoferrin, transferrin, cathepsin G, cystatin, CAP18, pepsinogen C, ribosomal protein S30, etc. (See, e.g., Robert E. W. Hancock *et al.*, 2000.)

Chicks raised in environments with poor sanitation have small intestines that possess high numbers of leukocytes in the villi and thickened lamina propria with larger Peyers patches. Chicks raised with poor sanitation also have markedly higher levels of circulation IL-1 than chicks raised with excellent sanitation. Presumably, the high burden of microbes, dust and dander chronically stimulate the immune system and induce the release of cytokines such as IL-1. As described above, high circulating levels of inflammatory cytokines result in slower growth. Feeding antibiotics to chicks in a dirty environment decreases the amount of circulation IL-1 to levels more similar to chicks raised in the clean environment. Feeding antibiotics result in little or no improvements in growth rate or changes in circulating IL-1 levels in clean environments. Thus, one of the mechanisms by which antibiotics improve growth rates is to decrease the number and severity of bacterial interactions with the animal. By decreasing the number of bacteria-host interactions, antibiotics decrease the degree of activity of the immune system and consequently decrease the levels of IL-1 and other monokines. This normalizes the metabolic rate and results in poor feed conversion efficiency and less muscular animals.

The concerns about antibiotics in animal feed, however, are well known, including such issues as the development of antibiotic-resistant microbes, requiring new and higher levels of antibiotics. Of perhaps greater concern is the threat of antibiotic

resistant genes being spread in the environment, potentially spreading even to human intestinal flora and/or human bacterial pathogens.

It has been proposed that acute phase proteins be used as markers for clinical and subclinical disease in production animals in order to identify conditions for optimal growth. Given the public pressure to reduce the use of ST antibiotics as growth promotants, there is a need to develop alternative strategies to maintain the health and growth of production animals.

III. Compositions Containing Anti-Microbial Proteins

The present invention provides animal feed compositions (also termed "improved animal feed compositions") containing one or more anti-microbial proteins, and methods of making such compositions. In practicing the invention, an anti-microbial protein, e.g., an acute-phase milk anti-microbial protein, is stably expressed in transgenic monocot seeds, the transgenic seeds are processed and added to animal feed. Accordingly, it is an object of the invention to provide animal feed compositions comprising anti-microbial proteins that allow for enhanced animal growth and production performance, without the side effects that result from long-term use of ST levels of small-molecule antibiotics.

By the present the invention a method is provided for maintaining healthy microflora in production animals by supplementing animal feed with transgenic grain or other plant material comprising one or more anti-microbial proteins. Exemplary anti-microbial proteins are the milk acute-phase anti-microbial proteins such as lysozyme and/or lactoferrin, particularly human lysozyme and/or lactoferrin.

Although the mechanism is not part of the invention, when fed to production animals the anti-microbial proteins described herein, are able to regulate the microflora of an animal in such a fashion that the animal avoids the typical low-level immune responses to subclinical infection. Accordingly, the supplemented animal feed compositions of the invention may be fed to production animals in order to regulate the animal gut microflora, thereby contributing to increased feed efficiency, and increased weight gain in the animals.

The invention is based on the expression at high expression levels of one or more anti-microbial proteins, exemplified by human lactoferrin (hLF) and human lysozyme, among others, under the control of a seed specific promoter in monocot seeds. The description provided herein includes a comparison of anti-microbial proteins produced by transgenic plants to the native form of the same protein, information on the

stability of the recombinant protein and the advantages of using grain containing such anti-microbial proteins in animal feed products.

The invention relies on the use of heterologous nucleic acid constructs which include the coding sequence for one or more anti-microbial proteins. Exemplary anti-microbial proteins, lysozyme and lactoferrin, are an integral part of the immune system of multicellular animals. They are found in epithelial secretions (tears, mucous, gastric juice) and blood plasma of mammals, birds, reptiles, amphibia, and a variety of invertebrates. They are also enriched in mammalian milk and avian eggs, where they serve as primary antimicrobial proteins. Furthermore, lysozyme is a major component of the secretory granules of neutrophils and macrophages and is released at the site of infection in the earliest stages of the immune response, and lactoferrin is found at high concentrations within specific granules of polymorphonuclear leukocytes.

Lysozyme along with lactoferrin and immunoglobulins, e.g., IgA, are widely recognized to be the predominant immunological factors in milk that manage the establishment of a protective commensal microflora population and control the growth of pathogens in the gastrointestinal tract of mammals.

While the capacity of lysozyme and lactoferrin to survive the digestive processes has long been appreciated in neonatal animal, several studies have examined this property in adult animals. (See, e.g., Mestecky *et al.*, 1998.) Thirty minutes following a single oral dose of lysozyme to mice, 10% of the dose was isolated in an enzymatically active form from the mid-jejunum and lysozyme activity could be measured throughout the intestine.

Furthermore, chicken lysozyme fed to rats can be observed by immunohistochemistry along the villi of the intestines, indicating a lack of proteolysis during transit through the digestive tract. Such studies demonstrate that native lysozyme and lactoferrin can survive the proteolytic and denaturing environment of the digestive tract. (See Example 6). Below is a discussion of several anti-microbial proteins suitable for use in the invention, classed as (Group 1) anti-microbial milk proteins, (Group 2), acute-phase proteins (other than Group-1 protein), (Group 3) anti-microbial peptides, and (Group 4), other anti-microbial proteins. Coding sequences for many of these proteins are given below. Other coding sequences, either for anti-microbial proteins from other sources, or other anti-microbial proteins, may be found in public gene databases, such as the GENBANK database, accessible through internet address "www.ncbi.nlm.gov/BLAST/".

Group 1. Anti-microbial milk proteins

Lysozyme, called muramidase or peptidoglycan N-acetylmuramoyl-hydrolase (EC 3.2.1.17) contains 130 amino acid residues (human lysozyme) and is a protein of 14.7 kDa in size. Human lysozyme is non-glycosylated and possesses unusual stability
5 *in vitro* and *in vivo* due to its amino acid and secondary structure.

Lysozyme is one of the most abundant proteins present in human milk with a concentration of about 400 µg/ml. The concentration of lysozyme is approximately 0.13 µg/ml in cow's milk (almost 3000 times less than found in human milk), 0.25 µg/ml in goat's milk, 0.1 µg/ml in sheep's milk and almost absent in rodent's milk (Chandan RC,
10 1968). Lysozyme is also found in other mammalian secretions, such as tears and saliva.

The protective role of lysozyme has been observed to include lysis of microbial cell walls, adjuvant activity of the end products peptidoglycan lysis, direct immunomodulating effects on leukocytes, and neutralization of bacterial endotoxins.
15 The bacteriostatic and bactericidal actions of lysozyme were originally discovered by Flemming in 1922 and have been studied in detail. Lysozyme is effective against both gram positive and gram negative bacteria, as well as some types of yeasts. The antimicrobial effects of lysozyme often act synergistically with other defense molecules, including immunoglobulin and lactoferrin. Furthermore, structural changes in the cell
20 wall due to lysozyme render bacteria more susceptible to phagocytosis by macrophages and neutrophils.

The hydrolysis of microbial peptidoglycans results in the release of the cleavage product, muramyl dipeptide, which is a potent adjuvant and is the active component of Freund's complete adjuvant. Muramyl dipeptide enhances IgA production, macrophage
25 activation, and rapid clearance of a variety of bacterial pathogens *in vivo*. Lysozyme itself is also immunomodulatory. It directly interacts with the cell membrane of phagocytes to increase their uptake of bacteria. Lysozyme also augments the proliferative response of mitogen stimulated lymphocytes to interleukin-2 and increases the rate of synthesis of IgG and IgM by more than 5-and 2- fold respectively.
30 Furthermore, the immunomodulatory action of lysozyme is not dependent upon enzymatic activity and is retained following denaturation. When lysozyme is fed to mice, it increases the number of intraepithelial and mesenteric lymph node lymphocytes that display antigens.

Lysozymes (either from human or non-human sources) act as enzymes that
35 cleave peptidoglycans, and ubiquitous cell wall component of microorganisms, in

particular bacteria. Specifically, lysozymes are 1,4- -acetylmuramidases that hydrolyze the glycoside bond between N-acetylmuramic acid and N-acetylglucosamine. Gram-positive bacteria are highly susceptible to lysozyme due to the polypeptidoglycan on the outside of the cell wall. Gram-negative strains have a single polypeptidoglycan layer covered by lipopolysaccharides and are therefore less susceptible to lysis by lysozyme, however, the sensitivity can be increased by the addition of EDTA (Schütte and Kula, 1990). Lysozyme also exhibits antiviral activity, as exemplified by the significant reduction in recurrent occurrences of genital and labial herpes after oral treatment of patients with lysozyme (Jollès, 1996). More recently, lysozyme from chicken egg whites, human milk and human neutrophils has been shown to inhibit the growth of HIV-1 in an *in vitro* assay (Lee-Huang *et al.*, 1999). In addition, an anti-fungal activity has been demonstrated for lysozymes using oral isolates of *Candida albicans* (the most common fungal causative agent of oropharyngeal infection in humans; (Samaranayake *et al.*, 1997). Lysozyme thus functions as a broad spectrum antimicrobial agent.

The ability of lysozyme to bind bacterial endotoxins, especially LPS, confers an important anti-microbial property to the molecule. Lysozyme binds electrostatically to the lipid A component of bacterial endotoxins at a 1:3 molar ratio. The resulting conformational change in endotoxin keeps it from interacting with macrophage receptors and dampens the release of pro-inflammatory cytokines such as interleukin-1 (IL-1), interleukin-6 (IL-6), and tumor necrosis factor (TNF). Thus, lysozyme exhibits anti-inflammatory activity during pathogen challenges.

The current major commercial source for lysozyme is chicken egg whites. Sequence analysis shows that lysozyme from chicken egg whites exhibits only partial homology (60%) with that synthesized by humans. Chicken and human lysozyme do not cross-react with their respective antibodies (Faure *et al.*, 1970), indicating significant structural differences between these two lysozymes. Human lysozyme has been purified from breast milk (Boesman-Finkelstein *et al.*, 1982; Wang *et al.*, 1984), neutrophils (Lollike *et al.*, 1995), and urine of hemodialysis patients (Takai *et al.*, 1996). Breast milk remains the main source for isolation of human lysozyme, but the supply is limited. Precautions are required for isolation of the enzyme from human sources to avoid contamination with viral and microbial pathogens.

Recombinant human lysozyme has been produced in the mammary gland of transgenic mice. The enzyme retained its antimicrobial activity, but the final concentration in the milk was low (Maga *et al.*, 1998; Maga *et al.*, 1994; Maga *et al.*, 1995). Human lysozyme has been expressed in *Aspergillus oryzae* (*A. oryzae*)

(Tsuchiya *et al.*, 1992) yeast (*S. cerevisiae*; Castañón *et al.*, 1988; Jigami *et al.*, 1986; and Yoshimura *et al.*, 1988) and in small amounts in tobacco leaves (Nakajima *et al.*, 1997). However, the expression level of recombinant human lysozyme in these organisms could be very low, and the cost of these forms may be prohibitive for food applications. In addition, human lysozyme produced in microorganisms may require extensive purification before it can be used in foods, particularly for infants and children.

In contrast to many other proteins, lysozyme is highly resistant to digestion in the gastrointestinal tract. *In vitro* studies have demonstrated that both molecules are resistant to hydrolysis by pepsin in the pH range found in the stomach. Furthermore, partial denaturation of lysozyme increases its bactericidal activity against some types of bacteria, and low pH, such as found in the stomach, increases the bactericidal effects of lysozyme. A proteolytic fragment (amino acids 98-112 of chicken egg white lysozyme) completely lacking enzymatic activity has been found to be the active bactericidal component of lysozyme. Additionally, a fragment of lactoferrin, known as lactoferricin, is formed by limited proteolytic digestion and has been shown to have extremely effective antibacterial activity.

The rice produced human lysozyme of the present invention exhibits acid pH resistance, as well as resistance to pepsin and pancreatin to make it resistant to digestion in the gastrointestinal tract. The excellent thermostability provides the feasibility to pasteurize products that include the recombinant human lysozyme.

Lactoferrin is an iron-binding protein found in the granules of neutrophils where it apparently exerts an antimicrobial activity by withholding iron from ingested bacteria and fungi; it also occurs in many secretions and exudates (milk, tears, mucus, saliva, bile, etc.). In addition to its role in iron transport, lactoferrin has bacteriostatic and bactericidal activities, in addition to playing a role as an anti-oxidant (Satue-Gracia *et al.*, 2000).

The mature human lactoferrin (LF) polypeptide consists of 692 amino acids, consists of a single-chain polypeptide that is relatively resistant to proteolysis, is glycosylated at two sites (N138 and N478) and has a molecular weight of about 80 kD. Human lactoferrin (hLF) is found in human milk at high concentrations (at an average of 1-3 mg/ml), and at lower concentration (0.1-0.3 mg/ml), in exocrine fluids of glandular epithelium cells such as bile, tears, saliva etc.

The primary functions of lactoferrin have been described as iron regulation, immune modulation and protection from infectious microbes. Lactoferrin can bind two ferric ions and has been shown to have biological activities including bacteriostatic

(Bullen *et al.*, 1972), bactericidal (Arnold, *et al.*, 1980) and growth factor activity *in vitro*. Further, lactoferrin can promote the growth of bacteria that are beneficial to the host organism by releasing iron in their presence. Additional studies have recently shown lactoferrin to have antiviral activity towards cytomegalovirus, herpes simplex virus, 5 rotavirus and HIV both *in vitro* and *in vivo*. (See, e.g., Fujihara *et al.*, 1995; Grover *et al.*, 1997; and Harmsen *et al.*, 1995.)

Lactoferrin, like transferrin, has a strong capacity to bind free iron under physiological conditions due to its tertiary structure, which consists of two globular lobes linked by an extended alpha-helix. The ability of lactoferrin to scavenge iron from the 10 physiological environment can effectively inhibit the growth of "more than 90% of all microorganisms" by depriving them of a necessary component of their metabolism, which will inhibit their growth *in vivo* and *in vitro*.

Unrelated to iron binding, the bactericidal activity of lactoferrin stems from its ability to destabilize the outer membrane of gram-negative bacteria through the 15 liberation of lipopolysaccharides that constitute the cell walls of the bacteria. Additionally, lactoferrin has recently been shown to bind to prions, a group of molecules common in *E. coli*, causing permeability changes in the cell wall. Studies in germfree piglets fed lactoferrin before being challenged with *E. coli* show significant decrease in mortality compared to the control group.

20 Recombinant LF (rLF) has been produced as a fusion protein in *Aspergillus oryzae* (Ward *et al.*, 1992) and in the baculovirus expression system (Salmon *et al.*, 1997). The *Aspergillus*-produced protein will require a high degree of purification as well as safety and toxicity testing prior to using it as a food additive (Lönnerdal, 1996). Lactoferrin has also been expressed in tobacco (*Nicotiana tabacum* L. cv Bright Yellow) 25 cell culture (Mitra and Zhang, 1994), tobacco plants (Salmon *et al.*, 1998) and potato (*Solanum tuberosum*) plants (Chong and Langridge, 2000). In tobacco cell culture the protein was truncated, whereas in tobacco and potato plants the rLF was processed correctly, but its expression level was very low (0.1% of total soluble protein) (Chong and Langridge, 2000). However, the expression level of recombinant human lactoferrin 30 in these organisms could be very low, and the cost of these forms may be prohibitive for food applications. In addition, human lactoferrin produced in microorganisms may require extensive purification before it can be used in foods, particularly for infants and children.

35 In contrast to most other proteins, lactoferrin has also been shown to be resistant to proteolytic degradation *in vitro*, with trypsin and chymotrypsin remarkably ineffective

in digesting lactoferrin, particularly in its iron-saturated form. Some large fragments of lactoferrin were formed, but proteolysis was clearly limited.

Lactoperoxidase is an enzyme which catalyzes the conversion of hydrogen peroxide to water. This enzyme is found in human milk, and plays host defensive roles through antimicrobial activity. When hydrogen peroxide and thiocyanate are added to raw milk, the SCN's oxidized by the enzyme-hydrogen peroxide complex producing bactericidal compounds which destroy Gram-negative bacteria (Shin).

Kappa-casein is a group of readily digested caseins, which account for almost half of the protein content in human milk, are important as nutritional protein for breast-fed infants. It has also been advocated that part of the antimicrobial activity of human milk resides in the caseins, most likely the glycosylated kappa-casein.

Alpha-1-antitrypsin is a protease inhibitor that acts to prevent the digestion of the milk proteins, for example lysozyme and lactoferrin, in newborn infant gut, is the presence of protease inhibitors, such as α -1-antitrypsin (AAT) in human milk. AAT belongs to the class of serpin inhibitors, has a molecular mass of 52 kD, and contains about 15% carbohydrate (Carrell et al, 1983). Concentrations of AAT in human milk range from 0.1 to 0.4 mg/mL, and they appear to decrease with infant age (Davidson and Lönnerdal, 1979; McGilligan et al., 1987). While the binding affinity of AAT is highest for human neutrophil elastase, it also has affinity for pancreatic proteases such as chymotrypsin and trypsin (Beatty et al., 1980). The role of AAT in milk is possible that AAT may prolong the survival of other anti-microbial proteins through inhibition of pancreatic proteases. While anti-microbial proteins have been expressed in systems such as transgenic cows and *Aspergillus* (Lönnerdal, 1996), transgenic rice provides a more attractive vehicle for the production of recombinant human AAT for food applications. High levels of expression are possible by using the combination of regulatory elements such as promotor, signal peptide, and terminator, and optionally, transcription factors, as disclosed herein.

Lactadherin is a nonimmunological component in human milk that helps can protect breast-fed infants against infection by microorganisms. One of the major protective glycoproteins is lactadherin. Protection against certain virus infections by human milk is associated with lactadherin. (Newburg, 1999, 1998; Peterson; Hamosh).

Immunoglobulins, e.g., IgA, are present in human act to confer resistance to a variety of pathogens to which the mother may have been exposed. (See, for example, Humphreys; Kortt; Larrick; Maynard; and Peeters.

Group 2. Acute-phase proteins

C-reactive protein (CRP) is a protein generally having a very low blood concentration, which rises up to two thousand times following inflammatory processes [J. J. Morley and I. Kushner, Am. N.Y. Acad. Sci., 389, 406-418 (1989)].

5 Serum amyloid A (SAA) is known to be an acute phase protein whose concentration increases about a thousand-fold, usually within 24 hours, during any inflammatory disorder.

Ferritin is a large (480 kD) intracellular iron storage protein which converts ferrous (Fe^{2+}) iron to the ferric (Fe^{3+}) state and sequesters up to 4500 iron atoms in the ferric state per molecule of ferritin. Hepatic ferritin content and serum ferritin concentrations increase rapidly after administration of interleukin-6 to rats. Kobune et al., Hepatology (1994); 19: 1468-1475. These effects of pro-inflammatory cytokines on ferritin synthesis are part of the hypoferremic response which occurs early in inflammation. Rogers, "Genetic regulation of the iron transport and storage genes: links with the acute phase response." In Iron and Human Disease. Lauffer, editor. (1992); 15 CRC Press, Boca Raton, pp. 77-104.

Haptoglobin (Hp) is a hemoglobin-binding serum protein which plays a major role in the protection against heme-driven oxidative stress (Langlois M R and Delanghe J R (1996) Clin Chem 42: 1589-1600; Delanghe J R et al. (1998) AIDS 12: 1027-1032; 20 Gutteridge J M. (1987) Biochim Biophys Acta 917: 219-223; Miller Y I et al. (1997) Biochem 36: 12189-12198; Vercellotti G M et al. (1994) Art Cell Blood Substit Imm Biotech 22: 207-213).

Seromucoids are glycoproteins derived from blood serum.

25 Ceruloplasmin (Cp) is a copper-containing glycoprotein which plays an important role in the metabolism of copper. The ceruloplasmin level in serum of a patient suffering from Wilson's disease, or an infectious disease, or of a pregnant woman is different from that of a normal human, so that quantification of human ceruloplasmin may be used for the diagnosis of these diseases and pregnancy.

30 15-keto-13,14-dihydro-prostaglandin F2 alpha (PGFM) is a prostanoid produced by the placental. Prostanoids are a family of autacoids (formed from arachidonic acid) thought to play an important role during implantation, in the progress and maintenance of pregnancy, and during the initiation and progress of labor (Angle and Johnston, 1990).

35 Fibrinogen (Fb) is a large protein molecule that normally circulates in the blood plasma in the dissolved state. Under attack from the enzyme thrombin, the fibrinogen

molecules link up, spontaneously aligning themselves into a long thread like polymer or network called fibrin which is the primary ingredient of blood clots. Fibrinogen itself comprises 6 chains including two copies of an alpha, beta and gamma chain.

Alpha-1-acid glycoprotein (AGP) is a human plasma glycoprotein which is produced by the liver. AGP is an acute-phase reactant, i.e., the concentration of AGP in the blood increases following inflammation. AGP synthesis increases several fold during an acute phase response (Baumann, H. et al. J. Biol. Chem. 256:10145-10155 (1981); Ricca, G. A. et al. J. Biol. Chem. 256:11199-11202 (1981); Koj, A. et al. Biochem J. 206:545-553 (1982); Koj, A. et al. Biochem J. 224:505-514 (1984)). The major acute phase inducers of AGP synthesis are the cytokines interleukin-1 (IL-1) and interleukin-6 (IL-6), which act additively to induce transcription of the AGP gene.

Mannose binding protein (MBP) is thought to play a role in the disposal of pathogenic organisms. MBP works both by opsonizing pathogen, and by activating the complement cascade. MBP consists of several monomers that assemble into one larger multimer.

Lipopolysaccharide binding protein (LBP) is a 60 kD glycoprotein synthesized in the liver and present in normal human serum. LBP belongs to the group of plasma proteins called acute phase proteins, including C-reactive protein, fibrinogen and serum amyloid A, that increase in concentration in response to infectious, inflammatory and toxic mediators. LBP expression has been induced in animals by challenge with lipopolysaccharide (LPS), silver nitrate, turpentine and *Corynebacterium parvum* [Geller et al., Arch. Surg. 128(1): 22-28 (1993); Gallay et al., Infect. Immun. 61(2): 378-383 (1993); Tobias et al., J. Exp. Med. 164: 777-793 (1986)].

Alpha-2 macroglobulin and various defensins

Alpha-2 macroglobulin is a 718 kDa homotetrameric glycoprotein and is a well characterized as an extracellular proteinase inhibitor.

The milk proteins lysozyme and lactoferrin, discussed above, are also included in this group.

Group 3: Antimicrobial Peptides

As many as 450 antimicrobial peptides have been found from amphibians, insects, mammals, plants, microorganisms and fishes. These antimicrobial peptides are known to be different in their sizes and amino acid sequences, but similar in their antimicrobial mechanism. Representative antimicrobial peptides include cecropin, magainin, bombinin, defensin, tachyplesin, and buforin. All of these are composed of 17-24 amino acids, showing antimicrobial activity against

a broad spectrum of microorganisms, including Gram-negative bacteria, Gram-positive bacteria, protozoa and fungi. Some of these peptides are effective against both cancer cells and viruses. For instance, magainin, consisting of 23 amino acids, was isolated from the skin of an amphibian and is reported not only to defend against pathogenic bacteria, but to kill human lung cancer cells (Zasloff, M. (1987) Proc. Natl. Acad. Sci., U.S.A. 84, 5449-5453).

Antimicrobial peptides comprising six conserved cysteines form the defensin family. This family is composed of antimicrobial peptides which are present in numerous species, which are abundant and which are about 3-4 kDa (Ganz and Lehrer, 1994). These peptides are formed of 30 to 40 amino acids, of which six invariant cysteines which form three intramolecular disulfide linkages. They have complex conformation, are amphipathic, rich in beta antiparallel sheets but lack alpha helices (Lehrer and Ganz, 1992). The antimicrobial action of defensins is thought to result from their insertion into the membranes of the target cells, allowing the formation of voltage-dependent channels. White et al. (1995) describe the possible mechanisms of membrane insertion and of formation of multimeric pores by the defensins, which allow the permeabilization of the membranes of the target cells, for example microbial or tumor cells. Defensins have an antimicrobial action on a broad spectrum of microorganisms in vitro (Martin et al., 1995). This activity spectrum, which is particularly broad, comprises bacteria, Gram-positive and Gram-negative bacteria, several fungi, mycobacteria, parasites including spirochetes and several enveloped viruses including the HSV and HIV viruses. They are also cytotoxic for several categories of normal and malignant cells, including cells resistant to TNF-alpha and to the cytolytic NK factor (Kagan et al., 1994).

Indolicidin is a peptide that has been isolated from bovine neutrophils and has antimicrobial activity, including activity against viruses, bacteria, fungi and protozoan parasites. Subbalakshmi C et al., *Biochem Biophys Res Commun*, 2000, 274(3):714-6.

Cecropin is an antimicrobial peptide that is isolated from the silkworm. Choi et al., *Comp Biochem Physiol C Toxicol Pharmacol*, 2000, 125:287-97.

Magainins are a group of short peptides originally isolated from frog skin and thought to function as a natural defense mechanism against infection due to their antimicrobial properties. Li et al., *Planta*, 2001, 212:635-9.

Tachyplesin I (T-SS) is a membrane-permeabilizing antimicrobial peptide discovered from horseshoe crab hemolymph. Kobayashi S et al., *Biochemistry*, 2001, 40:14330-5.

Parasin I (Park et al., FEBS Lett, 1998, 437:258-62 and U.S. Patent No. 6,316,594). Parasin I is a potent antimicrobial peptide which is secreted in response to an epidermal injury by *Parasilurus asotus*, a catfish.

Buforin I is a 39-residue antimicrobial peptide derived from the N-terminal of toad histone H2A [Kim et al. (1996) Biochem. Biophys. Res. Commun. 229, 381-387].

PMAP-23 (Park et al., *Biochem Biophys Res Commun*, 2002, 290:204-12).

PMAP-23 is a cathelicidin-derived antimicrobial peptide identified from porcine leukocytes. PMAP-23 demonstrates potent antimicrobial activity against Gram-negative and Gram-positive bacteria without hemolytic activity.

Moronecidin: Lauth, et al., *J Biol Chem*, 2002, 277:5030-5039. Moronecidin is a 22-residue, C-terminally amidated antimicrobial peptide which is isolated from the skin and gill of hybrid striped bass. Two isoforms, differing by only one amino acid, are derived from each parental species, white bass (*Morone chrysops*) and striped bass (*Morone saxatilis*). Molecular masses (2543 and 2571 Da), amino acid sequences (FFHHIFRGIVHVGKTIH(K/R)LVTGT SEQ ID NO:42), cDNA, and genomic DNA sequences were determined for each isoform. A predicted 79-residue moronecidin prepropeptide consists of three domains: a signal peptide (22 amino acids), a mature peptide (22 amino acids), and a C-terminal prodomain (35 amino acids). The synthetic, amidated white bass moronecidin exhibited broad spectrum antimicrobial activity that was retained at high salt concentration. The moronecidin gene consists of three introns and four exons. Peptide sequence and gene organization were similar to pleurocidin, an antimicrobial peptide from winter flounder. A TATA box and several consensus-binding motifs for transcription factors were found in the region 5[prime prime or minute] to the transcriptional start site. Moronecidin gene expression was detected in gill, skin, intestine, spleen, anterior kidney, and blood cells by kinetic reverse transcription (RT)-PCR. Thus, moronecidin is a new [alpha]-helical, broad spectrum antimicrobial peptide isolated from the skin and gills of hybrid striped bass.

Anoplin: Konno et al, *Biochim Biophys Acta*, 2001, 1550:70-80. Anoplin is an antimicrobial peptide which was purified from the venom of the solitary wasp *Anoplius samariensis*. Anoplin is composed of 10 amino acid residues, Gly-Leu-Leu-Lys-Arg-Ile-Lys-Thr-Leu-Leu-NH₂ and is the smallest among the linear alpha-helical antimicrobial peptides hitherto found in nature. Anoplin is the first antimicrobial component to be found in the solitary wasp venom and it may play a key role in preventing potential infection by microorganisms during prey consumption by their larvae. Biological evaluation using the synthetic peptide revealed that this peptide exhibited potent activity

in stimulating degranulation from rat peritoneal mast cells and broad-spectrum antimicrobial activity against both Gram-positive and Gram-negative bacteria.

Gambicin: Vizioli J. et al., *Proc Natl Acad Sci U S A*, 2001, 98:12630-5.

Gambicin is a antimicrobial peptide which is excreted by the mosquito. The 616-bp
5 gambicin ORF encodes an 81-residue protein that is processed and secreted as a 61-aa
mature peptide. Gambicin lacks sequence homology with other known proteins. Like
other *Anopheles gambiae* antimicrobial peptide genes, gambicin is induced by natural or
experimental infection in the midgut, fatbody, and hemocyte-like cell lines. Within the
midgut, gambicin is predominantly expressed in the anterior part. Both local and
10 systemic gambicin expression is induced during early and late stages of natural malaria
infection. In vitro experiments showed that the 6.8-kDa mature peptide can kill both
Gram-positive and Gram-negative bacteria, has a morphogenic effect on a filamentous
fungus, and is marginally lethal to *Plasmodium berghei* ookinetes.

SAMP-29: Shin SY et al., *Biochem Biophys Res Commun* 2001 Jul

15 27;285(4):1046-51. AMP-29 is a cathelicidin-derived antimicrobial peptide deduced
from sheep myeloid mRNA.

Group 4-Other anti-microbial proteins

CAP37: Cationic antimicrobial protein of Mr 37 kDa (CAP37) is a neutrophil-
derived inflammatory mediator on endothelial cell function.

20 Granulysin: A mechanism by which T cells contribute to host defense against
microbial pathogens is release of the antimicrobial protein granulysin. Ochoa et al., *Nat*
Med, 2001, 7:174-9.

SLPI: Secretory leukocyte protease inhibitor (SLPI) exhibits antimicrobial activity.
(Shugars et al., *Gerontology*, 2001, 47:246-53.)

25 Calprotectin: Squamous mucosal epithelial cells constitutively express
calprotectin in the cytoplasm. Cells expressing calprotectin resist invasion by *Listeria*
monocytogenes and *Salmonella enterica* serovar Typhimurium. Nisapakultorn K et al.,
Infect Immun, 2001, 69:3692-6.

30 CAP18: Cationic antimicrobial protein (CAP18) is an 18 kDa antimicrobial
protein.

Ubiquicidin: Ubiquicidin is a cationic, small (Mr 6654) protein which displays
marked antimicrobial activity against *Listeria monocytogenes* and *Salmonella*
typhimurium and activity against *Escherichia coli*, *Staphylococcus aureus*, and an
avirulent strain of *Yersinia enterocolitica*. Hiemstra et al., *J Leukoc Biol*, 1999, 66:423-8.

BAMP-1: Bovine antimicrobial protein-1 (BAMP-1) is a 76 amino acid residue antimicrobial protein which has been isolated from fetal calf serum. BAMP-1 showed a weak growth-inhibitory activity against *Escherichia coli* and yeasts tested in phosphate-buffered saline (PBS).: J Biochem (Tokyo) 1998 Apr;123(4):675-9.

5 Ace-AMP1 is an antifungal protein extracted from onion seeds. This cationic protein contains 93 amino acid residues and four disulfide bridges. Tassin et al., *Biochemistry*, 1998, 37:3623-37.

Small granules of horseshoe crab hemocytes contain two known major antimicrobial substances, tachyplesin and big defensin (S5). Kawabata et al., J
10 *Biochem* (Tokyo), 1996, 120:1253-60.

Ac-AMP2 is a lectin-like small protein with antimicrobial and antifungal activity isolated from *Amaranthus caudatus*. Verheyden et al., *FEBS Lett*, 1995, 370:245-9.

Ah-AMP1 (Fant et al., *Proteins*, 1999, 37:388-403). *Aesculus hippocastanum* antimicrobial protein 1 (Ah-AMP1) is a plant defensin isolated from horse chestnuts. The
15 plant defensins have been divided in several subfamilies according to their amino acid sequence homology. Ah-AMP1 is a member of subfamily A2 and inhibits growth of a broad range of fungi.

CAP18 (Cationic antimicrobial protein of 18 kDa) is an antimicrobial protein found in human and rabbit granulocytes.

20 IV. Expression Vectors For Generation Of Transgenic Plants Expressing Anti-microbial Proteins

Expression vectors for use in the present invention are chimeric nucleic acid constructs (or expression vectors or cassettes), designed for operation in plants, with
25 associated upstream and downstream sequences.

In general, expression vectors for use in practicing the invention include the following operably linked components that constitute a chimeric gene: (i) a transcriptional regulatory region from a monocot gene having a seed maturation-specific promoter, (ii) operably linked to said transcriptional regulatory region, a leader DNA
30 sequence encoding a monocot seed-specific transit sequence capable of targeting a linked polypeptide to an endosperm-cell organelle, such as the leader sequence for targeting to a protein-storage body, and (iii) a protein-coding sequence encoding an anti-microbial protein

The chimeric gene, in turn, is typically placed in a suitable plant-transformation
35 vector having (i) companion sequences upstream and/or downstream of the chimeric

gene which are of plasmid or viral origin and provide necessary characteristics to the vector to permit the vector to move DNA from bacteria to the desired plant host; (ii) a selectable marker sequence; and (iii) a transcriptional termination region generally at the opposite end of the vector from the transcription initiation regulatory region.

5 Exemplary methods for constructing chimeric genes and transformation vectors carrying the chimeric genes are given in Example 1.

A. Promoters

In one aspect of this embodiment, the expression construct includes a
10 transcription regulatory region (promoter) which exhibits specifically upregulated activity during seed maturation. Examples of such promoters include the maturation-specific promoter region associated with one of the following maturation-specific monocot storage proteins: rice glutelins, oryzins, and prolamines, barley hordeins, wheat gliadins and glutenins, maize zeins and glutelins, oat glutelins, and sorghum kafirins, millet
15 pennisetins, and rye secalins. Exemplary regulatory regions from these genes are exemplified by SEQ ID NOS: 18-24, as identified below.

Of particular interest is the expression of the nucleic acid encoding an anti-microbial protein from a transcription initiation region that is preferentially expressed in plant seed tissue. Examples of such seed preferential transcription initiation sequences
20 include those sequences derived from sequences encoding plant storage protein genes or from genes involved in fatty acid biosynthesis in oilseeds. Exemplary preferred promoters include a glutelin (Gt-1) promoter, as exemplified by SEQ ID NO: 18, which effects gene expression in the outer layer of the endosperm and a globulin (Glb) promoter, as exemplified by SEQ ID NO: 19, which effects gene expression in the center
25 of the endosperm. Promoter sequences for regulating transcription of gene coding sequences operably linked thereto include naturally-occurring promoters, or regions thereof capable of directing seed-specific transcription, and hybrid promoters, which combine elements of more than one promoter. Methods for construction such hybrid promoters are well known in the art.

30 In some cases, the promoter is derived from the same plant species as the plant cells into which the chimeric nucleic acid construct is to be introduced. Promoters for use in the invention are typically derived from cereals such as rice, barley, wheat, oat, rye, corn, millet, triticale or sorghum.

Alternatively, a seed-specific promoter from one type of monocot may be used regulate transcription of a nucleic acid coding sequence from a different monocot or a non-cereal monocot.

Numerous types of appropriate expression vectors, and suitable regulatory sequences are known in the art for a variety of plant host cells. The transcription regulatory or promoter region is chosen to be regulated in a manner allowing for induction under seed-maturation conditions. Examples of such promoters include those associated with the following monocot storage proteins: rice glutelins, oryzins, and prolamines, barley hordeins, wheat gliadins and glutelins, maize zeins and glutelins, oat glutelins, and sorghum kafirins, millet pennisetins, and rye secalins. Exemplary promoter sequences are identified herein as SEQ ID NOS: 18-24. Other promoters suitable for expression in maturing seeds include the barley endosperm-specific B1-hordein promoter (Brandt, A., et al., (1985), Glub-2 promoter, Bx7 promoter, Gt3 promoter, Glub-1 promoter and Rp-6 promoter, particularly if these promoters are used in conjunction with transcription factors. The primary structure of a B1 hordein gene from barley is provided in Carlsberg Res. Commun. 50, 333-345).

B. Signal/targeting/transport Sequences

In addition to encoding the protein of interest, the expression cassette or heterologous nucleic acid construct may encode a signal/targeting/transport peptide that allows processing and translocation of the protein, as appropriate. Exemplary signal/targeting/transport sequences, particularly for targeting proteins to intracellular bodies, such as vacuoles, are signal/targeting sequences associated with the monocot maturation-specific genes: glutelins, prolamines, hordeins, gliadins, glutenins, zeins, albumin, globulin, ADP glucose pyrophosphorylase, starch synthase, branching enzyme, Em, and lea. Exemplary sequences encoding a leader sequence for protein storage body are identified herein as SEQ ID NOS: 24-30.

In one preferred embodiment, the method is directed toward the localization of recombinant milk protein expression in a given subcellular compartment, in particular a protein-storage body, but also including the mitochondrion, endoplasmic reticulum, vacuoles, chloroplast or other plastidic compartment. For example, when recombinant milk protein expression is targeted to plastids, such as chloroplasts, in order for expression to take place the construct also employ the use of sequences to direct the gene to the plastid. Such sequences are referred to herein as chloroplast transit peptides (CTP) or plastid transit peptides (PTP). In this manner, when the gene of

interest is not directly inserted into the plastid, the expression construct additionally contains a gene encoding a transit peptide to direct the gene of interest to the plastid. The chloroplast transit peptides may be derived from the gene of interest, or may be derived from a heterologous sequence having a CTP. Such transit peptides are known
5 in the art. See, for example, Von Heijne *et al.*, 1991; Clark *et al.*, 1989; della-Cioppa *et al.*, 1987; Romer *et al.*, 1993; and Shah *et al.*, 1986. Additional transit peptides for the translocation of the protein to the endoplasmic reticulum (ER) (Chrispeels, K., 1991), nuclear localization signals (Raikhel, 1992), or vacuole may also find use in the constructs of the present invention.

10 Another exemplary class of signal/targeting/transport sequences are sequences effective to promote secretion of heterologous protein from aleurone cells during seed germination, including the signal sequences associated with α -amylase, protease, carboxypeptidase, endoprotease, ribonuclease, DNase/RNase, (1-3)- β -glucanase, (1-3)(1-4)- β -glucanase, esterase, acid phosphatase, pentosamine, endoxylanase, β -
15 xylopyranosidase, arabinofuranosidase, β -glucosidase, (1-6)- β -glucanase, peroxidase, and lysophospholipase.

Since many protein storage proteins are under the control of a maturation-specific promoter, and this promoter is operably linked to a leader sequence for targeting to a protein body, the promoter and leader sequence can be isolated from a single
20 protein-storage gene, then operably linked to a milk-protein storage protein in the chimeric gene construction. One preferred and exemplary promoter-leader sequence is from the rice Gt1 gene, having an exemplary sequence identified by SEQ ID NO:15. Alternatively, the promoter and leader sequence may be derived from different genes. One preferred and exemplary promoter/leader sequence combination is the rice Glb
25 promoter linked to the rice Gt1 leader sequence, as exemplified by SEQ ID NO: 16.

C. Protein Coding Sequences

The construct also includes the nucleic acid coding sequence for a heterologous protein, under the control of a promoter, preferably a seed-specific promoter. In
30 accordance with the present invention, polynucleotide sequences which encode a heterologous anti-microbial protein, such as lysozyme or lactoferrin, include splice variants, fragments of such proteins, fusion proteins, modified forms or functional equivalents thereof, collectively referred to herein as "anti-microbial protein-encoding nucleic acid sequences".

Such "anti-microbial protein-encoding nucleic acid sequences" may be used in recombinant expression vectors (also termed heterologous nucleic acid constructs), that direct the expression of an anti-microbial protein in appropriate host cells.

Due to the inherent degeneracy of the genetic code, a number of nucleic acid sequences which encode substantially the same or a functionally equivalent amino acid sequence may be generated and used to clone and express a given anti-microbial protein, as exemplified herein by the codon optimized coding sequences used to practice the invention (further described below). Thus, for a given anti-microbial protein-encoding nucleic acid sequence, it is appreciated that as a result of the degeneracy of the genetic code, a number of coding sequences can be produced that encode the same protein amino acid sequence. For example, the triplet CGT encodes the amino acid arginine. Arginine is alternatively encoded by CGA, CGC, CGG, AGA, and AGG. Therefore such substitutions in the coding region fall within the range of sequence variants covered by the present invention. Any and all of these sequence variants can be utilized in the same way as described herein for a "reference" anti-microbial- encoding nucleic acid sequence.

A "variant" anti-microbial protein-encoding nucleic acid sequence may encode a "variant" human anti-microbial amino acid sequence which is altered by one or more amino acids from the native anti-microbial protein sequence, both of which are included within the scope of the invention. Similarly, the term "modified form of", relative to a given anti-microbial protein, means a derivative or variant form of an anti-microbial protein or the coding sequence therefor. That is, a "modified form of" a anti-microbial protein has a derivative sequence containing at least one nucleic acid or amino acid substitution, deletion or insertion. The nucleic acid or amino acid substitution, insertion or deletion may occur at any residue within the sequence, as long as the encoded amino acid sequence maintains the biological activity of the native anti-microbial protein, e.g., the bactericidal effect of lysozyme.

A "variant" anti-microbial protein-encoding nucleic acid sequence may encode a "variant" anti-microbial protein sequence which contains amino acid insertions or deletions, or both. Furthermore, a variant anti-microbial protein coding sequence may encode the same polypeptide as the reference polynucleotide or native sequence but, due to the degeneracy of the genetic code, has a nucleic acid coding sequence which is altered by one or more bases from the reference or native polynucleotide sequence.

The variant nucleic acid coding sequence may encode a variant amino acid sequence which contains a "conservative" substitution, wherein the substituted amino

acid has structural or chemical properties similar to the amino acid which it replaces and physicochemical amino acid side chain properties and high substitution frequencies in homologous proteins found in nature (as determined, e.g., by a standard Dayhoff frequency exchange matrix or BLOSUM matrix). In addition, or alternatively, the variant
5 nucleic acid coding sequence may encode a variant amino acid sequence which contains a "non-conservative" substitution, wherein the substituted amino acid has dissimilar structural or chemical properties to the amino acid which it replaces.

Standard substitution classes include six classes of amino acids based on common side chain properties and highest frequency of substitution in homologous
10 proteins in nature, as is generally known to those of skill in the art and may be employed to develop variant anti-microbial protein-encoding nucleic acid sequences. A "variant" anti-microbial protein-encoding nucleic acid sequence may encode a "variant" anti-microbial protein sequence which contains a combination of any two or three of amino acid insertions, deletions, or substitution.

15 Anti-microbial protein-encoding nucleotide sequences also include "allelic variants" defined as an alternate form of a polynucleotide sequence which may have a substitution, deletion or addition of one or more nucleotides, which does not substantially alter the function of the encoded polypeptide.

The polynucleotides for use in practicing the invention include sequences which
20 encode anti-microbial proteins and splice variants thereof, sequences complementary to the protein coding sequence, and novel fragments of the polynucleotide. The polynucleotides may be in the form of RNA or in the form of DNA, and include messenger RNA, synthetic RNA and DNA, cDNA, and genomic DNA. The DNA may be double-stranded or single-stranded, and if single-stranded may be the coding strand or
25 the non-coding (anti-sense, complementary) strand.

As will be understood by those of skill in the art, in some cases it may be advantageous to use an anti-microbial protein-encoding nucleotide sequences possessing non-naturally occurring codons. Codons preferred by a particular eukaryotic host (Murray *et al.*, 1989) can be selected, for example, to increase the rate of
30 heterologous protein expression or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, than transcripts produced from naturally occurring sequence. Codon-optimized sequences for use in practicing the invention are further described below.

An anti-microbial protein-encoding nucleotide sequence may be engineered in
35 order to alter the protein coding sequence for a variety of reasons, including but not

limited to, alterations which modify the cloning, processing and/or expression of the protein by maturing seeds.

Heterologous nucleic acid constructs may include the coding sequence for a given anti-microbial protein, a variant, fragment or splice variant thereof: (i) in isolation; (ii) in combination with additional coding sequences; such as fusion protein or signal peptide, in which the anti-microbial protein coding sequence is the dominant coding sequence; (iii) in combination with non-coding sequences, such as introns and control elements, such as promoter and terminator elements or 5' and/or 3' untranslated regions, effective for expression of the coding sequence in a suitable host; and/or (iv) in a vector or host environment in which the anti-microbial protein coding sequence is a heterologous gene.

Depending upon the intended use, an expression construct may contain the nucleic acid sequence which encodes the entire anti-microbial protein, or a portion thereof. For example, where anti-microbial protein sequences are used in constructs for use as a probe, it may be advantageous to prepare constructs containing only a particular portion of the anti-microbial protein encoding sequence, for example a sequence which is discovered to encode a highly conserved anti-microbial protein region.

Codon-optimized coding sequences for human milk anti-microbial proteins are identified herein by SEQ ID NOS: 1, 3, and 7-14. An expression vector used to practice the invention has at least 70%, preferably 80%, 85%, 90% or 95% or more sequence identity to the codon-optimized sequences.

Coding sequences for exemplary non-human acute-phase proteins are identified herein by SEQ ID NOS: 36, and 46-56. An expression vector used to practice the invention has at least 70%, preferably 80%, 85%, 90% or 95% or more sequence identity to the identified sequences.

Coding sequences for exemplary anti-microbial peptides are identified herein by SEQ ID NOS: 34-68, 40-41, and 43. An expression vector used to practice the invention has at least 70%, preferably 80%, 85%, 90% or 95% or more sequence identity to the identified sequences.
codon-optimized sequence.

Coding sequences for other anti-microbial proteins are identified herein by SEQ ID NOS: 37, 45, and 57-59. An expression vector used to practice the invention has at least 70%, preferably 80%, 85%, 90% or 95% or more sequence identity to the identified sequences.

Additional codon sequences for anti-microbial proteins are available from public gene databases, such as GENBANK, accessible through "www.ncbi.nlm.gov/GENBANK/".

5 D. Codon Optimization

It has been shown that production of recombinant protein in transgenic barley grain was enhanced by codon optimization of the gene (Horvath *et al.*, 2000; Jensen *et al.*, 1996). The intent of codon optimization was to change an A or T at the third position of the codons of G or C. This arrangement conforms more closely with codon usage in
10 typical rice genes (Huang *et al.*, 1990a).

In order to obtain a high expression level for human lysozyme in rice cells, the coding sequence was codon optimized. The G + C content was thus increased from 46% to 68%. The codon optimized lysozyme coding sequence for use in practicing the invention is presented as SEQ ID NO:1.

15 Similarly, in order to obtain high level expression level of human lactoferrin (hLF) in rice cells, the native hLF coding sequence was codon optimized. Out of 693 codons used in the lactoferrin gene, 413 codons were changed by one or two nucleotides. The amino acid sequence of LF was unchanged. The codon optimized lactoferrin coding sequence for use in practicing the invention is presented as SEQ ID NO:3.

20 Codon optimized sequences for other human milk proteins are given as follows: for lactoferricin, SEQ ID NO: 7; for EGF, SEQ ID NO: 8; for IGF-1, SEQ ID NO: 9; for lactohedrin, SEQ ID NO: 10; for kappa-casein, SEQ ID NO: 11; for haptocorrin, SEQ ID NO: 12; for lactoperoxidase, SEQ ID NO: 13; for and for alpha-1-antitrypsin, SEQ ID NO: 14.

25

E. Transcription Factor Coding Sequences

In one embodiment of the invention, the transgenic plant is also transformed with the coding sequence of one or more transcription factors capable of stimulating the expression of a maturation-specific promoter. Specifically, the embodiment involves the
30 use of the maize *Opaque 2* (O2) and prolamin box binding factor (PBF) together with the rice endosperm bZip (Reb) protein as transcriptional activators of monocot storage protein genes. Exemplary sequence for these three transcription factors are given identified below as SEQ ID NOS: 31-33. Transcription factor sequences and constructs applicable to the present invention are detailed in co-owned PCT application No.

PCT/US01/14234, International Publication number WO 01/83792 A1, published November 8, 2001, which is incorporated herein by reference.

Transcription factors are capable of sequence-specific interaction with a gene sequence or gene regulatory sequence. The interaction may be direct sequence-specific binding in that the transcription factor directly contacts the gene or gene regulatory sequence or indirect sequence-specific binding mediated by interaction of the transcription factor with other proteins. In some cases, the binding and/or effect of a transcription factor is influenced (in an additive, synergistic or inhibitory manner) by another transcription factor. The gene or gene regulatory region and transcription factor may be derived from the same type (e.g., species or genus) of plant or a different type of plant. The binding of a transcription factor to a gene sequence or gene regulatory sequence may be evaluated by a number of assays routinely employed by those of skill in the art, for example, sequence-specific binding may be evaluated directly using a label or through gel shift analysis.

As detailed in the cited PCT application, the transcription factor gene is introduced into the plant in a chimeric gene containing a suitable promoter, preferably a maturation-specific seed promoter operably linked to the transcription factor gene. Plants may be stably transformed with a chimeric gene containing the transcription factor by methods similar to those described with respect to the milk-protein gene(s). Plants stably transformed with both exogenous transcription factor(s) and milk-protein genes may be prepared by co-transforming plant cells or tissue with both gene constructs, selecting plant cells or tissue that have been co-transformed, and regenerating the transformed cells or tissue into plants. Alternatively, different plants may be separately transformed with exogenous transcription factor genes and milk-protein genes, then crossed to produce plant hybrids containing by added genes.

F Additional Expression Vector Components

Expression vectors or heterologous nucleic acid constructs designed for operation in plants, comprise companion sequences upstream and downstream to the expression cassette. The companion sequences are of plasmid or viral origin and provide necessary characteristics to the vector to permit the vector to move DNA from bacteria to the plant host, such as, sequences containing an origin of replication and a selectable marker. Typical secondary hosts include bacteria and yeast.

In one embodiment, the secondary host is *E. coli*, the origin of replication is a colE1-type, and the selectable marker is a gene encoding ampicillin resistance. Such

sequences are well known in the art and are commercially available as well (e.g., Clontech, Palo Alto, Calif.; Stratagene, La Jolla, CA).

The transcription termination region may be taken from a gene where it is normally associated with the transcriptional initiation region or may be taken from a different gene.

5 Exemplary transcriptional termination regions include the NOS terminator from *Agrobacterium* Ti plasmid and the rice α -amylase terminator.

Polyadenylation tails (Alber *et al.*, 1982) may also be added to the expression cassette to optimize high levels of transcription and proper transcription termination, respectively. Polyadenylation sequences include, but are not limited to, the *Agrobacterium* octopine synthetase signal, Gielen, *et al.*, 1984 or the nopaline synthase of the same species, Depicker, *et al.*, 1982.

Suitable selectable markers for selection in plant cells include, but are not limited to, antibiotic resistance genes, such as, kanamycin (*nptII*), G418, bleomycin, hygromycin, chloramphenicol, ampicillin, tetracycline, and the like. Additional selectable markers include a bar gene which codes for bialaphos resistance; a mutant EPSP synthase gene which encodes glyphosate resistance; a nitrilase gene which confers resistance to bromoxynil; a mutant acetolactate synthase gene (ALS) which confers imidazolinone or sulphonylurea resistance; and a methotrexate resistant DHFR gene.

The particular marker gene employed is one which allows for selection of transformed cells as compared to cells lacking the DNA which has been introduced. Preferably, the selectable marker gene is one which facilitates selection at the tissue culture stage, e.g., a kanamycin, hygromycin or ampicillin resistance gene.

The vectors of the present invention may also be modified to include intermediate plant transformation plasmids that contain a region of homology to an *Agrobacterium tumefaciens* vector, a T-DNA border region from *Agrobacterium tumefaciens*, and chimeric genes or expression cassettes (described above). Further, the vectors of the invention may comprise a disarmed plant tumor inducing plasmid of *Agrobacterium tumefaciens*.

In general, a selected nucleic acid sequence is inserted into an appropriate restriction endonuclease site or sites in the vector. Standard methods for cutting, ligating and *E. coli* transformation, known to those of skill in the art, are used in constructing vectors for use in the present invention. (See generally, Maniatis, *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2d Edition (1989); Ausubel, *et al.*, (c) 1987, 1988, 1989, 1990, 1993, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley

& Sons, New York, N.Y; and Gelvin, S.B., *et al.*, eds. PLANT MOLECULAR BIOLOGY MANUAL, (1990), all three of which are expressly incorporated by reference, herein.

V. Generation of Transgenic Plants

5 Plant cells or tissues are transformed with expression constructs (heterologous nucleic acid constructs, e.g., plasmid DNA into which the gene of interest has been inserted) using a variety of standard techniques. Effective introduction of vectors in order to facilitate enhanced plant gene expression is an important aspect of the invention. It is preferred that the vector sequences be stably integrated into the host
10 genome.

The method used for transformation of host plant cells is not critical to the present invention. The transformation of the plant is preferably permanent, *i.e.* by integration of the introduced expression constructs into the host plant genome, so that the introduced constructs are passed onto successive plant generations. The skilled
15 artisan will recognize that a wide variety of transformation techniques exist in the art, and new techniques are continually becoming available.

Any technique that is suitable for the target host plant may be employed within the scope of the present invention. For example, the constructs can be introduced in a variety of forms including, but not limited to, as a strand of DNA, in a plasmid, or in an
20 artificial chromosome. The introduction of the constructs into the target plant cells can be accomplished by a variety of techniques, including, but not limited to calcium-phosphate-DNA co-precipitation, electroporation, microinjection, *Agrobacterium*-mediated transformation, liposome-mediated transformation, protoplast fusion or microprojectile bombardment. The skilled artisan can refer to the literature for details
25 and select suitable techniques for use in the methods of the present invention. Exemplary methods for plant transformation are given in Example 2.

When *Agrobacterium* is used for plant cell transformation, a vector is introduced into the *Agrobacterium* host for homologous recombination with T-DNA or the Ti- or Ri-plasmid present in the *Agrobacterium* host. The Ti- or Ri-plasmid containing the T-DNA
30 for recombination may be armed (capable of causing gall formation) or disarmed (incapable of causing gall formation), the latter being permissible, so long as the *vir* genes are present in the transformed *Agrobacterium* host. The armed plasmid can give a mixture of normal plant cells and gall.

In some instances where *Agrobacterium* is used as the vehicle for transforming
35 host plant cells, the expression or transcription construct bordered by the T-DNA border

region(s) is inserted into a broad host range vector capable of replication in *E. coli* and *Agrobacterium*, examples of which are described in the literature, for example pRK2 or derivatives thereof. See, for example, Ditta *et al.*, 1980 and EPA 0 120 515, expressly incorporated by reference herein. Alternatively, one may insert the sequences to be
5 expressed in plant cells into a vector containing separate replication sequences, one of which stabilizes the vector in *E. coli*, and the other in *Agrobacterium*. See, for example, McBride *et al.*, 1990, wherein the pRiHRI (Jouanin, *et al.*, 1985, origin of replication is utilized and provides for added stability of the plant expression vectors in host *Agrobacterium* cells.

10 Included with the expression construct and the T-DNA is one or more selectable marker coding sequences which allow for selection of transformed *Agrobacterium* and transformed plant cells. A number of markers have been developed for use with plant cells, such as resistance to chloramphenicol, kanamycin, the aminoglycoside G418, hygromycin, or the like. The particular marker employed is not essential to this
15 invention, with a particular marker preferred depending on the particular host and the manner of construction.

For *Agrobacterium*-mediated transformation of plant cells, explants are incubated with *Agrobacterium* for a time sufficient to result in infection, the bacteria killed, and the plant cells cultured in an appropriate selection medium. Once callus forms, shoot
20 formation can be encouraged by employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be grown to seed and the seed used to establish repetitive generations and for isolation of the recombinant protein produced by the plants.

25 There are a number of possible ways to obtain plant cells containing more than one expression construct. In one approach, plant cells are co-transformed with a first and second construct by inclusion of both expression constructs in a single transformation vector or by using separate vectors, one of which expresses desired genes. The second construct can be introduced into a plant that has already been
30 transformed with the first expression construct, or alternatively, transformed plants, one having the first construct and one having the second construct, can be crossed to bring the constructs together in the same plant.

A. Plants

Host cells of the present invention include plant cells, both monocotyledenous and dicotyledenous. In one preferred embodiment, the plants used in the methods of the present invention are derived from monocots, particularly the members of the taxonomic family known as the *Gramineae*. This includes all members of the grass family of which the edible varieties are known as cereals. The cereals include a wide variety of species such as wheat (*Triticum* sps.), rice (*Oryza* sps.) barley (*Hordeum* sps.) oats, (*Avena* sps.) rye (*Secale* sps.), corn (maize) (*Zea* sps.) and millet (*Pennisetum* sps.). In practicing the present invention, preferred grains are rice, wheat, maize, barley, rye, triticale. Also preferred are dicots exemplified by soybean (*Glycine* spp.)

In order to produce transgenic plants that express anti-microbial protein, monocot plant cells or tissues derived from them are transformed with an expression vector comprising the coding sequence for a anti-microbial protein. Transgenic plant cells obtained as a result of such transformation express the coding sequence for a anti-microbial protein, such as lysozyme or lactoferrin. The transgenic plant cells are cultured in medium containing the appropriate selection agent to identify and select for plant cells which express the heterologous nucleic acid sequence. After plant cells that express the heterologous nucleic acid sequence are selected, whole plants are regenerated from the selected transgenic plant cells. Techniques for regenerating whole plants from transformed plant cells are generally known in the art.

Transgenic plant lines, e.g., rice, wheat, corn or barely, can be developed and genetic crosses carried out using conventional plant breeding techniques.

Production of recombinant proteins in monocot seeds, e.g., rice (*Oryza sativa* L.) seeds has the advantages that (a) high level expression make it an economically practical strategy, and (b) rice is a normal part of the diet of infants and children, has good nutritional value and low allergenicity. Thus, the use of rice as the basis for a food supplement is unlikely to introduce any risk and thereby eliminates the need for a high degree of purification when included in infant formula.

In addition, rice is the staple food crop of more than half the world's population. Recent reports on the production of provitamin A (beta-Carotene) in rice seeds exemplifies the need for value added food crops especially in the developing world (Ye *et al.*, 2000) where rice is used as major food crop.

VI. Detecting Expression of Recombinant Anti-microbial Proteins

Transformed plant cells are screened for the ability to be cultured in selective media having a threshold concentration of a selective agent. Plant cells that grow on or in the selective media are typically transferred to a fresh supply of the same media and
5 cultured again. The explants are then cultured under regeneration conditions to produce regenerated plant shoots. After shoots form, the shoots are transferred to a selective rooting medium to provide a complete plantlet. The plantlet may then be grown to provide seed, cuttings, or the like for propagating the transformed plants. The method provides for efficient transformation of plant cells with expression of a gene of
10 autologous or heterologous origin and regeneration of transgenic plants, which can produce a recombinant anti-microbial protein.

The expression of the recombinant anti-microbial protein may be confirmed using standard analytical techniques such as Western blot, ELISA, PCR, HPLC, NMR, or mass spectroscopy, together with assays for a biological activity specific to the particular
15 protein being expressed.

Example 3 describes the characterization of human lysozyme produced in the seeds of transgenic rice plants. Analyses used to confirm that recombinant lysozyme produced in transgenic rice is essentially the same as the native form of the protein both in physical characteristics and biological activity included, SDS-PAGE, reverse IEF gel
20 electrophoresis, Western blot analysis, enzyme linked immunosorbant assay (ELISA), enzymatic activity assay and bactericidal activity assay using indicator strains, *Micrococcus luteus* and *E.coli* strain JM109.

Example 4 describes the characterization of human lactoferrin produced in the seeds of transgenic rice plants. Analyses used to confirm that recombinant lactoferrin
25 produced in transgenic rice is essentially the same as the native form of the protein both in physical characteristics and biological activity included, Southern blot, Western blot, ELISA, N-Terminal Amino Acid Sequencing, analysis of glycosylation and determination of sugar content, a determination of the isoelectric point, pH dependent iron release of rLF, bacteriostatic activity assay of rLF using enteropathogenic *E. coli* as the indicator
30 strain.

VII. Improved feed and methods

The invention provides, in one aspect, an improved feed containing a flour, extract, or malt seed composition obtained from mature monocot seeds and one or more
35 seed-produced anti-microbial proteins in substantially unpurified form. Exemplary anti-

microbial proteins are lysozyme and lactoferrin, where lysozyme is preferably present in an amount between about 0.05 and 0.5 grams protein/kg feed, and lactoferrin, in an amount between 0.2 to 2 grams/protein/kg feed. The feed to which the seed composition is added may be any standard feed suitable for a given production animals, e.g., chickens or cattle, and typically includes plant-produced components, such as grains or grain flour from corn, rice, barley and/or wheat. Exemplary feed for chickens is given in Example 7. In accordance with the invention, the animal feed to which the seed composition is added contains little or no small-molecule antibiotic(s), i.e., antibiotic levels that are themselves ineffective to optimize weight gain in the production animal.

To determine the amount of seed composition to be added to the feed, one first determines the approximate desired weight percent or concentration of the given anti-microbial protein or proteins in the final feed material. Typically, the amount of protein anti-microbial protein effective for optimal weight gain in production animals will be in the range 0.05 to 5 mg protein/ gram dry weight of feed, e.g., between about 0.005 to 0.5 weight percent of the anti-microbial protein. Effective levels can be determined, for example, by feed studies of the type reported in Example 7, where increasing amounts of anti-microbial protein, in the range indicated above, are added to feed, and weight gain and intestinal characteristics determined after several days-weeks of feeding.

The amount of seed composition added to the feed will, of course, depend on the concentration of the anti-microbial protein in the seed composition, and this concentration can be readily determined by standard protein assay methods, as described in Examples 3 and 4 below. For a grain composition, the level of heterologous protein present will be roughly that produced in seeds, e.g., between 0.1 to 1% of total seed weight. In this case, the final amount of grain composition added to the feed may be in the range 1-30%, depending on the final required level of anti-microbial protein. For an extract composition, the heterologous protein may be concentrated to form up to 5-40% or more of the total extract weight, so a much smaller percentage of extract composition will be required to achieve the desired level of anti-microbial protein, for example, between 0.1 to 10% by weight, typically, 1-5% of the total feed weight. A malt concentration, which will contain a significant percent of malt sugars, in addition to native and heterologous proteins, will typically contain an amount of anti-microbial protein that is intermediate between that of grain and the extract, and the amount added to feed will, accordingly, typically fall within the ranges of the other two compositions.

As an example, it may be desired to have a final concentration of lysozyme between about 0.05 and 0.5 grams protein/kg feed, and lactoferrin, in an amount

between 0.2 to 2 grams/protein/kg feed. Thus, if a seed composition is found to contain 10g/kg lysozyme, about 20 grams of the composition would be added to make up a liter of formula with a final lysozyme concentration of about 0.2 g/liter.

Below are described methods for preparing each of the three types of milk-protein-containing seed compositions.

A. Flour composition

The flour composition is prepared by milling mature monocot plant seeds, using standard milling and, optionally, flour purification methods, e.g., in preparing refined flour. Briefly, mature seeds are dehusked, and the dehusked seeds then ground into a fine flour by conventional milling equipment.

The flour may be added to feed in powdered, particulate, or liquid form, in an amount typically between 1 to 30% or more of the dry weight of the feed, depending on the desired final amount of anti-microbial protein needed, as described above. One exemplary flour composition includes at least lactoferrin and/or lysozyme, where lysozyme is preferably present in an amount between about 0.05 and 0.5 grams protein/kg feed, and lactoferrin, in an amount between 0.2 to 2 grams/protein/kg feed, in addition to other anti-microbial proteins. Flour containing two or more anti-microbial proteins may be prepared by combining flour from seeds that separately produce the different proteins, for example, equal amounts of a flour containing lysozyme and a flour containing lactoferrin. Alternatively, a multi-protein composition can be prepared as seed flour from plants monocot plants co-transformed with chimeric genes expressing different milk proteins, e.g., lactoferrin and lysozyme.

B. Extract composition

This composition is prepared by milling flour to form a flour, extracting the flour with an aqueous buffered solution, and optionally, further treating the water-soluble extract to partially concentrate the extract and/or remove unwanted components. Details of exemplary methods for producing the extract composition are given in Example 6.

Briefly, mature monocot seeds, such as rice seeds, are milled to a flour, and the flour then suspended in a buffered solution. The flour suspension is incubated with shaking for a period typically between 30 minutes and 4 hours, at a temperature between 20-55°C. The resulting homogenate is clarified either by filtration or centrifugation. The clarified filtrate or supernatant may be further processed, for example by ultrafiltration or dialysis or both to remove contaminants such as lipids, sugars and salt. Finally, the

material may be dried, e.g., by lyophilization, to form a dry cake or powder. A variety of aqueous media are suitable for the extraction buffer, including phosphate buffered saline (PBS) and ammonium bicarbonate, as demonstrated in Example 6. Volatile buffers like ammonium bicarbonate, in which the salt components of the buffer are volatilized on drying, may obviate an additional salt removal step, and thus offer a significant processing advantage.

The extract combines advantages of high protein yields, essentially limiting losses associated with protein purification. At the same time, the anti-microbial proteins are in a form readily usable and available upon ingestion of the animal feed. One feature for use in a feed product is the low amount of seed starch present in the extract. In particular, the extract may increase the concentration of recombinant protein from about 0.5% in conventional approaches to over about 25% in the extract approach. Some extract approach even reached 40% depending on the expression level of recombinant protein. In addition, the extract approach removes starch granules, which require high gelling temperature, for example above about 75°C. Consequently, the extract approach provides more flexibility in processing the rice grain and the recombinant proteins into feed. In the limiting case where the extract is processed to near protein purification or homogeneity, the amount of anti-microbial protein present may be in the range 70-95% of the extract.

The extract can be formulated and added to animal feed in powdered, or dry particleized or tabletized or powder form. In one embodiment, the extract is added to feed in an amount between about 0.5 to 10% by weight, preferably 1-5% by dry weight of the feed. An exemplary feed contains both lactoferrin and lysozyme, where lysozyme is preferably present in an amount between about 0.05 and 0.5 grams protein/kg feed, and lactoferrin, in an amount between 0.2 to 2 grams/protein/kg feed. The extract may alternatively, or in addition, include one or more of the other anti-microbial proteins noted above.

As above, extract containing two or more milk proteins may be prepared by combining extracts from seeds that separately produce the different proteins, or by processing seeds from plants co-transformed with chimeric genes expressing different milk proteins, e.g., lactoferrin and lysozyme.

C. Malt composition

In accordance with another embodiment, the invention provides a malt extract or malt syrup ("malt") in which seed starches have been largely reduced to malt sugars,

and the anti-microbial protein(s) are in an active, bioavailable form. A variety of feed products can be produced, depending on the type of malt used, the mashing program and the ways in which the wort is subsequently handled. If materials other than barley malt are used in the mash (such as starch from other grains), the resulting product is referred to herein as a sweetened malt.

Malt extracts, which may have a syrupy consistency or may be powders, are made by mashing ground malt, usually barley malt, in conventional brewery equipment, collecting the wort and concentrating it or drying it. Modern production of food malt extracts and malt syrups has evolved into three basic grain stages: steeping, germination, and drying of the germinated seed, followed by three more steps involving liquefaction of the germinated grain, mashing of the germinated grain, lautering (filtering), and evaporation. Many variations of malt extracts or syrups are possible. Flavor, color, solids, enzymatic activity, and protein are the basic characteristics that can be adjusted during production to provide malts specific for given food applications. (See, generally, Eley; Hickenbottom, 1996, 1997a, 1997b, 1983; Lake; Moore; Moe; Sfat; Doncheck; Briggs, 1981, 1998; and Hough).

C1. Steeping

After the barley of choice has been cleaned of foreign material, it is graded to size and transferred to steep tanks equipped with water inlet and outlet pipes. Compressed air is fed from the tank bottom for vigorous aeration and mixing for the barley/water mixture. When the barley has reached a water content of 43-45%, steeping is stopped.

C2. Germination

The steeped barley is moved to germination floors or rooms depending on the particular malt house's capabilities and allowed to germinate under controlled temperature, air, and moisture conditions. Total germination varies from four to seven days, depending on the barley type, density end use of the malt, and the controls or germination method used. All aspects of germination must be kept in constant balance to ensure proper kernel modification and yield.

Many enzymatic systems are activated during germination. Two of the systems are the oxidative and reductive systems involved with the respiration phase. Other enzymes break down the endosperm cell structure, which in itself is a measure of germination rate when the pentose production is evaluated. The proteolytic enzymes release or active beta-amylase and also work on the proteins present to render them soluble. In fact, about 40% of the total protein is made soluble in water. Optimum germination activates a balanced enzyme system, which hydrolyzes the starch present.

C3. Kilning

Drying or kilning, when done at the proper time and optimum degree of starch modification, stops the germination. The heat also catalyzes additional reactions, notably flavor and color development. The heating step is carried out at conventional
5 kilning time and temperature conditions well known to those in the field. When drying is complete, the sprouts and other extraneous materials are removed, and the kernels are then ready for further processing.

C4. Malt Extracts And Syrups

10 The malted barley (kernel) is coarsely ground in crushers and fed into mash tuns where it is mixed with water. During a series of time and temperature changes, some of the starch is converted into fermentable sugars by action of the natural alpha- and beta-amylases, better known as the diastatic system. If cereal adjuncts are to be added,
15 which result in malt syrups with mellow and sweeter flavors than the extracts, they are added at this stage usually derived from the cereal grains, corn and rice, although barley, wheat, rye, millet and sorghum are sometimes used, derived from mature seeds that produce the desired recombinant milk proteins.

Once the mash batch has achieved the correct degree of hydrolysis, it is transferred to lauter tuns. The lauter tun has a slotted or false bottom a few inches above the real
20 bottom to allow for filtration and is also equipped with some means of agitation. During this extraction stage, the amylolytic enzymes liquefy additional insoluble starches, converting them to maltose and dextrans. At the same time, the proteolytic enzymes attach certain proteins converting them into simpler, soluble forms. After the appropriate conditions have been met, the liquid phase, or wort, is drawn from the lauter tuns into
25 evaporators.

Evaporation of the wort is conducted under vacuum where it is converted into a syrup of about 80% solids. Depending on the temperatures used, malt extracts or syrups of high, medium, or zero enzymatic activity can be produced. Color and flavor
30 also can be controlled during this stage. The finishing steps of filtering, cooling, and packaging complete the malt extract/syrup process.

C5. Transgenic Malt Extract

For a transgenic malt extract, the starting barley is a transgenic barley engineered to produce one or more anti-microbial proteins in the endosperm either in
35 grain maturation or in the malting process, or at both times. Malting and processing

times and conditions are adjusted so that the bioactivity of the target recombinant molecules is preserved and the bioavailability of the target recombinant molecule is maximized. The resulting malt extract may be added directly to animal feed, in a desired weight ratio. Studies conducted in support of the present invention demonstrate that

5 recombinant anti-microbial proteins lysozyme and lactoferrin retain activity after malting for up to 288 hrs. Thus, the malting procedure is useful in that it allows for extraction of microbial proteins in bioavailable form, with little loss of protein activity, and in a final composition that can serve as a source of sugar in the feed.

10 C6. Transgenic Sweetened Malt

For a transgenic sweetened malt syrup or extract, the starting barley can be a non-transgenic barley, or a transgenic barley, or a mixture of both. The barley is processed as described, except that during the mashing process, a cereal adjunct is added in a form that it is converted during the mashing process with the concurrent retention and

15 generation of bioavailability and bioactivity of the target recombinant molecule found within the transgenic cereal adjunct. The use of a transgenic cereal adjunct enables the production in the malt syrup of the target recombinant molecule expressed in the transgenic grain endosperm.

Preferred malt extracts or syrups contain lactoferrin and/or lysozyme. The malt

20 may alternatively, or in addition, include one or more of the antimicrobial proteins noted above, such as the antimicrobial milk proteins lactohedrin, kappa-casein, haptocorrin, lactoperoxidase, and alpha-1-antitrypsin. As above, malt containing two or more antimicrobial proteins may be prepared by combining or preparing malts from seeds that separately produce the different proteins, or by preparing a malt from the seeds of plants

25 co-transformed with chimeric genes expressing different milk proteins, e.g., lactoferrin and lysozyme.

D. Improved Feed Method

In accordance with another important aspect of the invention, it has been

30 discovered (see Examples 6 and 7) that the nutritionally enhanced feed of the invention provides the same or improved intestinal health and weight gain as feed supplemented with small-molecule antibiotics. The improved method includes substituting conventional feed with a feed supplemented with one or more small-molecule antibiotics or a feed that has been nutritionally enhanced by addition of the seed composition of present

35 invention, containing one or more anti-microbial proteins in substantially unpurified form.

Preferably, conventional feed is substituted with feed nutritionally enhanced by the addition of lysozyme or lactoferrin in the seed composition. Preferably, the feed may be supplemented with antibiotics such as roxarsone and bactiracin methylene disalicylate. The provision of feed supplemented with one or more small molecule antibiotics or a feed enhanced by the seed composition comprising one or more antimicrobial proteins results in a feed method with improved feed efficiency and improved intestinal health of the animals to which it is administered.

From the foregoing, it can be appreciated how various objects and features of the invention are met. The results presented herein demonstrate that milk proteins may be expressed at high levels in the seeds of transgenic plants. Once produced, the protein-containing grain of the invention finds utility in improved animal feed that may be fed to production animals. When fed to animals, the improved animal feed of the invention finds utility in increasing the growth rate and feed conversion efficiency of the animals, particularly for animals in a typical production environment (*i.e.*, a low-sanitation environment), thereby reducing or eliminating the need for administration of ST antibiotics. The production of high levels of anti-microbial proteins in grains, exemplified herein by rice, provides the distinct advantage that feed supplements may be prepared with little or no purification. In one preferred approach, the transgenic grain is ground (*e.g.*, into flour) and directly added to a production-animal feed, without additional processing.

Transgenic seeds are ideal bioreactors, combining low production costs and low or minimal downstream processing costs prior to use. Seed grain proteins can accumulate to 9-19% of grain weight (Lásztitym 1996); the endosperm proteins are synthesized during grain maturation and stored in protein bodies for use in the germination and seedling growth of the next plant generation; grains can be stored for years without loss of functionality, and therefore the downstream processing can be conducted independently of growing seasons.

Below are described methods for making grain, extract and malt compositions from mature monocot seeds containing seed-produced milk proteins. The composition is added to a feed or feed product in powdered, liquid, or particulate form, in an amount sufficient to bring the amount of added milk protein to effective levels.

All publications, patents and patent applications are herein expressly incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

EXAMPLE 1

Expression Vectors for Generation of transgenic Plants

In general, expression vectors were constructed using standard molecular biological techniques as described in Ausubel *et al.*, 1987. The vectors contain a
5. heterologous protein coding sequence for lactoferrin or lysozyme under the control of a rice tissue-specific promoter, as further described below.

A. An Expression Vector For Human Lysozyme Expression In Transgenic Rice Cells

The synthesized lysozyme gene was cloned into an API base vector pAPI137 by
10 conventional molecular cloning techniques (Sambrook *et al.*, 1989). Plasmid pAPI137 contains the *RAmy3D* promoter (Huang *et al.*, 1993), the codons for the *RAmy3D* signal peptide and the *RAmy3D* terminator. The *RAmy3D* promoter, isolated from the rice amylase gene family, is activated in rice calli by sugar starvation (Huang *et al.*, 1993). The human lysozyme gene was placed between the sequences of the *RAmy3D* signal
15 peptide and the *RAmy3D* terminator to give plasmid pAPI156 (Fig. 1) having a size of 4829 bp. Plasmid, pAPI76 carrying the bacterial hygromycin B phosphotransferase (hpt) gene was used for co-transformation of the calli to allow selection of the transformants.

Briefly, pAPI176 was created as following: A DNA fragment was amplified from a
20 rice alpha-amylase gene *RAmy1A* (Huang *et al.*, 1990b) and cloned into pBluescript KS+ at the *SmaI/EcoRI* restriction sites and the resulting plasmid was called p1AT. The PCR amplified fragment contained 297 bp of *RAmy1A* terminator. A *BamHI* DNA fragment from pGL2 (Shimamoto *et al.*, 1989) was cloned into the *BamHI* site of p1AT and the resulting plasmid was called pAPI174. Finally a *SacI/XbaI* fragment amplified
25 from glucanase gene, *Gns9* (Romero *et al.*, 1998) was inserted into pAPI174 using the same restriction sites. The PCR generated, *Gns9* promoter fragment was confirmed by DNA sequencing. The resulting plasmid was named as pAPI176.

B. An Expression Vector For Human Lactoferrin Expression In Transgenic Rice

30 The hLF gene (Rey, 1990) was codon optimized and synthesized by Operon Technologies (CA, USA). The plasmid containing the codon-optimized gene was called Lac-ger. Lac-ger was digested with *SmaI/XhoI* and the fragment containing the lactoferrin gene was cloned into pAPI141 that was partially digested with *NaeI* and completely digested with *XhoI*. The resulting plasmid was named pAPI 164. For

expression of hLF in rice seeds, the codon optimized gene was operably linked to the rice endosperm specific glutelin (Gt1) promoter and NOS terminator (Fig. 7).

The Gt1 promoter was cloned based on the published DNA sequence (Okita *et al.*, 1989). Genomic DNA was isolated from a rice variety called M202 and the Gt1 promoter isolated by PCR. The primers used to amplify the fragment were MV-Gt1-F1:
5' ATC GAA GCT TCA TGA GTA ATG TGT GAG CAT TAT GGG ACC ACG 3' (SEQ ID NO:5) and Xba-Gt1-R1: 5' CTA GTC TAG ACT CGA GCC ATG GGG CCG GCT AGG GAG CCA TCG CAC AAG AGG AA 3' (SEQ ID NO:6). The PCR amplifications were carried out using the GeneAmp PCR system (model 2400, Perkin-Elmer) operated according to the manufacturers instructions. Basic cycling conditions were 30 cycles, after a 2 minute pre-denaturing step at 95 °C, with a 30 second denaturing step at 95 °C, a 30 second annealing step at specific temperature, and a 2 minute extension step at 72 °C. The final extension step was 5 minutes at 72 °C, followed by 4 °C soaking step. Reaction components per 50 µl volume, were 1 µg of genomic DNA or 1 ng of plasmid DNA, 2.5 µl of 5 µM primer mixture, 5 µl of 10 mM dNTP, 2.5 units of *Taq* polymerase (Perkin-Elmer), 5 µl of 10X PCR buffer (Perkin-Elmer).

The amplified fragment was then cloned in both orientations with the resulting plasmids designated pCRGT1 and pCRGT1R. pCRGT1R was digested with *HindIII/Xba I* and the fragment cloned into pAPI 135 containing the kanamycin gene. The resulting plasmid was named as pAPI141.

To visualize and confirm the tissue specificity of the cloned promoter obtained by PCR, a GUS gene and NOS terminator was obtained from pRAJ275 (ClonTech, USA) and digested with *EcoRI/NcoI*. The fragment was inserted into pAPI141 and the resulting plasmid designated pAPI142.

The plasmid pAPI176 was constructed in three steps. A DNA fragment was amplified from a rice alpha-amylase gene, *Ramy1A*, (Huang *et al.*, 1990b) and cloned into pBluescript KS+ at the *SmaI/EcoRI* restriction sites. The PCR amplified fragment contained 297 bp of *Ramy1A* terminator. The resulting plasmid was called p1AT. A *BamHI* DNA fragment from pGL2 (Shimamoto *et al.*, 1989) was cloned in to *BamHI* site of p1AT. The *BamHI* fragment contained most of the hygromycin phosphotransferase (Hph) gene with deletion of four amino acids at C-terminus. This plasmid was called as pAPI174. Finally, a *SacI/XbaI* fragment amplified from glucanase gene, *Gns9* (17) was inserted into pAPI174 using same restriction sites. The PCR generated, *Gns9*, promoter fragment was confirmed by DNA sequencing. The resulting plasmid was named as pAPI176.

EXAMPLE 2

Generation of Transgenic Plant Cells Expressing Anti-Microbial Proteins or Peptides

The procedure of microprojectile-mediated rice transformation (Chen *et al.*, 1998 and Sivamani *et al.*, 1996) was followed with modifications. Calli was raised from
5 TP309 mature rice seeds, with calli two to four mm in diameter selected and placed on N6 media supplemented with 0.3 M mannitol and 0.3 M sorbitol for 20 hours before bombardment. Biolistic bombardment was carried out with the biolistic PDC-1000/He system (Bio-Rad, USA). Plasmids pAPI164 and pAPI176 were gold coated and co-bombarded at a ratio of 6:1 with a helium pressure of 1100psi. Two-day old bombarded
10 calli were then transferred to N6 selection media supplemented with 20 mg/l hygromycin B and allowed to grow in the dark at 26°C for 45 days.

In order to develop transgenic rice plants, the selected calli were transferred to pre-regeneration and regeneration media. When regenerated plants became 1-3 cm in height, the plantlets were transferred to rooting media consisting of half concentration of
15 MS and 0.05 mg/l NAA. After two weeks, plantlets with developed roots and shoots were transferred to soil and kept under the cover of plastic container for a week. The plants were allowed to grow about 12 cm tall and shifted to the green house where they were grown up to maturity.

20 A. Generation of Human Lysozyme Expressing Transgenic Rice Cells and Plants

The synthetic human lysozyme (hLys) gene under the control of the *RAmy3D* promoter and terminator in the pAPI156 plasmid was used to generate sixty independent transformants by particle bombardment-mediated transformation.

25 Particle bombardment mediated transformation of rice was carried out as described above. (See Chen *et al.*, 1998.) Briefly, rice calli derived from TP309 were bombarded with gold particles coated with plasmids pAPI156 and pAPI176 in a ratio of 6:1 using the helium biolistic particle delivery system, PDS 1000 (Bio-Rad, CA). Transformed calli were selected in the presence of hygromycin B (35 mg/L) on N6
30 (Sigma, MO).

Selected cell lines were maintained in culture media with 3% sucrose (Huang *et al.*, 1993). Lysozyme expression was induced by sugar starvation. Briefly, AA medium (containing 3% sucrose) was removed by aspiration, followed by washing the cells three times with AA minus sucrose (AA-S). The cells were then incubated with AA-S at 40%
35 (v/v) density for three and a half days to obtain the optimal level of lysozyme expression.

Transformants expressing lysozyme were identified by immunoblot analysis, turbidimetric rate determination with *Micrococcus lysodeikticus* or ELISA. Calli were ranked according to the expressed lysozyme level. Suspension cell cultures from the top lines were established following the procedure described previously (Huang *et al.*, 1993). The amount of total protein (Bradford assay) and lysozyme (ELISA) was evaluated in selected calli (Table 1).

Table 1. Expression Level Of Human Milk Lysozyme In Transformed Calli

Cell line	Calli (g)	Total protein (μ g)	Lysozyme (μ g)	Lysozyme/protein (%)
156-1	0.39	2626.5	65.7	2.5
156-5	0.38	5510	68.9	1.25
156-16	0.4	4815	120.4	2.5
156-19	0.44	2440	30.5	1.25
156-28	0.49	4910	24.6	0.5
156-43	0.56	8150	101.9	1.25
156-47	0.37	2472	6.2	0.25

Transformed calli were selected as described above, then transferred to pre-regeneration and regeneration media. When regenerated plants became 1-3 cm in height, the plantlets were transferred to rooting media which consisted of half concentration of MS and 0.05 mg/l NAA. After two weeks, plantlets with developed roots and shoots were transferred to soil and kept under the cover of plastic container for a week. The plants were allowed to grow about 12 cm tall and shifted to the green house where they were grown up to maturity (R0 plants). Figure 2 illustrates the seed specific expression of human lysozyme in transgenic plants and Fig. 6 shows the expression level of human lysozyme in powdered R3 seeds taken from transgenic rice plants. In completing this analysis, the human lysozyme in powdered R3 seeds was extracted by mixing rice powder (prepared by grinding with a Kitchen Miller) with phosphate buffer saline (PBS) containing 0.35 N NaCl at 1 gm/40 mL for one hour. The whole homogenate was allowed to settle, 1 mL homogenate was removed and centrifuged at 4°C and 14000 rpm for 15 min. The supernatant was removed and diluted as needed for lysozyme assay by ELISA.

Embryos from individual R1 seed (derived from R0 plants) that showed a level of lysozyme expression that was greater than 10 μ g/seed were saved and used to generate R1 plants. Briefly, seeds were dissected into embryo and endosperm portions. The endosperm was ground and assayed for lysozyme expression (as further described below). Embryos were sterilized in 50% commercial bleach for 25 minutes and washed

with sterile H₂O three times for 5 minutes each. Sterilized embryos were placed in a tissue culture tube that contained MS solid medium. Embryos germinated and plantlets having about three inches shoots and healthy root systems were obtained in two weeks. The plantlets were then transferred to pots to obtain mature plants (R1).

5

B. Generation of Human Lactoferrin Expressing Transgenic Rice Cells and Plants

The synthetic human lysozyme gene under the control of the Gt1 promoter in the pAPI164 plasmid was used to generate over 100 independent transformants by particle bombardment-mediated transformation.

10

Particle bombardment mediated transformation of rice was carried out as described in (Chen *et al.*, 1998). Briefly, rice calli derived from TP309 were bombarded with gold particles coated with plasmids pAPI164 and pAPI76 in a ratio of 6:1 using the helium biolistic particle delivery system, PDS 1000 (Bio-Rad, CA).

15

Two day old bombarded calli were then transferred to N6 selection media supplemented with 20 mg/l hygromycin B and allowed to grow in the dark at 26°C for 45 days. The selected calli were then transferred to pre-regeneration and regeneration media. When regenerated plants became 1-3 cm in height, the plantlets were transferred to rooting media which consisted of half concentration of MS and 0.05 mg/l NAA. After two weeks, plantlets with developed roots and shoots were transferred to soil and kept under the cover of plastic container for a week. The plants were allowed to grow about 12 cm tall and shifted to the green house where they were grown up to maturity.

20

25

EXAMPLE 3

Characterization Of Recombinant Human Lysozyme (rLys) Produced by Transgenic Rice Cells and Plants

A. Purification, SDS-PAGE and reverse IEF gel electrophoresis

Induced calli or harvested cells from suspension cell cultures were ground with cold phosphate buffered-saline (PBS) with a protease inhibitor cocktail (2 µg/ml aprotonin, 0.5 µg/ml leupeptin, 1 mM EDTA and 2 mM Pefabloc). The protease inhibitor cocktail was excluded from the buffer used subsequently during the purification of the enzyme, since the inhibitors did not increase the lysozyme expression yield. Grinding was conducted with a pre-chilled mortar and pestle at approximately 2 ml buffer/g calli

30

or cells. A clear homogenate was obtained by subjecting the resulting extract to centrifugation at 16,000 x g for 10 minutes at 4 C.

SDS-PAGE was carried out using an 18% precast gel (Novex, CA). The resulting gel was stained with 0.1% Coomassie brilliant blue R-250 at 45% methanol and 10% glacial acetic acid for three hours. Gel destaining was conducted with 45% methanol and 10% glacial acetic acid until the desired background was reached.

Reverse IEF gel electrophoresis was carried out using a precast Novex pH 3-10 IEF gel according to the manufacturer's instructions (Novex, CA). About 30 µg of lysozyme was loaded onto the gel and electrophoresed at 100 V for 50 minutes followed by application of 200 V for 20 minutes. The gel was then fixed in 136 mM sulphosalicylic acid and 11.5% TCA for 30 minutes and stained in 0.1% Coomassie brilliant blue R-250, 40% ethanol, 10% glacial acetic acid for 30 minutes. The destaining solution contained 25% ethanol and 8% acetic acid.

15 B. Western blot analysis

A SDS-PAGE gel was electroblotted to a 0.45 µm nitrocellulose membrane using a Mini Trans-Blot Electrophoretic Transfer Cell (Bio-Rad, CA) and subsequently subjected to immuno-blotting analysis. The blot was blocked with 5% non-fat dry milk in PBS, pH 7.4 for at least two hours followed by three washes with PBS, pH 7.4 for 10 minutes each. The primary rabbit polyclonal antibody against human lysozyme (Dako A/S, Denmark) was diluted at 1:2000 in the blocking buffer and the blot was incubated in the solution for at least one hour. The blot was then washed with PBS three times for 10 minutes each. The secondary goat anti-rabbit IgG (H+L)-alkaline phosphatase conjugate (Bio-Rad, CA) was diluted in the blocking buffer at 1:4000. The membrane was then incubated in the secondary antibody solution for one hour and then washed three times. Color development was initiated by adding the substrate system BCIP-NBT (Sigma) and the process was stopped by rinsing the blot with H₂O once the desirable intensity of the bands had been achieved.

30 C. Enzyme Linked Immunosorbant Assay (ELISA)

An indirect sandwich ELISA was developed to quantify total lysozyme expressed in rice calli or cells and used as an alternative assay to determine the lysozyme expression yield. A direct sandwich ELISA for lysozyme quantification has been previously reported (Lollike *et al.*, 1995, Taylor, 1992), however an alternate assay was developed as a key reagent used in the assay is no longer commercially available.

In carrying out the assay, rabbit anti-human lysozyme antibody (Dako D/K, Denmark) was used to coat a 96 well plate at 1:5000 diluted in PBS overnight at room temperature. After washing with PBS, the plate was blocked with 5% normal donkey serum (Jackson ImmunoResearch Laboratories, PA) in PBS for one hour. The plate
5 was washed again with PBS. Lysozyme samples were diluted in 0.05% Tween in PBS and captured by adding to the plate and incubating for one hour. After washing the plate with PBS, sheep anti-human lysozyme at 1:1000 diluted with 0.05% Tween in PBS was added and incubated for one hour. The plate was washed again with PBS. Peroxidase-conjugated affinipure donkey anti-sheep IgG (H+L) diluted in 0.05% Tween
10 in PBS at 1:10,000 was added and incubated for one hour. After a final wash of the plate with PBS, color was developed by incubating the plate with TMB substrate (Sigma, MO) for 5-15 minutes and the absorbance read at 655 nm.

D. Enzymatic Activity Assay For Lysozyme

15 An indirect sandwich ELISA was developed to quantify total lysozyme expressed in rice calli or cells and used as an alternative assay to determine the lysozyme expression yield. A direct sandwich ELISA for lysozyme quantification has been previously reported (Lollike *et al.*, 1995, Taylor, 1992 #284), however an alternate assay was developed as the alkaline phosphatase (AP) conjugated sheep anti-human
20 lysozyme is no longer commercially available.

In carrying out the assay, rabbit anti-human lysozyme antibody (Dako D/K, Denmark) was used to coat the plate at 1:5000 diluted in PBS overnight at room temperature. After washing with PBS, the plate was blocked with 5% normal donkey serum (Jackson ImmunoResearch Laboratories, PA) in PBS for one hour. The plate
25 was washed again with PBS. Lysozyme samples were then diluted in 0.05% Tween in PBS and captured by incubating for one hour. After washing the plate with PBS, sheep anti-human lysozyme at 1:1000 diluted with 0.05% Tween in PBS was added and incubated for one hour. The plate was washed again with PBS. Peroxidase-conjugated affinipure donkey anti-sheep IgG (H+L) diluted in 0.05% Tween in PBS at 1:10,000 was
30 added and incubated for one hour. After a final wash of the plate with PBS, color was developed by incubating the plate with TMB substrate (Sigma, MO) for 5-15 minutes and Absorbance was read at 655 nm.

E. Enzymatic Activity Assay For Lysozyme

A reliable and quantitative method was developed to analyze the expression level of enzymatically active lysozyme. The turbidimetric assay was developed using a 96-well microtiter plate format and based on the standard lysozyme assay that is carried out spectrophotometrically in cuvettes. A microtiter plate based method previously described for the detection of lysozyme release from human neutrophils had a detection range of 1-100 ng/ml (Moreira-Ludewig and Healy, 1992). The assay conditions were modified to maintain the linearity of detection up to 3.0 µg/ml.

The enzymatic activity of lysozyme was routinely determined by spectrophotometric monitoring of the decrease in turbidity at 450 nm of a suspension of *Micrococcus luteus* (*M. lysodeikticus*) cells (Shugar, 1952). Specifically, 250 µl of a 0.015% (w/v) *Micrococcus luteus* cell suspension was prepared in 66 mM potassium phosphate, pH 6.24 (buffer A). Cell suspensions were equilibrated at room temperature and the reaction was initiated by adding 10 µl samples containing lysozyme with concentrations from 0 to 2.4 µg/ml. Lysozyme activity was determined in a kinetic mode for 5 minutes at 450 nm. The concentration of lysozyme was then calculated by reference to the standard curve constructed with human milk-derived lysozyme.

The enzymatic activity of human milk lysozyme and the rice calli derived lysozyme of the invention was compared. As shown in Figure 3, the lysozyme effected reduction of the turbidity of *Micrococcus luteus* cell suspensions at 450 nm was very similar for lysozyme from the two sources, while buffer alone did not have any effect on the reduction of turbidity.

Three selected suspension cell culture lines were induced to express lysozyme and the yield estimated in parallel by ELISA and the enzymatic activity assay described above (Table 2). T-test analysis showed that there was no significant difference between the lysozyme concentration measured by ELISA and enzymatic activity assay ($p < 0.05$). These results demonstrate that active recombinant human milk lysozyme is synthesized and maintained in rice callus cells and can be isolated without losing its ac

Table 2. Comparison Of Lysozyme Yields Estimates By Enzymatic Activity Assay And ELISA.

Cell line	Lysozyme yield by enzymatic activity assay (lysozyme/total protein $\mu\text{g}/\text{mg}$)	Lysozyme yield by ELISA (lysozyme/total protein $\mu\text{g}/\text{mg}$)
156-5	25.8 \pm 6.3	30.3 \pm 3.9
156-16	32.1 \pm 5.7	32.9 \pm 3.2
156-31	47.0 \pm 6.2	42.3 \pm 7.0

5. F. Recombinant Human Lysozyme has Bactericidal Function

The sensitive lysis of *Micrococcus luteus* cells in a turbidimetric assay (Fig. 3) indicates that recombinant human lysozyme possesses enzymatic activity and functions as a bactericide. To confirm this with a gram-negative bacterium, a bactericidal assay was carried out using an *E.coli* strain (JM109) as a test organism (data not shown).

- 10 In carrying out the assay, an aliquot of overnight JM109 culture was grown in LB medium until mid log phase. A standard inoculum of mid-log phase JM109 at 2×10^5 CFU (colony forming units)/ml was used in the bactericidal assay. Buffer (20 mM Sodium phosphate, pH 7.0, 0.5 mM EDTA) alone, buffer containing human milk lysozyme or rice cell culture derived lysozyme at about 30 $\mu\text{g}/\text{ml}$ were sterilized by
- 15 filtration. The mixture of cells and lysozyme solution was then incubated at 37 C for the specified length of time. One-fifth of the mixture volume was plated onto the LB agar plates and incubated overnight at 37 C in order to determine the number of colony forming units. At the concentration of 30 $\mu\text{g}/\text{ml}$, recombinant human lysozyme exhibited a similar bactericidal effect as lysozyme from human milk. There was no reduction of
- 20 colony forming units using an extract from the non-transgenic control.

G. Purification of Lysozyme From Rice Calli, Suspension Cultures And Transgenic Rice

- Five rice calli lines expressing high levels of lysozyme were propagated and
- 25 induced by sucrose starvation. The calli or cells were ground by a TissueMizer in extraction buffer (PBS, 0.35 M NaCl) at 2 ml buffer/g of wet calli. The resulting tissue homogenate was centrifuged at 25,000 x g for 30 minutes at 4 C. The supernatant was removed and subjected to filtration through a pre-filter and then through a 0.45 μm nitrocellulose filter.

- 30 Approximately 1 liter of filtered supernatant from 500 grams of induced wet calli were then dialyzed against 50 mM sodium phosphate, pH 8.5 at 4 C overnight. The

supernatant was loaded onto a 200 ml SP Sepharose fast flow column (XK26/40, Pharmacia) equilibrated with the loading buffer (50 mM sodium phosphate, pH 8.5) at a flow rate of four ml/min. The column was then washed with the same buffer until a baseline of A280 was achieved. Lysozyme was eluted by 0.2 M NaCl in the loading buffer and fractions containing lysozyme activity were pooled, concentrated and reappplied to a Sephacryl-100 column equilibrated and run with PBS at a flow rate of one ml/min. Proteins were eluted and separated by using PBS at a flow rate of one ml/min. Pure lysozyme fractions were identified by activity assay and total protein assay (Bradford). Finally the purity of lysozyme was confirmed by SDS-PAGE.

The five lines with the highest lysozyme expression level were selected and propagated continuously in petri dishes or shake flasks for lysozyme isolation and purification. A crude extract from rice callus contains both recombinant human lysozyme and large amounts of native rice proteins. Since the calculated pI of lysozyme is approximately 11, a strong cation exchange column, SP-Sepharose fast flow (Pharmacia), was chosen as the first column to separate the rice proteins from recombinant human lysozyme. Most of the rice proteins did not bind to the column when equilibrated with 50 mM sodium phosphate, pH 8.5. The recombinant human lysozyme, on the other hand, bound to the column and was eluted by 0.2 M NaCl. Rice proteins that co-eluted with recombinant human lysozyme, were separated from lysozyme by gel filtration through a Sephacryl S-100 column and highly purified recombinant human lysozyme was obtained.

R2 rice seeds from transgenic plants were dehusked and milled to flour using conventional methods. Lysozyme was extracted by mixing the rice flour with 0.35 N NaCl in PBS at 100 grams/liter at room temperature for one hour. The resulting mixture was subjected to filtration through 3 μ m of a pleated capsule, then through 1.2 μ m of a serum capsule and finally through a Suporcap 50 capsule with a 0.8 μ m glass filter on top of 0.45 μ m filter (Pall, MI).

The clear rice extract (1 liter) was then dialyzed against 50 mM sodium phosphate, pH 8.5 at 4°C overnight and the dialyzed sample was loaded onto a cation exchange resin SP-Sepharose (Pharmacia Amersham), which was pre-conditioned with 50 mM sodium phosphate, pH 8.5 before loading. After loading, the column was washed with the same buffer until a base line A280 reading was achieved, then lysozyme was eluted with 0.2 N NaCl in 50 mM sodium phosphate, pH 8.5. Fractions containing lysozyme were pooled and reappplied to a Sephacryl S-100 column (Bio-Rad; equilibrated and run with PBS). Pure lysozyme fractions were identified by enzymatic

assay and total protein assay (Bradford) and the purity of lysozyme was confirmed by SDS-PAGE.

H. Attributes Of Recombinant Human Lysozyme Produced In Rice

(i). N-Terminal Amino Acid Sequencing

Recombinant human lysozyme (rLys) isolated from mature rice seed as described above, was separated by 18% SDS-PAGE followed by electroblotting to a PVDF membrane (Bio-Rad, CA). The lysozyme band was identified by staining the membrane with 0.1% Coomassie Brilliant Blue R-250 in 40% methanol and 1% glacial acetic acid for 1 minute. The stained PVDF membrane was immediately destained in 50% methanol until the band was clearly visible. After the blot was thoroughly washed with H₂O and air-dried, it was sequenced with a sequencer ABI 477 by Edman degradation chemistry at the Protein Structure Laboratory of the University of California at Davis. The results showed that the rLys produced in transgenic rice seed had an identical N-terminal sequences to the human lysozyme, as follows:

Recombinant Lys-----LysValPheGluArg(.)GluLeuAlaArgThr

Human Lys-----LysValPheGluArgCysGluLeuAlaArgThr

Additionally, a number of structural and functional attributes of human lysozyme and recombinant lysozyme produced in rice were found to be the same, including molecular weight, pI, bactericidal effect with E. coli, thermal and pH stability and specific activity.

(ii). Thermal and pH stability of lysozyme

A human lysozyme standard and lysozyme from rice were diluted to a final concentration of 50 µg/ml in PBS and subjected to the following thermal treatment in a sequential mode: (1): 62 C for 15 minutes; (2): 72 C for 20 seconds; (3): 85 C for 3 minutes and finally; (4): 100 C for 8 seconds. Studies were conducted with 100 µl per tube and repeated three times. Aliquots were saved at the end of each treatment and the remaining lysozyme activity was measured by activity assay.

For studies on pH stability, lysozyme was dissolved in 0.9% NaCl at 100 µg/ml at pH 7.3, 5, 4, 3 and 2. The solutions were incubated at 24°C for one hour. Experiments were conducted with 200 µl per tube and repeated three times. Remaining lysozyme was detected by lysozyme activity assay.

For biotechnological applications of the recombinant human lysozyme, its thermal and pH stability as well as its resistance to proteases is of decisive importance.

In the results presented in Fig 5, recombinant lysozyme exhibited the same degree of thermal stability in the temperature range from 62 C to 100 C as human milk lysozyme. The pH stability of recombinant and native lysozyme was comparable over the range of pH 2-7.3 (Fig. 5).

5

(iii). Determination of *in vitro* protease resistance of lysozyme

Lysozyme was dissolved in 0.9% NaCl at 100 µg/ml. The pH of the solution was reduced to 3, 4 and 5 with HCl. Pepsin (Sigma, MO) (pepsin: lysozyme = 1:22 (w/w)) was added and the solutions were incubated at 37°C for one hour. Then the pH of all treatments was raised to pH 7 with bicarbonate. Pancreatin (Sigma, MO) (pancreatin : lysozyme = 1:110 (w/w)) was added to the neutral solution and incubated at 37 C for two hours. The remaining lysozyme activity was measured by activity assay.

In *in vitro* digestion experiments with pepsin and pancreatin, the native and recombinant human lysozyme displayed very similar resistance to pepsin and pancreatin digestion (Fig. 11). Under these conditions, human albumin was degraded as demonstrated by SDS-PAGE (data not shown).

(iv). Biochemical Characterization of Lysozyme

After recombinant human lysozyme was purified to near homogeneity, several biochemical characterizations were carried out to compare human milk lysozyme with recombinant human milk lysozyme derived from rice cells. The results summarized in Table 3 show that by SDS-PAGE, native human milk lysozyme and recombinant lysozyme migrated to the same position (Fig. 4).

Nucleotides encoding the rice *Ramy3D* signal peptide were attached to the human lysozyme gene in the expression vector pAPI156. Determination of the N-terminal amino acid sequence of the purified recombinant human lysozyme revealed an N-terminal sequence identical with that of native human lysozyme, as detailed above. Rice cells thus cleave the correct peptide bond to remove the *RAmy3D* signal peptide, when it is attached in the human lysozyme precursor.

The overall charge of recombinant and native human lysozyme were compared by isoelectric-focusing (IEF) gel electrophoresis and pI values determined. Since lysozyme is a basic protein with a calculated pI of 10.20, the pI comparison studies were carried out by reverse IEF gel electrophoresis. Recombinant and native human lysozyme displayed identical pI, indicating the same overall charge (data not shown).

Recombinant human lysozyme derived from transgenic rice had a specific activity similar to the native lysozyme (200,000 units/mg (Sigma, MO), whereas, lysozyme from chicken egg whites had the expected 3-4 fold lower specific activity (Sigma, MO).

5

Table 3. Comparison of Biochemical Characteristics of Human Milk Lysozyme and Recombinant Lysozyme

Lysozyme source	N-terminal sequence	Size (kDa)	Glycosylation	Specific activity (units/mg)	pI
Human milk	KVFER C ELART	14	No	201,526	10.2
rice	KVFER(-) ELART	14	No	198,000	10.2

10

The results described above demonstrate the ability to use rice calli or cell suspension cultures as a production system to express human lysozyme from milk. Over 60 individual transformants were screened by immunoblot, enzymatic activity assay and ELISA. Yields of recombinant human milk lysozyme reached 4% of soluble cell proteins. Although the mechanism is not part of the invention, the high expression level may be explained by the utilization of the strong *RAmy3D* promoter (Huang *et al.*, 1993) and the codon-optimized gene.

15

When the gene construct for the human alpha-1-antitrypsin (AAT) precursor containing the rice *RAmy3D* signal peptide was expressed in rice cells, the precursor AAT was processed and secreted and AAT was detected only in the media (Terashima *et al.*, 1999a; Terashima *et al.*, 1999b). In contrast, the majority of recombinant human lysozyme produced by rice calli according to the methods described above was not detected in the culture media, but remained associated with the calli.

20

The plant derived human milk lysozyme obtained by the methods of the present invention was identical to endogenous human lysozyme in electrophoretic mobility, molecular weight, overall surface charges and specific bactericidal activity.

25

EXAMPLE 4

Characterization Of Recombinant Human Lactoferrin (rLF) Produced by Transgenic Rice Plants

A. Southern blot analysis.

5 About three grams of young leaf were collected and ground with liquid nitrogen into a very fine powder. The DNA was isolated according to the procedure as described in Dellaporta SL *et al.*, 1983, and purified by phenol-chloroform extraction. Approximately 5 µg of *EcoRI* and *HindIII* digested DNA from each line was used to make blot for Southern analysis. The ECL™ direct nucleic acid labeling and detection
10 system (Amersham, USA) was used for analysis.

The lactoferrin gene copy number was estimated to be from 2 to 10 as determined by Southern blot hybridization using *EcoRI* and *HindIII* digested genomic DNA. The API164-12-1 (R₀) transgenic plant line was subjected to Southern analysis together with ten Western blot positive, field grown R₁ lines. A typical Southern blot
15 shows that there are at least three fragments above the original plasmid derived plant transformation unit (3156 bp). All the LF inserts appear to be inherited from the original R₀ transgenic plant event (lane 2) to R₁ (lane 3 – 11) generation.

20 B. Protein Isolation and Western blot.

Rice seeds were ground with 1 ml of 0.35 N NaCl in phosphate buffer saline (PBS), pH 7.4 using an ice-cold mortar and pestle and the resulting homogenate was centrifuged at 15000 rpm for 15 min at 4°C. The supernatant was used as a protein extract and about 1/25 or 1/50 of the salt soluble content was loaded onto a 10% pre
25 cast gel (Novex, USA) and electrophoresis was carried according to the manufacturer's instructions. For total protein detection, the polyacrylamide gel was stained with 0.1% Coomassie brilliant blue R-250 (dissolved in 45% methanol and 10% glacial acetic acid) for at least three hours and destained with 45% methanol and 10% glacial acetic acid until the desired background was achieved.

30 For Western blot analysis, SDS-PAGE gels were electroblotted onto a 0.45 µm nitrocellulose membrane with a Mini-Trans-Blot Electrophoretic Transfer Cell System (Bio-Rad, USA) and subsequently subjected to immuno-blotting analysis. The blot was blocked with 5% non-fat dry milk in PBS for at least two hours followed by three washes with PBS for 10 minutes each. The primary rabbit polyclonal antibody against hLF
35 (Daka A/S, Denmark) was diluted at 1:2500 in the blocking buffer and the blot was

incubated in the solution for one hour. The blot was washed with PBS for three times with 10 minutes each. The secondary goat anti-rabbit IgG (H+L)-alkaline phosphatase conjugated (Bio-Rad, USA) was diluted in the blocking buffer at 1:5000 ratio. The membrane was incubated in the secondary antibody solution for one hour and followed by three washes with PBS. Color development was initiated by adding the substrate system BCIP-NBT (Sigma, USA) and the process was stopped by rinsing the blot with H₂O once the desirable intensity of the bands was achieved.

Fifteen R₀ plants were grown to maturity, seeds were harvested from 11 fertile plants and individual seeds analyzed by Western blot to detect the expression of rLF.

Coomassie blue staining was carried out to compare the mobility of rLF with native human lactoferrin (hLF) (Fig. 8), with 40 µg of total protein loaded onto each lane, along with 40 ng of native purified hLF per lane as the positive control.

Estimation of total rLF by ELISA indicated that from 93 µg to 130 µg rLF was expressed in transformed rice seeds. A typical Western blot analysis (Fig. 9) illustrates that both rLF and native hLF migrate at approximately the same rate with the molecular weight about 80 kDa, consistent with that determined by other researchers (Wang *et al.*, 1984).

C. Protein purification.

Rice seeds from R₂ homozygous generation were dehusked and milled to flour conventionally. Recombinant lactoferrin was extracted by mixing the rice flour with 0.35 N NaCl in PBS at 100 g/l at room temperature for two hours. The resulting mixture was centrifuged at 15,000 rpm for one hour at 4°C. The collected supernatant was subjected to the following steps of filtration before loading onto a Sepharose column. First, the supernatant was run through a few layers of cheesecloth. Then the filtrate was passed sequentially through an 8µm paper, 1 µm paper and a 0.25µm nitrocellulose membrane. The clear protein solution was loaded onto a ConA Sepharose column (Pharmacia, XK 26) which had been equilibrated with 0.5 N NaCl in 20 mM Tris, pH 7.4 (binding buffer) at a flow rate at 4 ml/min. After the loading was complete, the column was washed with binding buffer until the baseline at A_{280nm} was achieved. Lactoferrin was eluted with 0.1N mannoside in the binding buffer. Fractions containing lactoferrin were pooled and loaded onto a second column SP-Sepharose (Bio-Rad, USA) which has been equilibrated with 0.4 N NaCl in 50 mM sodium phosphate, pH 8.0 (binding buffer) at the flow rate 4 ml/min. Then the column was washed with the binding buffer until the baseline at A_{280 nm} was obtained. Lactoferrin was eluted by 1 N NaCl in 50 mM sodium

phosphate, pH 8.0 and the fractions containing LF were pooled and dialyzed against PBS. Finally the purity of LF was assessed by SDS-PAGE and stored at - 80°C.

D. Enzyme Linked Immunosorbant Assay (ELISA).

5 ELISA was conducted using seed extracts, isolated as described above, with total protein assayed using the Bradford method (Bradford, 1976). The ELISA was based on a typical sandwich format generally known in the art. Briefly, 96 well plates were coated with rabbit anti-human lactoferrin antibody (Daka A/S, Denmark), then rLF and control samples were added to individual wells of the plate and incubated for 1 hour
10 at 35°C. Rabbit anti-human lactoferrin horseradish peroxidase conjugate (Biodesign, USA) was then added to each well and incubated for 1 hour at 35°C, followed by addition of the tetramethylbenzidine substrate (Sigma, USA) and incubation for 3 minutes at room temperature. The reaction was stopped by adding 1N H₂SO₄ to each well. The plates were read at dual wavelengths of 450 and 650 nm in a Microplate
15 Reader (Bio-Rad, model 3550) and the data was processed by using Microplate Manager III (Bio-Rad). The results of an analysis of 10 homozygous selected lines showed that from 93 µg to 130 µg rLF was expressed per seed.

E. Selection of plants for advance generations.

20 At least 20 - 40 seeds from 11 independent lines were analyzed. Individual R₁ seeds were cut into half and endospermic halves were subjected to analysis by Western blot with the positive corresponding embryonic halves germinated on 3% sucrose medium with 0.7% agar. The seedlings were transplanted to the field for R₁ generation. Out of 11 individual lines, 3 lines were expressed. A total of 38 plants were grown in the
25 field derived from the 3 expressed mother lines. Based on the agronomic character (Table 4) of those 38 plants, 28 plants were selected.

It was observed that all the Western positive R₁ seeds were opaque to pinkish in color in comparison to control seeds, so this criteria was applied in screening the R₂ seeds. Mature R₂ seeds were harvested at maturity and dehusked. The pinkish R₂
30 seeds were confirmed by Western dot blot and ELISA as expressing rLF (data not presented). Finally 10 homozygous R₂ lines were selected and grown in the field in order to advance the generation.

Table 4. Comparison Of Phenotypic Characteristics Of Native TP-309 And Transformed TP-309 Rice Seeds

Source	Effective tiller	Blank grain (%)	1000 seed weight (g)	µg of rLF/seed
TP-309	43	5.0	25	
Homozygous transgenic lines	42	19.7	20.2	125

During R₂ and R₃ generation the percentage of blank seeds was higher in homozygous transgenic lines than in the non-transgenic control. This affected the 1000 seed weight. However, in the R₄ generation no significant differences in phenotypic character were observed in homozygous transgenic lines when compared to non-transformed TP309 (Table 4).

F. Attributes Of Recombinant Human Lactoferrin Produced In Rice

Physical characterization of the rLF showed there was no significant difference between the rLF and a commercially available purified form of hLF based on N-terminal amino acid sequencing, and physical characteristics of rLF such as molecular weight as determined by MALDI-MS, HPLC profile of which showed a comparable peptide map, pH dependent iron release and bacteriostatic activity, using the analyses described below.

(i). N-Terminal Amino Acid Sequencing. Purified rLF from rice seeds was resolved by 10% SDS-PAGE, followed by electroblotting to PVDF membrane (Bio-Rad, USA). The target band was identified by staining the membrane with 0.1% Coomassie brilliant blue R-250 in 40% methanol and 1% glacial acetic acid for 1 minute. The stained PVDF membrane was immediately destained in 50% methanol until the band is clearly visible. The blot was thoroughly washed with ddH₂O and air dried. Finally this sample was sent to the Protein Structure Laboratory in University of California at Davis (CA, USA) for sequencing analysis.

(ii). Detection of glycosylation and determination of sugar content

Glycosylation of the recombinant human lactoferrin produced in rice was analyzed by an immunoblot kit for glycoprotein detection (Bio-Rad, USA) per instructions from the manufacturer. An increase of molecular weight of lactoferrin due to carbohydrate content was determined by Matrix Assisted Laser Desorption Ionization-Mass spectrometry (MALDI-MS) (PE Applied Biosystems, Voyager System).

Recombinant lactoferrin produced in rice is glycosylated as evident from the binding to Con A resin, the positive staining by glycoprotein detection kit as well as the larger detected mass as compared to the calculated mass (76.2 kDa) based on the peptide backbone. MALDI-MS showed that seed derived recombinant lactoferrin has molecular weight of 78.5 kD while human milk lactoferrin is 80.6 kDa (Table 5). The difference could be due to the lesser degree of glycosylation in the rice seed-derived lactoferrin.

(iii). Determination of isoelectric point of Lactoferrin.

Reverse isoelectric focusing (IEF) gel electrophoresis was carried out with a precast Novex IEF gel, pH 3-10 according to the manufacturer's instruction. About 30 µg of purified rLF was loaded and the running condition was 100 V for 50 minutes and 200 V for 20 minutes. The gel was then fixed in 136 mM sulphosalicylic acid and 11.5% TCA for 30 minutes, stained in 0.1% Coomassie brilliant blue R-250, 40% ethanol, 10% glacial acetic acid for 30 minutes and destained in a solution containing 25% ethanol and 8% acetic acid.

(iv). Comparison of physical characteristics of rLF with native hLF.

The HPLC profile of native and rLF showed a comparable peptide map. This confirmed that LF from the two sources have an identical amino acid sequence (data not presented). Additional comparisons confirm that human lactoferrin produced in transgenic rice closely resembles native human lactoferrin, as evidenced by (1) the N-terminal sequence of purified rLF from homozygous R₂ seeds and hLF (Dakao A/S, Denmark), which were shown to be identical (Table 5); (2) the isoelectric point (pI) of native and rice seed derived LF which is the same, indicating that they have similar surface charges (Table 5); (3) the pH dependent iron release of rLF which was shown to be closely related to that of native hLF (Fig. 10); and (4) the bacteriostatic activity of rHLf which was shown to be similar to that of native human lactoferrin (nHLf) on enteropathogenic *E. coli* (EPEC; Fig. 11) and confirmed the presence of active recombinant LF in extracts derived from transformed rice seeds.

Table 5. Physical characterization data for human (hLF) and rice seed derived recombinant lactoferrin (rLF)

LF source	Size (kDa)	N-terminal sequence	pI	Glycosylated	Sugar content (%)
hLF	80.6	GlyArgArgArgArgSerValGlnTrpCysAla	8.2	YES	5.5
rLF	78.5	GlyArgArgArgArgSerValGlnTrp()Ala	8.2	YES	2.9

5 (v). Iron content and nutrient value determination of rice seeds

The iron content of R₂ homozygous seeds was determined. Two grams of dry mature seeds from each transformed and non transformed line were weighed and wet-ashed with HNO₃ and H₂O₂ solution at 110°C (Goto *et al.*, 1999). The ash was dissolved in 1N HCl solution. The iron content was then measured by absorbance of Fe-O-phenanthroline at 510 nm, using a Sigma kit (Sigma, USA) per instructions of manufacturer.

The different values of nutrient facts of homozygous transgenic seeds and non transgenic seeds were measured by standard procedure at A & L Western Agricultural Laboratories (Modesto, CA, USA).

15 The different values of nutrient facts of homozygous transgenic seeds and non transgenic seeds were measured by standard procedure at A & L Western Agricultural Laboratories (Modesto, CA, USA).

A comparative analysis of transgenic lactoferrin-expressing rice seeds with non transformed native Teiwei-309 showed that there is no significant difference between transformed and non transformed seeds in nutrient value with the exception that the concentration of iron is 50% greater (Table 6). The increased level of iron may be the reason for the opaqueness and pink coloration of the rLF expressing transgenic rice seeds.

25 There was no difference noticed during the seed germination of transgenic seeds, the phenotype of R₂ R₃ and R₄ plants was vigorous and the seed yield was similar to that of non-transgenic Teiwei-309 plants (data not shown).

Table 6. Comparison of Nutrition Value (in mg) Per 100 Gram of Non Transformed and Transformed Rice Seeds

Source	Carbohydrate	Protein	Fat	Ca	K	Na	Fe	Water	Calories
TP-309	76.0	8.7	2.4	9	370	<10	0.8	11.3	369
Homozygous transformed lines	75.7	8.7	2.2	8	330		1.2	11.8	367

(vi). Tissue specificity and stability of rLF

5 An endosperm specific rice glutelin promoter was used to express recombinant lactoferrin in maturing or matured seeds. To confirm the tissue specificity of the expressed lactoferrin, protein was extracted from root, shoot, leaf beside mature seed and subjected to Western blot and the results indicated that there was no detectable expression of rLF except in the seed/endosperm (results not shown). Furthermore, the
10 presence of rLF in 5 day old germinated seeds showed the stability of stored rLF within the plant cell during germination.

(vii). Iron Saturation And pH Dependent Iron Release

Lactoferrin was incubated with 2M excess ferric iron (FeCl_3 :NTA = 1:4) and
15 sodium bicarbonate ($\text{Fe}:\text{HCO}_3^- = 1:1$) for 2h at room temperature. Excess free iron was removed by using a PD-10 desalting column (Pharmacia, USA) and the iron saturation level was determined by the A_{280}/A_{456} ratio. Both native hLF and rLF were completely saturated by iron. Holo hLF was incubated in buffers with a pH between 2 and 7.4, at room temperature for 24 h. Free iron released from hLF was removed and the iron
20 saturation level was determined by A_{280}/A_{456} ratio.

The results showed that iron release was similar for both hLF and rLF. Iron release began around pH 4 and was completed around pH 2 (Fig. 10). The iron binding was reversible since iron-desaturated rLF was re-saturated by raising the pH to 7 (data not shown). The similarity in pH dependent iron release of rLF to that of the hLF
25 standard demonstrated that rLF is able to adapt the appropriate tertiary structure for proper iron binding and release (Salmon, Legrand *et al.* 1997).

(viii). Antimicrobial Activity: Effect Of *In vitro* Digestion

Lactoferrin is known to inhibit the growth of a variety of bacterial species based
30 on its iron chelation and direct bactericidal properties. The anti-microbial effect of rLF extracted from rice seeds was tested following treatment using an *in vitro* digestion

model with an enzymatic system containing pepsin (an enzyme active in stomach) and pancreatin (an enzyme active in deodenum).

LF proteins were dissolved in PBS at 1mg/ml, and either left untreated, pepsin treated (0.08mg/ml at 37°C for 30 min), or pepsin/pancreatin treated (0.016 mg/ml at 37°C for 30 min). LF proteins were sterilized by passing through a membrane filter with a pore size of 0.2 μ m [Rudloff, 1992]. The filter sterilized LF (0.5 μ g/ml) was incubated with 10⁴ colony forming unit (CFU) enteropathogenic *E. coli* (EPEC)/ μ l in 100 μ l sterile synthetic broth (1.7% : AOAC) containing 0.1% dextrose and 0.4 ppm ferrous sulfate at 37°C for 12h and colony forming units (CFU) were determined.

Starting with an enteropathogenic *E. coli* (EPEC) concentration of 10⁴ CFU (colony forming units), the untreated samples of rLF reached up to 10^{6.5} CFU after 12 h of incubation at 37°C in comparison to hLF, which produced up to 10⁶ CFU. An *in vitro* digestion model using an enzymatic system containing pepsin (enzyme active in stomach) and pancreatin (enzyme active in deodenum) with moderate shaking to imitate the transit of protein through infant gut [Rudloff, 1992] was used. rLf and nHLf were treated with active pepsin and pancreatic enzymes and exposed to 10⁴ CFU EPEC cells for 12 h at 37°C (Fig. 11). Both the native human lactoferrin standard (nHLf) and the recombinant rice-derived lactoferrin (rLf) remained active in inhibiting growth of enteropathogenic *E. coli*, indicating that both nHLf and rHLf are resistant to protease digestion.

EXAMPLE 5

Resistance of the Recombinant Lysozyme and Lactoferrin to Intestinal Digestion *In vivo*

The established protective actions of lysozyme and lactoferrin require that they survive in a biologically active form following passage through the stomach and into the small intestine and both human lysozyme and lactoferrin are known to possess sufficient protection against denaturation and hydrolysis to be efficacious in humans. The pH gradients and the digestive enzymes in chickens and humans are similar suggesting that the molecules should survive passage through the chicken's digestive tract. In order to evaluate the resistance of recombinant rice-derived lysozyme and lactoferrin to the digestive process of humans or chickens, the survival of lysozyme and lactoferrin at 2 ages: 3 days of age and 21 days of age was evaluated. At 3 days of age, the digestive tract of chicks is immature and levels of digestive enzymes are low. At 21 days, the rate of digestive enzyme production is at adult levels and digestibility of

proteins is maximal; providing a maximal challenge to the survival of the target molecules.

At 21 days of age, one chick per pen is killed and the contents of their intestines removed. IN carrying out the analysis, segments of approximately 1.5 cm in length are
5 flushed with saline and fixed in 10% buffered formalin (pH 7.0), embedded with paraffin, sectioned (5 mm), stained with hematoxylin-eosin, and mounted. This procedure is performed by a commercial laboratory (California Veterinary Services, West Sacramento California). The histological sections are evaluated for: thickness of the lamina propria; villus height- from the base of the lamina propria to the apex of the villus;
10 crypt depth between adjacent villi. Morphometric data is collected on 10 different villi per animal on each of the two different serial sections. Measurements are made and analyzed by computer-aided light microscope analysis at magnifications between 10 to 1000x using Image-Pro-Plus analysis software for the PC.

Separate studies may be carried out to determine the survivability of lysozymes
15 and lactoferrin in the digestive tract of chickens.

In carrying out the study, on the day of the test, food is removed for 4 hours to insure that a 20g aliquot of the experimental diets are consumed with 10 minutes. At 15, 20, 40 and 60 minutes after consumption of experimental diets, 4 chicks per treatment are killed. The gastrointestinal tract is removed and divided into 6 sections:
20 proventriculus, gizzard, duodenum, jejunum, ileum, and hindgut. Each section is flushed with 1 volume of physiological saline and the contents collected into sterile plastic tubes.

Following removal of solids by centrifugation, soluble proteins can be tested for activity. In case of lysozyme, the test is a measurement of hydrolysis of *Micrococcus lysodeikticus* (Sigma Chemical Company) using chicken lysozyme as a standard. In
25 the case of lactoferrin, the material is tested using published protocols for determining lactoferrin binding to CACO-2 cells. Human lactoferrin (Sigma Chemical Company) is used as a standard. Chromium marker concentration can be determined by flame atomic absorption spectrophotometry, with concentrations expressed as µg/µg of
30 chromium.

EXAMPLE 6

Effect of Lysozyme Supplemented Feed on the Feed Utilization and Growth of Chickens

Various feed samples were prepared to test the effect of transgenic rice-derived
35 rLys, supplemented animal feed on feed utilization and animal growth (Table 7).

Table 7: Composition of Feed Samples

Diet #	Type	Components
1	negative control	A standard corn-based diet containing no antibiotic supplements
2	negative control	A corn-based diet containing no antibiotics, but supplemented with 15% non-transgenic control rice.
3	positive control	A corn-based diet containing a standard sub-therapeutic level of antibiotics (Bacitracin + Roxarsone)
4	Test diet	A non-antibiotic diet, substituted with 10% transgenic (TG) rice containing rLys to give a final lysozyme level of 0.03%
5	Test diet	A non-antibiotic diet, substituted with 10% transgenic rice containing rLys and 5% transgenic rice containing rLac to give final lysozyme and lactoferrin levels of 0.03% and 0.02% respectively

5 rLys concentrations are based on 2 g/kg lysozyme in the transgenic rice and each of the experimental diets also contains a non-absorbable marker (chromium) in order to follow passage of the non-absorbed digesta.

The presence of lysozyme in each of the feed compositions was confirmed by Western blot.

10 Chickens (male Cobb broiler chickens; n=48) were individually housed and fed the control diet *ad libitum* prior to being given a test diet at 3 days and 3 weeks of age. The control diet (Diet #1) is a standard reference chicken diet recommended by the National Academy of Science, with the exception that 10% ground rice is used in place of ground corn.

15 On the day of hatching, two hundred forty chickens (male Cobb broiler chickens) were assigned to each of the 5 treatment groups with 8 pens per treatment and 5 chicks per pen. Beginning on the first day post-hatch, chickens received experimental diets and water *ad libitum*. Chickens were weighed and feed intake recorded on days 1, 5, 10, 15, 20, 25, 30 and 35.

20 The cages used for the study had previously been used for 3 production cycles of chickens. They were not cleaned or disinfected between cycles of chicken production, permitting the build up of feces, dust, and dander. The rate of air-turnover in the room housing the chickens was reduced to levels similar to that seen in a commercial broiler production barn. These sanitation practices were intended to mimic commercial conditions where antibiotics are proven to provide increased growth and
25 feed conversion efficiency.

The results demonstrate that feeding growing chickens a diet containing recombinant lysozyme results in a feed conversion efficiency that is better than sub-therapeutic antibiotics. As shown in Table 8, the growth rate of chickens as measured in grams of weight gained per chick per day is comparable, even at the level of just 0.2% added transgenic grain, to that seen with antibiotic or chicken lysozyme controls. Similarly, growth as a factor of feed efficiency, or grams of weight gained per gram of feed, was on the order of that seen in antibiotic and chicken lysozyme controls.

Table 8: Weight and Feed Intake and Feed Efficiency of Chickens Fed Various Diets

Diet #	Diet composition	Gain (g/chick/day)	Feed Consumed	Feed Efficiency
		Day 0-21	Day 0-21	Day 0-21
1	Standard Corn-based Diet	33.82	51.08	0.732
2	Standard Diet with 15% rice	38.03	52.18	0.72
3	Standard Diet + antibiotics	34.43	50.48	0.754
4	Standard Diet + 10% TG rice containing rLys	33.53	48.5	0.766
5	Standard Diet + 10% TG rice containing rLys and 5% TG rice containing rLac	34.03	49.02	0.766

EXAMPLE 7

Effect of Supplemented Feed on Intestinal tract of Chickens

Two studies conducted to demonstrate that rice that has been genetically produced to express human lactoferrin (LF) or lysozyme (LZ) protects the intestinal tract similar to sub-therapeutic antibiotics.

A. Expression of Lactoferrin or Lysozyme in Rice

Two transgenic rice strains were produced to express either lactoferrin or lysozyme (Huang et al., 2002). Briefly, rice callus from the rice strain Taipei 309 was transformed with plasmids carrying genes for LF and LZ under the control of rice glutelin 1 gene promoter. Transgenic plants were screened for high level of expression of both recombinant proteins. Selected lines, 159-53 and 164-12 were propagated to produce sufficient amount of rice seed for these experiments. LF and LZ rice expressed 2.5 and 4.0 g/kg of recombinant protein as determined by ELISA. Taipei 309, which is the conventional rice (CONV) that served as the host for transgenic plant production, served

as a control. All rice was dehusked to yield brown rice and then ground using a comminuting machine (Fitzpatrick Inc, Chicago, IL).

B. Preparation and Management of poultry

5 One-day old male Cobb broiler chicks (Foster Farms, Delhi, CA) were raised in Petersime brooder batteries (Petersime Incubator Co., Gettysburg, OH) located in an environmentally controlled room (25 C) with 24 hrs of light. Chicks were provided water and commercial chick starter for ad libitum consumption. The batteries had not been
10 cleaned after their previous use in order to provide a level of sanitation conducive for an antibiotic response. When the chicks were 3 days of age, experimental chicks were selected for uniform body weight from a two-fold larger population and randomly assigned to dietary treatments. Chicks had ad libitum access to both feed and water and were exposed to a 24-hour light cycle. All experiments and procedures were approved by the Campus Animal Care and Use Committee.

15

C. Diet Formulation

Corn-soy-rice basal diets were formulated to meet or exceed the nutrient needs of young growing broiler chicks suggested by NRC (1994). All experimental diets were formulated to contain the same amount of rice by substituting transgenic rice for CONV
20 rice as shown in Tables 1 and 2. A range of levels of each test rice was chosen for study in order to determine a minimally efficacious level. The 10.0% LZ diet used in study 1 was analyzed to contain 176 mg/kg lysozyme. Following 6 months of storage at room temperature, it contained 152 mg/kg lysozyme, indicating that this protein was stable to storage.

Table 9: Composition of Diets from Studies 1 and 2

Ingredient, g/kg	Study 1 Corn-soy-rice	Study 2 Corn-soy-rice	Study 2 Corn-soy
Corn	354	408	551
Soy meal (48.5%)	320	343	342
Rice ¹	200	150	---
Poultry Grease	57.9	43.1	51.0
Meat with Bone Meal	40.5	---	---
Feather Meal	---	21.2	21.2
Calcium carbonate	---	7.6	8.0
Tricalcium phosphate	---	19.1	18.7
Calcium phosphate	18.5	---	---
Vitamin-mineral premix ²	1.00	1.00	1.00
Choline	0.70	0.84	0.91
Sodium chloride	4.03	2.21	2.04
D-L Methionine	2.78	2.03	1.97
L-Lysine	---	---	0.14
ME, kcal/kg	3,199	3,201	3,201
Crude protein, %	22.15	22.29	22.29
Crude fat, %	8.07	6.69	7.85
Available Lys, %	1.20	1.15	1.15
Available Met + Cys, %	0.95	0.88	0.88

¹ Three types of rice (conventional rice, or rice expressing lactoferrin or lysozyme) at variable levels depending upon the treatment.

- 5 ² Vitamins and minerals were provided in the form and level described in NRC (1994) Standard Reference Diet for Chicks.
Corn-soy diet for study 2 only.

Table 10: Types of Rice in Dietary Treatments Used in Studies 1 and 2

Dietary Treatment	% of the Diet	
	Conventional Rice	Transgenic Rice
Study 1		
CONV	20	0
0.1% LF	19.9	0.1
1.0% LF	19.0	1.0
5.0% LF	15.0	5.0
Human lactoferrin ¹	20	0
0.2% LZ	19.8	0.2
10.0% LZ	10.0	10.0
0.1% LF + 0.2% LZ	19.7	0.3
5.0% LF + 10% LZ	5.0	15.0
Antibiotics ²	20	0
Study 2		
Corn-soy	0	0
CONV	15	0
10% LZ	5.0	10.0
5.0% LF + 10% LZ	0	15.0
Antibiotics	15	0

¹ Incorporated at 0.2 g/kg diet.

² Antibiotic treatments contained roxarsone (3-nitro-4-hydroxyphenylarsonic acid; Pfzier, Inc. New York, NY) and bactiracin methylene disalicylate (Schering-Plough Animal Health, Omaha, NE 68127) at 0.25 g/kg diet and 0.5 g/kg diet, respectively.

5

D. Study 1

Study 1 compared 10 corn-soy diets containing 20% of various LF, LZ or conventional rice (CONV). 300 3-day old chicks were randomly assigned to one of 10 dietary treatments as shown in Table 2. Each dietary treatment consisted of six
10 replicates, with five chicks per replicate. Chick and feeder weights were determined on day 1 and 17.

Intestinal samples were taken on the last day of study 1 and 2. Sections (2.5 cm) from one chick per replicate were obtained from the duodenum at the apex of the pancreas, the jejunum at a position midway between Meckel's diverticulum and the
15 entrance of the bile ducts, the ileum at a position midway between Meckel's diverticulum and the ileum-cecal junction, and the ceca at a point midway along its length (study 2 only). Samples were flushed with saline, fixed in 10% buffered formalin (pH 7.0), embedded with paraffin, thin sectioned and stained with hematoxylin-eosin (IDEXX Veterinary Services, Inc., Sacramento, CA). For enumeration of intraepithelial and
20 lamina propria leukocytes sections were fixed in acetone, re-dried, incubated with mouse anti-chicken CD45 monoclonal antibody (Southern Biochemical Associates, Birmingham, Alabama) for 1 hour and then rinsed in PBS.

Sections were incubated with rabbit anti-mouse Ig tagged with peroxidase with 0.5% bovine serum albumin for 1 hour and rinsed. Peroxidase activity was developed by
25 incubating sections with 0.01% H₂O₂ and 3,3'-diamino-benzidine-tetrahydrachloride. The slides were counterstained with hematoxylin-eosin. The number of leukocytes in 10 villi per section and the number of leukocytes in the lamina propria underneath and within these 10 villi were enumerated. Cells with endogenous peroxidase activity (primarily heterophils) were also enumerated as described by Vervelde and Jeurissen (Vervelde,
30 L. and Jeurissen, S.H., 1993). For each intestinal sample, villi height, villi width, crypt depth, lamina propria thickness, number of lamina propria leukocytes, and number of intra-epithelial leukocytes were estimated using Image-Pro-Plus software (Media Cybernetics, Silver Spring, MD).

Data were analyzed for main effect of diet using the general linear model (Minitab; State College, PA). When main effects were significant ($P < 0.05$), differences due to dietary treatment were determined using Tukey's means comparisons.

Chicks fed diets containing 0.1% LF, 1% LF, 5% LF, human lactoferrin, 0.2% LZ, 10% LZ, or 0.1% LF + 0.2% LZ did not differ from chicks fed CONV in any of the parameters measured. Feed intake and body weight gain were not affected by dietary treatments ($P > 0.05$), and averaged 36.10 g/chick/day and 28.96 g/chick/day, respectively. Chicks fed 5% LF + 10% LZ had significantly greater feed conversion compared to chicks fed CONV (Table 3). Chicks fed Antibiotic also tended ($P = 0.058$) to have greater feed conversion compared to those fed CONV.

Table 11: Effect of Dietary Treatment on Feed Efficiency (Study 1)

Feed Efficiency (g body weight gain / g feed consumed)	
Dietary Treatment	
CONV	0.79 ± 0.1^a
5% LF + 10% LZ	0.84 ± 0.1^b
Antibiotics	0.82 ± 0.1^c

Values are means \pm SEM. Means in a column not sharing common superscripts are significantly different ($P < 0.05$).

Histological characteristics of the duodenum, jejunum, and ileum are presented in Table 4.

Table 12: Effect of Dietary Treatment on Intestine Histology (Study 1)

	Lamina propria thickness (μm)	Intra-epithelial leukocytes (#/villi)	Lamina propria leukocytes (#/villi)
Duodenum			
CONV	101 ^b	4.8	34.3
5% LF + 10% LZ	82 ^a	4.0	24.5
Antibiotic	85 ^{ab}	4.75	25.5
SEM ¹	3.06	0.73	3.35
P value	0.015	0.543	0.196
Jejunum			
CONV	101 ^b	8.0	38 ^b
5% LF + 10% LZ	97 ^{ab}	5.0	21 ^{ab}
Antibiotic	84 ^a	2.5	14 ^a
SEM ¹	3.26	1.06	3.67
P value	0.011	0.061	0.008
Ileum			
CONV	105	0.38	34
5% LF + 10% LZ	86	0.34	17
Antibiotic	83	0.39	18
SEM ¹	6.5	0.03	3.67
P value	0.111	0.494	0.068

¹Values are means of 6 observations.

Means in a column within intestinal segment not sharing common subscripts are significantly different ($P < 0.05$).

There were no significant differences in villus height, villus width, or crypt depth due to dietary treatments in any intestinal segment (data not shown). Chicks fed 5% LF + 10% LZ had significantly thinner lamina propria in the duodenum compared to those fed CONV ($P < 0.05$). Chicks fed Antibiotic had jejuni with significantly thinner lamina propria and lower counts of lamina propria leukocytes compared to chicks fed CONV ($P < 0.05$). Chicks fed 5% LF + 10% LZ or Antibiotics tended ($P = 0.068$) to have lower counts of lamina propria leukocytes in the ileum compared to those fed CONV.

E. Study 2

A second study was designed to confirm the results of study 1 using twice the number of replicates per treatment in order to examine more subtle effects of treatments. Study 2 compared 5 corn-soy diets containing experimental rice combinations totaling 155 rice. 360 3-day old chicks were randomly assigned to one of 5 dietary treatments (Table 2) with 12 replicates per treatment and 6 chicks per replicate (42 chicks per treatment). Chick and feeder weights were determined on day 1 and 19. Chicks fed the corn soy diet, which was devoid of rice, did not differ from chicks fed CONV in any of the parameters measured. There was no significant difference ($P > 0.05$) in body weight gain due to dietary treatments and averaged 37.89 g/chick/day. Chicks fed 5% LF + 10% LZ or 10% LZ had significantly lower feed intake compared to those fed CONV (Table 5). Chicks fed either 10% LZ, 5% LF + 10% LZ, or Antibiotics had greater feed efficiency compared to those chicks fed CONV (Table 5). As in study 1, there were no differences in body weight gain, food intake, or feed efficiency between chicks fed LF or LZ rice and those fed Antibiotics.

Table 13: Effect of Dietary Treatment on Food Intake and Feed Efficiency (Study 2)

Dietary Treatment	Food Intake (g/chick/day)	Feed Efficiency (g body weight gain/g feed consumed)
CONV	52.18 ± 1.09^b	0.72 ± 0.01^a
10% LZ	48.05 ± 1.11^a	0.77 ± 0.01^b
5% LF + 10% LZ	49.02 ± 1.23^a	0.77 ± 0.01^b
Antibiotic	50.48 ± 1.15^{ab}	0.75 ± 0.01^b

Values are means \pm SEM. Means in a column not sharing common superscripts are significantly different ($P < 0.05$).

Chicks fed 10% LZ, 5% LF + 10% LZ, or Antibiotics had significantly greater villus height in the duodenum compared to chicks fed CONV as shown in Table 6.

Chicks fed CS + 10% LZ or Antibiotics had significantly thinner lamina propria in the ileum and fewer leukocytes in the ileal lamina propria compared to chicks fed CONV. There were no other significant differences due to diet in villus width, crypt depth, or intra-epithelial leukocytes in any intestinal segment.

5

Table 14: Effect of Dietary Treatment on Intestine Histology (Study 2)

	Lamina propria thickness (μm)	Villus Height (μm)	Lamina propria leukocytes (#/villi)
Duodenum			
CONV	81	743 ^a	24
5% LF + 10% LZ	77	884 ^b	21.5
Antibiotic	82	882 ^b	23.3
SEM ¹	2.46	18.3	2.1
P value	0.69	0.016	0.93
Ileum			
CONV	99 ^b	365	28.3 ^b
5% LF + 10% LZ	88 ^{ab}	421	25.2 ^{ab}
Antibiotic	75 ^a	356	18.7 ^a
SEM ¹	3.2	12.6	1.5
P value	0.03	0.36	0.02b

¹ Values are means of 12 observations.

Means in a column not sharing common superscripts are significantly different (P,0.05).

The results described above demonstrate that the combination of 5% LF + 10% LZ was more efficacious at improving feed efficiency and histological indices of intestinal health than 10% LZ alone. LF alone was without effect. These results demonstrate the potential of rice expressing lactoferrin and lysozyme to serve as an alternative to antibiotics in broiler diets.

10

Table 15. Brief Description of the Sequences

Description	SEQ ID NO
<p>Codon optimized lysozyme coding sequence: AAAGTCTTCGAGCGGTGCGAGCTGGCCCGCACGCTCAAGCGGCTCGGCAT GGACGGCTACCGGGGCATCAGCCTCGCCAACTGGATGTGCCTCGCCAAAGT GGGAGTCGGGCTACAACACCCGCGCAACCAACTACAACGCCGGCGACCG CTCCACCGACTAGGGCATCTTCCAGATCAACTCCCGCTACTGGTGCAACGA CGGCAAGACGCCCGGGGGCCGTCAACGCCTGCCACCTCTCCTGCTCGGCC CTGCTGCAAGACAACATCGCCGACGCCGTGCGGTGCGCGAAGCGCGTCGT CCGCGACCCGCGAGGGCATCCGGGCGCTGGGTGGCCTGGCGCAACCGCTGC CAGAACCGGGACGTGCGCCAGTACGTCCAGGGCTGCGGCGTCTGA</p>	1
<p>Amino acid sequence based on codon optimized lysozyme coding sequence: KVFERCELARTLKRLGMDGYRGISLANWMCLAKWESGYNTRATNYNAGDRST DYGIFQINSRYWCNDGKTPGAVNACHLSCSALLQDNIADAVACAKRVVRDPQG IRAWVAWRNRCQNRDVRQYVQCGCV</p>	2
<p>Codon optimized lactoferrin coding sequence:GGGCGGCGGCGGCGCTCGGTGCAAGTGGTGGCGCCGTGTCCCAG CCCGAGGCGACCAAGTGCTTCCAGTGGCAGCGCAACATGCGGAAGGTGC GCGGCCCCGCGGTGAGCTGCATCAAGCGGGACTCCCCCATCCAATGCATC CAGGCCATCGCGGAGAACCGCGCCGACGCGGTACCCCTGGACGGCGGGT TCATCTACGAGCGGGGCTCGCCCCGTACAAGCTCCGCCCGGTGGCGGC GGAGGTGTACGGCACCGAGCGCCAGCCGCGCACGCACTACTACGCGGTG GCCGTGCTCAAGAAGGGCGGGTCTTCCAGCTCAACGAGCTGCAGGGCCT GAAGTCGTGCCACACGGGCTCCGGCGGACGGCGGGCTGGAACGTGCCC ATCGGCACCCTGCGCCCCCTTCTGAAGTGGACCGGCCCGCCGGAGCCGAT CGAGGCCCGCGTGGCCCCGCTTCTTACGCGCCTCCTGCGTCCCCGGCGCC GACAAGGGCCAGTTCCCGAACCTCTGCCGGCTCTGCGCCGGGACGGGCG AGAACAAGTGCGCCTTCTCCTCGCAGGAGCCGTACTTCTCCTACTCGGGC GCGTTCAAGTGCCCTCCGCGACGGGGCGGCGACGTGGCGTTCATCCGCG AGTCCACCCTGTTTCGAGGACCTCTCCGACGAGGCGGAGCGGGACGAGTAC GAGCTGCTGTGCCCCGACAACACCCGCAAGCCGGTGGACAAGTTCAAGGA CTGCCACCTGGCGCGGGTGCCCTCGCACGCGGTGCTCGCCCGCAGCGTC AACGGCAAGGAGGACGCGATCTGGAACCTCCTCCGCCAGGCCAGGAGAA GTTCCGGCAAGGACAAGTCCCCCAAGTTCCAGCTCTTCGGGAGCCCCAGCG GCCAGAAGGACCTCCTCTTCAAGGACTCCGCGATCGGCTTCTCCCGCGTC CCCCCGCGCATCGACTCCGGCCTGTACCTCGGCTCCGGGTACTTCACCGC GATCCAGAACCTCCGGAAGAGCGAGGAGGAGGTGGCGGCGCGGCGGGGCC CGCGTCGTGTGGTGCGCCGTGGGCGAGCAGGAGCTGCGGAAGTGCAACC AGTGAGCGGGCCTGAGCGAGGGGTGCGGTGACCTGCTCGTCCGCCAGCAC CACCGAGGACTGCATCGCGCTCGTCTCAAGGGGGAGGCCGACGCGATG AGCCTCGACGGGGGGTACGTCTACACCGCCGGCAAGTGCGGCCTGGTCC CGGTCTTGCGGAGAACTACAAGTCGCAGCAGTCCAGCGACCCCGACCCG AACTGCGTGGACCGCCCCGTGAGGGGTACCTCGCCGTGGCCGTGCTGC GCCGGTCCGACACCTCCCTGACGTGGAACAGCGTCAAGGGCAAGAAGAGC TGCCACACCGCCGTGGACCGCACCGCCGGCTGGAACATCCCGATGGGCC TCCTCTTCAACCAGACCGGCTCCTGCAAGTTCGACGAGTACTTCTCCAGT CCTGCGCCCCCGGCTCGGACCCCCGCTCCAACCTGTGCGCCCTCTGCATC GGGACGAGCAGGGCGAGAACAAAGTCGCTGCCAACAGCAACGAGCGGT ACTACGGCTACACGGGGGCTTCCGCTGCTGCGCGGAGAACGCCGGGA CGTCGCGTTCTGTAAGGACGTGACCGTGCTGCAAAACACGACGGGAACA ACAACGAGGCGTGGGCGAAGGACCTCAAGCTCGCCGACTTCGCCCTGCTG TGCCTCGACGGCAAGCGCAAGCCCGTCACCGAGGCGCGGTCTGCCACC TGGCGATGGCCCCCAACCACGCCGTGCTCTCCCGCATGGACAAGGTGAG CGCCTCAAGCAGGTGCTCCTGCACCAGCAGGCCAAGTTCGGCCGGAACGG</p>	3

CAGCGACTGCCCGGACAAGTTCTGCCTGTTCCAGTCGGAGACCAAGAACC TCCTCTTCAACGACAACACCGAGTGCCTGGCGCGCCTCCACGGCAAGACC ACCTACGAGAAGTACCTCGGCCCCGAGTACGTCGCCGGCATCACCAACCT CAAGAAGTGCTCCACCTCCCCCCTCCTGGAGGCGTGCGAGTTCCTCCGCA AGTGA	
Amino acid sequence based on codon optimized lactoferrin coding sequence: GRRRRSVQWCAVSQPEATKCFQWQRNMRKVRGPPVSCIKRDSPIQCIQIAIE NRADAVTLGGFIYEAGLAPYKLRPVAAEVYGTERRQPRTHYYAVAVVKKGGSF QLNELQGLKSCHTGLRRTAGWNVPIGTLRPFLNWTGPPEPIEAAVARFFSASC VPGADKGQFPNLCRLCAGTGENKCAFSSQEPYFSYSGAFKCLRDGAGDVAFI RESTVFEDLSDEAERDEYELLCPDNTRKPVDFKDKCHLARVPSHAVVARSVNG KEDAIWNLLRQAQEKFGKDKSPKFQLFGSPSGQKDLLFKDSAIGFSRVPPRIDS GLYLGSGYFTAQNLRKSEEEVAARRARVWCAVGEQELRKCNCQWSGLSEGS VTCSSASTTEDCIALVLKGEADAMSLDGGYVYTAGKCGLVPVLAENYKSQQSS DPDPNCVDRPVEGYLAVAVRRSDTSLTWNSVKGKKSCHTAVDRTAGWNIPM GLLFNQTGSCKFDEYFSQSCAPGSDPRSNLCALCIGDEQGENKCVPSNERY YGYTGAFRCLAENAGDVAFVKDVTVLQNTDGNNEAWAKDLKLADFALLCLD GKRKPVTEARSCHLAMAPNHAVVSRMDKVERLKQVLLHQQAKFGRNGSDCP DKFCLFQSETKNLLFNDNTECLARLHGKTTYEKYLGPPQYVAGITNLKKCSTSP LEACEFLRK	4
MV-Gt1-F1 primer: 5' ATC GAA GCT TCA TGA GTA ATG TGT GAG CAT TAT GGG ACC ACG 3'	5
Xba-Gt1-R1 primer: 5' CTA GTC TAG ACT CGA GCC ATG GGG CCG GCT AGG GAG CCA TCG CAC AAG AGG AA 3'	6

WHAT IS CLAIMED IS:

1. An improved feed for production animals, comprising
one or more plant-derived feed ingredients, substantially unsupplemented with
5 small-molecule antibiotics, and
as an additive a seed composition containing a flour, extract, or malt obtained
from mature monocot seeds and one or more heterologous seed-produced anti-
microbial proteins in substantially unpurified form.
- 10 2. The feed of claim 1, wherein the one or more seed-produced anti-microbial
protein(s) present in the food are milk proteins selected from the consisting of
lactoferrin, lysozyme, lactoferricin, lactohedrin, kappa-casein, haptocorrin,
lactoperoxidase, alpha-1-antitrypsin, and immunoglobulins.
- 15 3. The feed of claim 2, wherein said the seed-produced proteins are lysozyme
and lactoferrin.
4. The feed of claim 3, wherein lysozyme is present in an amount between
about 0.05 and 0.5 grams protein/kg feed, and lactoferrin, in an amount between 0.2 to
20 2 grams/protein/kg feed.
5. The feed of claim 1, wherein the one or more seed-produced anti-microbial
protein(s) present in the food are acute-phase, non-milk proteins selected from the
consisting of C-reactive protein, serum amyloid A; ferritin, haptoglobin, seromucoids,
25 ceruloplasmin, 15-keto-13,14-dihydro-prostaglandin F2 alpha, fibrinogen, alpha-1-acid
glycoprotein, mannose binding protein, lipopolysaccharide binding protein, alpha-2
macroglobulin and defensins.
6. The feed of claim 1, wherein the one or more seed-produced anti-microbial protein(s)
30 present in the food are antimicrobial peptides selected from the group consisting of cecropin,
magainin, defensins, tachyplesin, parasin I, buforin I, PMAP-23, moronecidin, anoplin, gambicin,
and SAMP-29.
7. The feed of claim 1, wherein the one or more seed-produced anti-microbial
35 protein(s) present in the food are antimicrobial proteins selected from the group

consisting: CAP37, granulysin, secretory leukocyte protease inhibitor, CAP18, ubiquicidin, bovine antimicrobial protein-1, Ace-AMP1, tachyplesin, big defensin, Ac-AMP2, Ah-AMP1, and CAP18.

- 5 8. The feed of claim 1, wherein
- (a) the flour is prepared by milling mature monocot seeds,
- (b) the extract is prepared by suspending milled flour in a buffered aqueous medium; and
- (c) the malt is prepared by (i) steeping barley seeds to a desired water content,
- 10 (ii) germinating the stepped barley, (iii) drying the germinated seeds, under conditions effective to stop germination, (iv) crushing the dried seeds, (v) optionally, adding crushed seeds from a non-barley monocot plant, (vi) forming a mixture of crushed seeds in water, and (vii) malting the crushed seed mixture until a desired malt is achieved, where at least one of the barley or non-barley monocot seeds is stably transformed to
- 15 produce such anti-microbial protein(s).

 9. The feed of claim 8, wherein step (v) includes adding to the crushed dried barley seeds, mature rice transgenic seeds that produce such anti-microbial protein.

- 20 10. In a method for achieving high growth rates in production animals, by feeding animals a feed supplemented with sub-therapeutic levels of one or more small-molecule antibiotics, an improvement comprising
- replacing the small-molecule antibiotic(s) in the feed with a seed composition containing a flour, extract, or malt obtained from mature monocot seeds and one or
- 25 more seed-produced heterologous anti-microbial proteins in substantially unpurified form.

11. The improved method of claim 10, wherein the one or more seed-produced protein(s) present in the feed additive are milk protein(s) selected from the consisting of
- 30 lactoferrin, lysozyme, lactoferricin, lactohedrin, kappa-casein, haptocorrin, lactoperoxidase, alpha-1-antitrypsin, and immunoglobulins.

 12. The improved method of claim 11, wherein the one or more seed-produced proteins are lysozyme and lactoferrin, wherein lysozyme where lysozyme is present in

an amount between about 0.05 and 0.5 grams protein/kg feed, and lactoferrin, in an amount between 0.2 to 2 grams/protein/kg feed.

13. The improved method of claim 10, wherein the one or more seed-produced
5 anti-microbial protein(s) present in the food are acute-phase, non-milk proteins selected from the consisting of C-reactive protein, serum amyloid A; ferritin, haptoglobin, seromucoids, ceruloplasmin, 15-keto-13,14-dihydro-prostaglandin F2 alpha, fibrinogen, alpha-1-acid glycoprotein, mannose binding protein, lipopolysaccharide binding protein, alpha-2 macroglobulin and defensins.

14. The improved method of claim 10, wherein the one or more seed-produced anti-
microbial protein(s) present in the food are antimicrobial peptides selected from the group consisting of cecropin, magainin, defensins, tachyplesin, parasin I, buforin I, PMAP-23, moronecidin, anoplin, gambicin, and SAMP-29.

15. The improved method of claim 10, wherein the one or more seed-produced
anti-microbial protein(s) present in the food are antimicrobial proteins selected from the group consisting: CAP37, granulysin, secretory leukocyte protease inhibitor, CAP18, ubiquicidin, bovine antimicrobial protein-1, Ace-AMP1, tachyplesin, big defensin, Ac-
20 AMP2, Ah-AMP1, and CAP18.

16. A method of producing a feed for production animals, comprising
(a) obtaining a monocot plant that has been stably transformed with a first
chimeric gene having (i) a transcriptional regulatory region from a monocot gene having
25 a seed maturation-specific promoter, (ii) operably linked to said transcriptional regulatory region, a leader DNA sequence encoding a monocot seed-specific transit sequence capable of targeting a linked polypeptide to an endosperm-cell organelle, and (iii) a protein-coding sequence encoding a heterologous anti-microbial protein,
(b) cultivating the transformed plant under seed-maturation conditions,
30 (c) harvesting mature seeds from the cultivated plant,
(d) extracting from the harvested seeds, a flour, extract, or malt composition containing the human milk protein in substantially unpurified form, and
(e) adding the composition to an animal feed that is substantially free of small-
molecule antibiotics.

17. The method of claim 16, wherein the seed-produced protein(s) present in the feed additive are milk protein(s) selected from the group consisting of human or animal lactoferrin, lysozyme, lactoferricin, lactohedrin, kappa-casein, haptocorrin, lactoperoxidase, alpha-1-antitrypsin, and immunoglobulins.

5

18. The method of claim 17, wherein said the seed-produced proteins present in the feed additive are lysozyme and lactoferrin, where lysozyme is present in an amount between about 0.05 and 0.5 grams protein/kg feed, and lactoferrin, in an amount between 0.2 to 2 grams/protein/kg feed.

10

19. The method of claim 17, wherein the protein coding sequence is selected from the group of codon-optimized sequences identified by SEQ ID NOS: 1, 3, 7, and 10-15.

15

20. The method of claim 16, wherein the one or more seed-produced anti-microbial protein(s) present in the food are acute-phase, non-milk proteins selected from the consisting of C-reactive protein, serum amyloid A; ferritin, haptoglobin, seromucoids, ceruloplasmin, 15-keto-13,14-dihydro-prostaglandin F2 alpha, fibrinogen, alpha-1-acid glycoprotein, mannose binding protein, lipopolysaccharide binding protein, alpha-2 macroglobulin and defensins.

20

21. The method of claim 20, wherein the protein coding sequence is selected from the group identified by SEQ ID NOS: 36 and 46-56.

25

22. The method of claim 16, wherein the one or more seed-produced anti-microbial protein(s) present in the food are antimicrobial peptides selected from the group consisting of cecropin, magainin, defensins, tachyplesin, parasin I, buforin I, PMAP-23, moronecidin, anoplin, gambicin, and SAMP-29.

30

23. The method of claim 22, wherein the protein coding sequence is selected from the group identified by SEQ ID NOS: 34-68, 40-41, and 43.

24. The method of claim 16, wherein the one or more seed-produced anti-microbial protein(s) present in the food are antimicrobial proteins selected from the group consisting of: CAP37, granulysin, secretory leukocyte protease inhibitor, CAP18,

35

ubiquicidin, bovine antimicrobial protein-1, Ace-AMP1, tachyplesin, big defensin, Ac-AMP2, Ah-AMP1, and CAP18.

25. The method of claim 24, wherein the protein coding sequence is selected
5 from the group identified by SEQ ID NOS: 37, 45, and 57-59.

26. The method of claim 16, wherein the anti-microbial protein(s) constitute at least 0.25 weight percent of the total protein in the harvested mature seeds.

10 27. The method of claim 16, wherein
(a) the flour is prepared by milling mature monocot seeds,
(b) the extract is prepared by suspending milled flour in a buffered aqueous medium; and
(c) the malt is prepared by (i) steeping barley seeds to a desired water content,
15 (ii) germinating the stepped barley, (iii) drying the germinated seeds, under conditions effective to stop germination, (iv) crushing the dried seeds, and (v) after mixing the crushed seeds with water, malting the crushed seed mixture until a desired malt is achieved.

20 28. The method of claim 27, wherein the malt is further prepared by adding to the crushed dried seeds, mature non-barley transgenic monocot seeds that produce a heterologous, anti-microbial protein.

25 29. The method of claim 16, wherein the monocot plant obtained is further transformed with a second chimeric gene having (i) a transcriptional regulatory region from a monocot gene having a seed maturation-specific promoter, (ii) operably linked to said transcriptional regulatory region, a transit DNA sequence encoding a monocot seed-specific transit sequence capable of targeting a linked polypeptide to an endosperm-cell organelle, and (iii) a protein-coding sequence encoding a second
30 heterologous, anti-microbial protein.

30. The method of claim 16, wherein the transcriptional regulatory region in the chimeric gene is from the promoter of a gene selected from the group of rice glutelins, rice globulins, oryzins, and prolamines, barley hordeins, wheat gliadins and glutenins,

maize zeins and glutelins, oat glutelins, and sorghum kafirins, millet pennisetins, and rye secalins genes.

5 31. The method of claim 30, wherein the leader sequence in the chimeric gene is from the gene selected from the group of rice glutelins, rice globulins oryzins, and prolamines, barley hordeins, wheat gliadins and glutenins, maize zeins and glutelins, oat glutelins, and sorghum kafirins, millet pennisetins, and rye secalins genes.

10 32. The method of claim 31, wherein the transcriptional regulatory region in the chimeric gene is a rice glutelin Gt1 promoter, and the leader DNA sequence is a rice glutelin Gt1 signal sequence capable of targeting a linked polypeptide to a protein storage body.

15 33. The method of claim 32, wherein glutelin Gt1 promoter and glutelin Gt1 signal sequence are included within the sequence identified by SEQ ID NO:15.


20 34. The method of claim 31, wherein the transcriptional regulatory region in the chimeric gene is a rice globulin Glb promoter, and the leader DNA sequence is a rice glutelin Gt1 signal sequence capable of targeting a linked polypeptide to a protein storage body.

35. The method of claim 34, wherein the globulin Glb promoter and glutelin Gt1 signal sequence are included within the sequence identified by SEQ ID NO:16.


25 36. The method of claim 16, wherein the transformed monocot plant further comprises a nucleic acid that encodes at least one transcription factor selected from the group consisting of Reb, O2 and PBF, and an active fragment thereof.

30 37. The method of claim 36, wherein the transcription factor is O2 and/or PBF.



1/45

**Fig. 1**


2/45

**Fig. 2**



3/45

**Fig. 3**


4/45

**Fig. 4**

5/45

Fig. 5A**Fig. 5B****Fig. 5C****Fig. 5D****Fig. 5E**


6/45



Transgenic Lines

Fig. 6

7/45

**Fig. 7**

8/45






Fig. 8


9/45

**Fig. 9**

10/45

**Fig. 10**


11/45

**Fig. 11**

12/45

Fig. 12A

Fig. 12B



pmol/million of cells




Fig. 12C

%




Fig. 12D





Fig. 13

14/45

**Fig. 14**

15/45




Fig. 15

16/45

Fig. 16A







Fig. 16B



17/45

**Fig. 17A**

18/45

**Fig. 17B**

19/45






Fig. 17C


20/45

**Fig. 18A**

21/45

**Fig. 18B**

22/45

**Fig. 19**

23/45

Sequence Range Egfactor: 4 to 165

		10		20		30		40		50								
Egfactor	AAC	TCC	GAC	TCG	GAG	TGC	CCC	CTC	TCC	CAC	GAC	GGT	TAC	TGC	CTC	CAC	GAC	GGG
	N	S	D	S	E	C	P	L	S	H	D	G	Y	C	L	H	D	G>

		3360		3370		3380		3390		3400								
Native Gene	AAT	GGT	GAC	TCT	GAA	TGT	CCC	CTG	TCC	CAC	GAT	GGG	TAC	TGC	CTC	CAT	GAT	GGT
Egfactor	AAC	TCC	GAC	TCG	GAG	TGC	CCC	CTC	TCC	CAC	GAC	GGT	TAC	TGC	CTC	CAC	GAC	GGG

	60		70		80		90		100		110							
Egfactor	GTC	TGC	ATG	TAC	ATC	GAG	GCC	CTC	GAC	AAG	TAC	GCC	TGC	AAC	TGC	GTC	GTG	GGC
	V	C	M	Y	I	E	A	L	D	K	Y	A	C	N	C	V	V	G>

		3410		3420		3430		3440		3450								
Native Gene	GTG	TGC	ATG	TAT	ATT	GAA	GCA	TTG	GAC	AAG	TAT	GCA	TGC	AAC	TGT	GTT	GTT	GGC
Egfactor	GTC	TGC	ATG	TAC	ATC	GAG	GCC	CTC	GAC	AAG	TAC	GCC	TGC	AAC	TGC	GTC	GTG	GGC

		120		130		140		150		160								
Egfactor	TAC	ATC	GGC	GAG	CGG	TGC	CAG	TAC	CGC	GAC	CTC	AAG	TGG	TGG	GAG	CTG	CGC	TGA
	Y	I	G	E	R	C	Q	Y	R	D	L	K	W	W	E	L	R	*>


		3460		3470		3480		3490		3500							
Native Gene	TAC	ATC	GGG	GAG	CGA	TGT	CAG	TAC	CGA	GAC	CTG	AAG	TGG	TGG	GAA	CTG	CGC
Egfactor	TAC	ATC	GGC	GAG	CGG	TGC	CAG	TAC	CGC	GAC	CTC	AAG	TGG	TGG	GAG	CTG	CGC

Epidermal Growth Factor


Number of codons in mature peptide: 53
 Number of codons changed: 27 (51%)
 Number of nucleotides changed: 30 (19%)

Fig. 20

24/45

**Fig. 21**

25/45

**Fig. 22**

26/45




Fig. 23

27/45

Sequence Range: 4 to 216

Insgfact	GGC CCA GAG ACC CTG TGC GGT CCG GAG CTG CTC GAC GCC CTC CAG TTC GTC TGC GGG GAC CCG GGC TTC TAC TTC
	G P E T L C G A E L V D A L Q P V C G D R G F Y F>
native gene	CGA CCG GAG ACG CTC TGC GGG GCT GAG CTG GTG GAT GCT CTT CAG TTC GTG TGT CGA GAC AGG GGC TTT TAT TTC>
Insgfact	GGC CCA GAG ACC CTG TGC GGT CCG GAG CTG CTC GAC GCC CTC CAG TTC GTC TGC GGG GAC CCG GGC TTC TAC TTC
Insgfact	AAC AAG CCA ACG GGC TAC GGG TCC TCC TCG CGC CGC GCC CCC CAG ACC GGC ATC GTG GAC GAG TGC TGC TTC CCC
	N K P T G Y G S S S R R A P Q T G I V D E C C F R>
native gene	AAC AAG CCC ACA GGG TAT GGC TCC AGC AGT CCG AGG CCG CCT CAG ACA GGC ATC GTG GAT GAG TGC TGC TTC CCG>
Insgfact	AAC AAG CCA ACG GGC TAC GGC TCC TCC TCG CGC CGC GCC CCC CAG ACC GGC ATC GTG GAC GAG TGC TGC TTC CCG
Insgfact	TCC TGC GAC CTC CCG CCG CTC GAG ATG TAC TGC GCC CCA CTC AAG CCC GGC AAG AGC GCC TGA
	S C D L R R L E M Y C A P L K P A K S A *>
native gene	AGC TCT GAT CTA ACG AGG CTG GAG ATG TAT TGC GCA CCC CTC AAG CCT GCC AAG TCA GCT>
Insgfact	TCC TGC GAC CTC CCG CCG CTC GAG ATG TAC TGC GCC CCA CTC AAG CCC GCC AAG AGC GCC

Fig. 24

28/45





Fig. 25

29/45

**Fig. 26**

30/45






Fig. 27


31/45

**Fig. 28**


32/45

**Fig. 29**


33/45

**Fig. 30**


34/45

**Fig. 31**


35/45

**Fig. 32**


36/45

**Fig. 33**


37/45

**Fig. 34**


38/45

**Fig. 35**


39/45

**Fig. 36A**


40/45

**Fig. 36B**



41/45

**Fig. 37**


42/45

**Fig. 38**


43/45

**Fig. 39A****Fig. 39B**

44/45

**Fig. 40**

45/45

**Fig. 41**

SEQUENCE LISTING

<110> Ventria Bioscience

<120> Feed Additive Compositions and Methods

<130> 50665-8021.W000

<140> Not Yet Assigned

<141> Filed Herewith

<150> US 60/269,188

<151> 2001-02-14

<150> US 09/847,232

<151> 2001-05-02

<160> 60

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 393

<212> DNA

<213> Homo sapiens

<400> 1

aaagtcttcg	agcgggtgcga	gctggccccgc	acgctcaagc	ggctcggcat	ggacggctac	60
cggggcatca	gcctcgccaa	ctggatgtgc	ctcgccaagt	gggagtcggg	ctacaacacc	120
cgcgcaacca	actacaacgc	cggcgaccgc	tccaccgact	acggcatctt	ccagatcaac	180
tcccgtact	ggtgcaacga	cggcaagacg	cccggggccg	tcaacgcctg	ccacctctcc	240
tgctcgcccc	tgctgcaaga	caacatcgcc	gacgcgctcg	cgtgcgcgaa	gcgcgtcgtc	300
cgcgacccgc	agggcatccg	ggcctgggtg	gcctggcgca	accgctgcca	gaaccgggac	360
gtgcgccagt	acgtccaggg	ctgcggcgtc	tga			393

<210> 2

<211> 130

<212> PRT

<213> Homo sapiens

<400> 2

Lys	Val	Phe	Glu	Arg	Cys	Glu	Leu	Ala	Arg	Thr	Leu	Lys	Arg	Leu	Gly
1				5					10					15	
Met	Asp	Gly	Tyr	Arg	Gly	Ile	Ser	Leu	Ala	Asn	Trp	Met	Cys	Leu	Ala
			20					25					30		
Lys	Trp	Glu	Ser	Gly	Tyr	Asn	Thr	Arg	Ala	Thr	Asn	Tyr	Asn	Ala	Gly
		35					40					45			
Asp	Arg	Ser	Thr	Asp	Tyr	Gly	Ile	Phe	Gln	Ile	Asn	Ser	Arg	Tyr	Trp
	50				55				60						
Cys	Asn	Asp	Gly	Lys	Thr	Pro	Gly	Ala	Val	Asn	Ala	Cys	His	Leu	Ser
65				70					75					80	
Cys	Ser	Ala	Leu	Leu	Gln	Asp	Asn	Ile	Ala	Asp	Ala	Val	Ala	Cys	Ala
			85				90							95	
Lys	Arg	Val	Val	Arg	Asp	Pro	Gln	Gly	Ile	Arg	Ala	Trp	Val	Ala	Trp
		100					105						110		
Arg	Asn	Arg	Cys	Gln	Asn	Arg	Asp	Val	Arg	Gln	Tyr	Val	Gln	Gly	Cys
	115					120						125			
Gly	Val														
	130														

<210> 3

<211> 2079

<212> DNA

<213> Homo sapiens

<400> 3

```

gggcggcggtt ggcgctcggt gcagtgggtgc gccgtgtccc agcccgagggc gaccaagtgc      60
ttccagtggc agcgcaacat gcggaagggtg cgcggcccgc cggtcagctg catcaagcgg      120
gactccccc tccaatgcat ccaggccatc gcggagaacc gcgccgacgc ggtcaccctg      180
gacggcggtt tcatctacga ggcgggggtc gcccgtaca agtccgccc ggtggcggtg      240
gaggtgtacg gcaccgagcg ccagccgcgc acgcactact acgcggtggc cgtcgtcaag      300
aaggcggggt ccttcagct caacgagctg cagggcctga agtcgtgcca cacgggcctc      360
cggcgacggc cgggctggaa cgtgcccac ggcacctgc gccccttctt gaactggacc      420
ggcccgccgg agccgatcga ggccgcccgt gcccgttctt tcagcgctc ctgcgtcccc      480
ggcgccgaca agggccagtt cccgaacctc tgccggctct gcgccgggac gggcgagaac      540
aagtgcgcct tctcctcgca ggagccgtac ttctcctact cgggcgcggt caagtgcctc      600
cgcgacgggg cggcgacgtt ggcgttcac cgcgagtcca ccgtgttcga ggacctctcc      660
gacgaggcgg agcgggacga gtacgagctg ctgtgcccc acaacacccg caagccggtg      720
gacaagttca aggactgcca cctggcgcg gtgcccctgc acgcggtcgt cgcgcgagc      780
gtcaacggca aggaggacgc gatctggaac ctctccgcc agggccagga gaagtccggc      840
aaggacaagt cccccaagt ccagctcttc gggagcccca gcggccagaa ggacctctc      900
ttcaaggact ccgcatcggt cttctccgc gtccccccgc gcatcgactc cggcctgtac      960
ctcggctccg ggtacttcac cgcgatccag aacctccgga agagcgagga ggaggtggcg      1020
gcgcggcggt cccgcgtcgt gtggtgcgc gtgggcgagc aggagctgcg gaagtgaac      1080
cagtggagcg gctgagcga ggggtcgggt acctgctcgt ccgccagcac caccgaggac      1140
tgcatcgcg tgcctctcaa gggggaggcc gacgcgatga gcctcgacgg ggggtacgtc      1200
tacaccgccc gcaagtgcgg cctgggtccc gtcttgccgg agaactaaa gtcgcagcag      1260
tccagcgacc ccgacccgaa ctgctgggac cgcgccgtcg agggctacct cgcctgggc      1320
gtcgtgcgcc ggtccgacac ctccctgac tggaacagcg tcaagggcaa gaagagctgc      1380
cacaccgccc tggaccgcac cgcgggtgg aacatccgca tgggcctcct cttcaaccag      1440
accggtcctt gcaagttcga cgagtacttc tccagtcct cgcgcccggt ctcggacccc      1500
cgctccaacc tgtgcgcctt ctgcatcggt gacgagcagg gcgagaacaa gtgcgtgccc      1560
aacagcaacg agcgggtacta cggctacac ggggccttcc gctgcctggc ggagaacgcc      1620
ggggacgtcg cgttcgtgaa ggacgtgacc gtgctgcaaa acacgagcgg gaacaacaac      1680
gaggcgtggg cgaaggacct caagctcgcc gaattcgccc tgcgtgtgct cgacggcaag      1740
cgcaagcccc tcaccgaggc gcggtcctgc cactggcga tggcccccaa ccagccgctc      1800
gtctcccgca tggacaaggt cgagcgctc aagcaggtgc tctgcaaca ccagggcaag      1860
ttcggccgga acggcagcga ctgcccggac aagttctgcc tgttccagtc ggagaccaag      1920
aacctcctct tcaacgacaa caccgagtgc ctggcgcgcc tccacggcaa gaccacctac      1980
gagaagtacc tcggcccgca gtacgtcgcc ggcacacca acctcaagaa gtgctccacc      2040
tccccctcc tggaggcggt cgagttcctc cgcaagtga      2079

```

<210> 4

<211> 690

<212> PRT

<213> Homo sapiens

<400> 4

```

Gly Arg Arg Arg Arg Ser Val Gln Trp Cys Ala Val Ser Gln Pro Glu
 1          5          10          15
Ala Thr Lys Cys Phe Gln Trp Gln Arg Asn Met Arg Lys Val Arg Gly
 20          25          30
Pro Pro Val Ser Cys Ile Lys Arg Asp Ser Pro Ile Gln Cys Ile Gln
 35          40          45
Ala Ile Ala Glu Asn Arg Ala Asp Ala Val Thr Leu Asp Gly Gly Phe
 50          55          60
Ile Tyr Glu Ala Gly Leu Ala Pro Tyr Lys Leu Arg Pro Val Ala Ala
 65          70          75          80
Glu Val Tyr Gly Thr Glu Arg Gln Pro Arg Thr His Tyr Tyr Ala Val
 85          90          95
Ala Val Val Lys Lys Gly Gly Ser Phe Gln Leu Asn Glu Leu Gln Gly
 100         105         110
Leu Lys Ser Cys His Thr Gly Leu Arg Arg Thr Ala Gly Trp Asn Val
 115         120         125
Pro Ile Gly Thr Leu Arg Pro Phe Leu Asn Trp Thr Gly Pro Pro Glu
 130         135         140
Pro Ile Glu Ala Ala Val Ala Arg Phe Phe Ser Ala Ser Cys Val Pro
 145         150         155         160

```

Gly	Ala	Asp	Lys	Gly	Gln	Phe	Pro	Asn	Leu	Cys	Arg	Leu	Cys	Ala	Gly
				165					170					175	
Thr	Gly	Glu	Asn	Lys	Cys	Ala	Phe	Ser	Ser	Gln	Glu	Pro	Tyr	Phe	Ser
			180					185					190		
Tyr	Ser	Gly	Ala	Phe	Lys	Cys	Leu	Arg	Asp	Gly	Ala	Gly	Asp	Val	Ala
		195					200					205			
Phe	Ile	Arg	Glu	Ser	Thr	Val	Phe	Glu	Asp	Leu	Ser	Asp	Glu	Ala	Glu
	210				215						220				
Arg	Asp	Glu	Tyr	Glu	Leu	Cys	Pro	Asp	Asn	Thr	Arg	Lys	Pro	Val	
	225				230				235					240	
Asp	Lys	Phe	Lys	Asp	Cys	His	Leu	Ala	Arg	Val	Pro	Ser	His	Ala	Val
			245						250					255	
Val	Ala	Arg	Ser	Val	Asn	Gly	Lys	Glu	Asp	Ala	Ile	Trp	Asn	Leu	Leu
			260					265					270		
Arg	Gln	Ala	Gln	Glu	Lys	Phe	Gly	Lys	Asp	Lys	Ser	Pro	Lys	Phe	Gln
	275						280						285		
Leu	Phe	Gly	Ser	Pro	Ser	Gly	Gln	Lys	Asp	Leu	Leu	Phe	Lys	Asp	Ser
	290					295					300				
Ala	Ile	Gly	Phe	Ser	Arg	Val	Pro	Pro	Arg	Ile	Asp	Ser	Gly	Leu	Gly
	305				310					315				320	
Ser	Gly	Tyr	Phe	Thr	Ala	Ile	Gln	Asn	Leu	Arg	Lys	Ser	Glu	Glu	Glu
				325					330					335	
Val	Ala	Ala	Arg	Arg	Ala	Arg	Val	Val	Trp	Cys	Ala	Val	Gly	Glu	Gln
			340					345					350		
Glu	Leu	Arg	Lys	Cys	Asn	Gln	Trp	Ser	Gly	Leu	Ser	Glu	Gly	Ser	Val
	355						360					365			
Thr	Cys	Ser	Ser	Ala	Ser	Thr	Thr	Glu	Asp	Cys	Ile	Ala	Leu	Val	Leu
	370					375					380				
Lys	Gly	Glu	Ala	Asp	Ala	Met	Ser	Leu	Asp	Gly	Gly	Tyr	Val	Tyr	Thr
	385				390					395					400
Ala	Gly	Lys	Cys	Gly	Leu	Val	Pro	Val	Leu	Ala	Glu	Asn	Tyr	Lys	Ser
				405					410					415	
Gln	Gln	Ser	Ser	Asp	Pro	Asp	Pro	Asn	Cys	Val	Asp	Arg	Pro	Val	Glu
			420					425					430		
Gly	Tyr	Leu	Ala	Val	Ala	Val	Val	Arg	Arg	Ser	Asp	Thr	Ser	Leu	Thr
		435					440					445			
Trp	Asn	Ser	Val	Lys	Gly	Lys	Lys	Ser	Cys	His	Thr	Ala	Val	Asp	Arg
	450					455					460				
Thr	Ala	Gly	Trp	Asn	Ile	Pro	Met	Gly	Leu	Leu	Phe	Asn	Gln	Thr	Gly
	465				470					475					480
Ser	Cys	Lys	Phe	Asp	Glu	Tyr	Phe	Ser	Gln	Ser	Cys	Ala	Pro	Gly	Ser
				485					490					495	
Asp	Pro	Arg	Ser	Asn	Leu	Cys	Ala	Leu	Cys	Ile	Gly	Asp	Glu	Gln	Gly
			500					505					510		
Glu	Asn	Lys	Cys	Val	Pro	Asn	Ser	Asn	Glu	Arg	Tyr	Tyr	Gly	Tyr	Thr
		515					520					525			
Gly	Ala	Phe	Arg	Cys	Leu	Ala	Glu	Asn	Ala	Gly	Asp	Val	Ala	Phe	Val
	530					535					540				
Lys	Asp	Val	Thr	Val	Leu	Gln	Asn	Thr	Asp	Gly	Asn	Asn	Asn	Glu	Ala
	545				550					555					560
Trp	Ala	Lys	Asp	Leu	Lys	Leu	Ala	Asp	Phe	Ala	Leu	Leu	Cys	Leu	Asp
				565					570					575	
Gly	Lys	Arg	Lys	Pro	Val	Thr	Glu	Ala	Arg	Ser	Cys	His	Leu	Ala	Met
			580					585					590		
Ala	Pro	Asn	His	Ala	Val	Val	Ser	Arg	Met	Asp	Lys	Val	Glu	Arg	Leu
		595					600					605			
Lys	Gln	Val	Leu	Leu	His	Gln	Gln	Ala	Lys	Phe	Gly	Arg	Asn	Gly	Ser
	610					615					620				
Asp	Cys	Pro	Asp	Lys	Phe	Cys	Leu	Phe	Gln	Ser	Glu	Thr	Lys	Asn	Leu
	625				630					635					640
Leu	Phe	Asn	Asp	Asn	Thr	Glu	Cys	Leu	Ala	Arg	Leu	His	Gly	Lys	Thr
				645					650					655	
Thr	Tyr	Glu	Lys	Tyr	Leu	Gly	Pro	Gln	Tyr	Val	Ala	Gly	Ile	Thr	Asn
			660					665					670		
Leu	Lys	Lys	Cys	Ser	Thr	Ser	Pro	Leu	Leu	Glu	Ala	Cys	Glu	Phe	Leu

675 680 685

Arg Lys
690

<210> 5
<211> 42
<212> DNA
<213> Artificial Sequence

<220>
<223> primer

<400> 5
atcgaagctt catgagtaat gtgtgagcat tatgggacca cg 42

<210> 6
<211> 53
<212> DNA
<213> Artificial Sequence

<220>
<223> primer

<400> 6
ctagtctaga ctcgagccat ggggccggct agggagccat cgcacaagag gaa 53

<210> 7
<211> 72
<212> DNA
<213> Artificial Sequence

<220>
<223> codon optimized lactoferricin coding sequence
based on Homo sapiens sequence

<400> 7
accaagtgt tccagtggca gcgcaacatg cggaaggtgc gcggcccgcc ggtcagctgc 60
atcaagcggg ac 72

<210> 8
<211> 162
<212> DNA
<213> Artificial Sequence

<220>
<223> codon optimized EGF coding sequence based on Homo
sapiens sequence

<400> 8
aactccgact cggagtgtccc cctctccac gacgggttact gcctccacga cgggggtctgc 60
atgtacatcg aggcctcga caagtacgcc tgcaactgcg tcgtgggcta catcggcgag 120
cggtgccagt accgcgacct caagtgggtg gagctgcgct ga 162

<210> 9
<211> 213
<212> DNA
<213> Artificial Sequence

<220>
<223> codon optimized IGF-1 coding sequence based on
Homo sapiens sequence

<400> 9
ggcccgagaga ccctctgcgg cgccgagctc gtggacgccc tccagttcgt gtgcggcgac 60
cgcggttct acttcaacaa gccgaccggc tacggcagca gcagccgccc cgccccgcag 120

accggcatcg tggacgagtg ctgcttccgc agctgcgacc tccgccgcct ggagatgtac 180
tgcgccccgc tcaagccccg caagagcgcc tga 213

<210> 10

<211> 1095

<212> DNA

<213> Artificial Sequence

<220>

<223> codon optimized lactohedrin coding sequence based
on Homo sapiens sequence

<400> 10

ctggacatct	gctcgaagaa	cccgtgccac	aacggcgggc	tctgcgagga	gatcagccag	60
gaggtgcg	gagacgtgtt	cccctcgta	acctgcacct	gcctgaaggg	ctacgccggg	120
aacctg	agacgaagt	cgtggagccc	ctggggatgg	agaacggcaa	catcgccaac	180
tccagatcg	ccgcctcctc	cgtgcggtg	accttcctcg	gcctccagca	ctgggtcccg	240
gagctggccc	ggctcaaccg	ggcgggcatg	gtgaacgcgt	ggacccccctc	gtccaacgac	300
gacaaccgt	ggatccaagt	gaacctgctc	cgccgcatgt	gggtcaccgg	cgtggtcacc	360
caaggcgcca	gccgcctggc	cagccacgag	tacctcaagg	ccttcaagg	cgcctacagc	420
ctcaaccggc	acgagttcga	cttcattccac	gacgtcaaca	agaagcaca	ggagttcgtg	480
ggcaactgga	acaagaacgc	ggtccacgtg	aacctcttcg	agacccccgt	cgaggcccag	540
tacgtccgcc	tctacccccac	gagctgccac	accgcctgca	cgctccgctt	cgagctgctg	600
gggtgcgagc	tgaacgggtg	cgcgaaccgc	ctggggctca	agaacaacag	catccccgac	660
aagcagatca	cggcctcgct	gtcgtacaag	acctggggcc	tgcacctctt	ctcgtggaac	720
ccgagctacg	cccggctgga	caagcagggc	aacttcaacg	cctgggtcgc	cgggagctac	780
gggaacgacc	agtggctcca	ggtggacctc	ggcagctcca	aggaggtcac	cggcatcatc	840
acgcaggggg	cccgcaactt	cggctccgtg	cagttcgtgg	cctcctacaa	ggtggcctac	900
tcgaacgaca	gcgccaactg	gaccgagtac	caggaccgcg	gcaccgggtc	cagcaagatc	960
ttccccggca	actgggacaa	ccacagccac	aagaagaacc	tgttcgagac	ccccatcctc	1020
gcccggtagc	tccgcatcct	ccccgtcgct	tggcacaacc	ggatcgcgct	cgggctggag	1080
ctcctcggtc	gctga					1095

<210> 11

<211> 489

<212> DNA

<213> Artificial Sequence

<220>

<223> codon optimized kappa-casein coding sequence based
on Homo sapiens sequence

<400> 11

gaggtccaaa	accagaagca	gcccgcctgc	cacgagaacg	acgagcgccc	cttctaccag	60
aagaccgcac	cctacgtccc	gatgtactac	gtcccgaaca	gctaccccta	ctacggtacg	120
aacctgtacc	agcgccgccc	ggccatcgct	atcaacaacc	cctacgtccc	ccggacctac	180
tacgcgaacc	cggccgtggt	gcggccccac	gcgcagatcc	cgcagcggca	gtacctgcca	240
aacagccacc	cccccaaccg	ggtgcggcgg	cccaacctcc	acccgagctt	catcgctatc	300
ccccccaaga	agatccagga	caagatcatc	atcccgaacca	tcaacaccat	cgccaccgtg	360
gagccgacgc	cagcccccg	gaccgagccc	acggtggaca	gcgtcgtgac	cccagaggcg	420
ttctccgaat	cgatcatcac	ctccaccccc	gagaccacca	cgggtggcgt	cacgccgcgc	480
acggcatga						489

<210> 12

<211> 1233

<212> DNA

<213> Artificial Sequence

<220>

<223> codon optimized haptocorrin coding sequence based
on Homo sapiens sequence

<400> 12

gagatctg	aggtctccga	ggagaactac	atccgcctca	agccccctct	gaacaccatg	60
atccagagca	actacaaccg	gggcacgtcg	gccgtgaacg	tcgtgctctc	cctgaagctc	120

gtgggcatcc	agatccagac	cctcatgcag	aagatgatcc	agcagatcaa	gtacaacgtg	180
aagagccgcc	tctcggacgt	gtccagcggc	gagctggcgc	tcacatcctt	cgcgctcggc	240
gtgtgccgga	acgcggagga	gaacctcatc	tacgactacc	acctcacgga	caagctggag	300
aacaagttcc	aggccgagat	cgagaacatg	gaggcccaca	acggcacccc	gctgaccaac	360
tactaccagc	tcagcctgga	cgctcctcgc	ctctgcctgt	tcaacgggaa	ctactccacc	420
gccgaggtgg	tcaaccactt	cacccccgag	aacaagaact	actacttcgg	ctcgagttc	480
tccgtggaca	ccggggccat	ggcgcctctg	gccctcacct	gcgtgaagaa	gtccctcatc	540
aacggccaga	tcaaggccga	cgagggtctc	ctgaagaaca	tctcgatcta	caccaagagc	600
ctcgtggaga	agatcctcag	cgagaagaag	gagaacgggc	tgatcggcaa	caccttctcg	660
accggcgagg	cgatgcaggc	cctgttcgtg	agcagcgact	actacaacga	gaacgactgg	720
aactgccagc	agacctcaa	cacggtcctg	accgagatca	gccagggcgc	gttcagcaac	780
cccaacgccg	ccgcccaggt	cctgccggcc	ctgatgggca	agaccttcct	cgacatcaac	840
aaggacagct	cctgcgtgtc	cgcgagcggc	aacttcaaca	tctccgccga	cgagccgata	900
acggtgacgc	cgcccagacg	ccagtcgtac	atctccgtga	actacagcgt	gcggatcaac	960
gagacctact	tcacgaacgt	gacggtcctc	aacggctcgg	tcttctctgag	cgtgatggag	1020
aaggcgacga	agatgaacga	cacgatcttc	ggcttcacga	tggaggagcg	cagctggggc	1080
ccctacatca	cctgcatcca	gggcctctgc	ggcaacaaca	acgaccgcac	ctactggggg	1140
ctgctgagcg	gcggcgagcc	gctgagccag	ggggccggca	gctacgtggt	ccgcaacggc	1200
gagaacctgg	aggtccggtg	gagcaagtac	tga			1233

<210> 13

<211> 2061

<212> DNA

<213> Artificial Sequence

<220>

<223> codon optimized lactoperoxidase coding sequence
based on Homo sapiens sequence

<400> 13

caaacgaccc	ggacgtcggc	gatctccgac	acgggtctcgc	aggccaaggt	gcaagtcaac	60
aaggcattcc	tggattcgcg	cacgcggctg	aagaccgcga	tgtcgtccga	gaccccgacg	120
agccggcagc	tgagcgagta	cctcaagcac	gcgaaggggc	ggacgcgcac	cgccatccgc	180
aatggccaag	tgtgggagga	atccctgaag	cggtgcggc	agaaggcgtc	gctcaccaac	240
gtgaccgacc	cgctccctga	cctgaccagc	ctctccctgg	aggtcggctg	cggcgccccg	300
gcgcccgtcg	tgcgtcgcga	cccctgctcg	ccataccgca	cgatcacggg	cgactgcaac	360
aaccggcgga	agccggcact	gggggctgcg	aaccgcgccc	tcgcgcgctg	gctccccgcc	420
gagtacgagg	acggcctcag	cctccccttc	ggttggaccc	ccggcaagac	gcgcaacggc	480
ttcccgtctc	cgctcgtctg	cgaggtcagc	aacaagatcg	tcggttacct	gaacgaggag	540
ggggtctctg	acaaaaaccg	ctccctcctc	ttcatgcagt	gggggcagat	cgtggaccac	600
gacctggact	tcgccccgga	cacggagctg	ggctccagcg	agtacagcaa	gacctagtcg	660
gacgaatact	gcattccagg	cgacaactgc	ttcccgatca	tgttcccccc	gaacgacccg	720
aaggcgggca	cccagggcaa	gtgcatgccg	ttcttccggg	caggcttcgt	ctgcccgcac	780
cccccgatca	agtcctcgc	gcgcgagcag	atcaacgcgc	tcacgtcctt	cctcgacgcc	840
agcttcgtct	acagcagcga	gccgtccctc	gccagecgcc	tccgcaacct	cagcagcccc	900
ctcggcctca	tggcgggtcaa	ccaggaggtg	tcggaccacg	gcctcccata	cctgcccgtac	960
gacagcaaga	agccgtcccc	ctgcgagttc	atcaacacca	ccgcgcgcgt	ccgtgcttc	1020
ctcgcggcg	attcgcgggc	gagcgagcac	atcctcctcg	ccacgagcca	cacctgttct	1080
ctccgcgagc	acaaccgcct	cgcccgggag	ctgaagcgcc	tcaaccgcga	gtgggacggc	1140
gagaagctct	accaggaggc	ccggaagatc	ctcggcgctt	tcgtccagat	catcaccttc	1200
cgggactacc	tccccatcct	gctcggtgac	cacatgcaga	agtggatccc	cccctaccaa	1260
ggctaactcg	agagcgtgga	cccgcgcctc	tccaacgtct	tcacgttcgc	gttccgcttc	1320
gggcacctgg	aggtgcgctc	gtcgtatgtc	cgcctcgacg	agaactacca	gccctggggc	1380
ccagagcccg	agctgcgcgt	ccacaccctg	ttcttcaaca	cctggcggat	ggtcaaggac	1440
ggcggcatcg	acccgctcgt	gcgcgggctc	ctggctaaga	agtcgaagct	catgaagcag	1500
aacaagatga	tgaccggcga	gctgcgcaac	aagctgttcc	agcccaccca	ccgcatccac	1560
gggttcgacc	tggctgcaat	caacaccacg	cggtgccgcg	accacggcca	gcccggctac	1620
aactcgtggc	gcgcgttctg	cgacctctcc	cagccacaga	cgctggagga	gctcaacacc	1680
gtgctcaaga	gcaagatgtc	cgccaagaag	ctgctcgggc	tctacggcac	gcccgcacac	1740
atcgacatct	ggatcggggc	catcgcggag	ccgctcgtgg	agcgcggggc	cgcggcccg	1800
ctgctcgcgt	gcctcctggg	caagcaattc	caacagatcc	gcgacgggga	ccggttctgg	1860
tgggagaacc	ccggcggtgt	caccaacgag	cagaaggatt	cgctccaaaa	gatgagcttc	1920
tcccgcctgg	tgtgcgacaa	caccgcgctc	accaaggtcc	cgcgcgaccc	attctggggc	1980
aactcctacc	cgtacgactt	cgtggactgc	tccgccatcg	acaagctcga	cctgtccccc	2040
tgggcatcgg	tgaagaactg	a				2061

<210> 14
 <211> 1185
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> codon optimized alpha-1-antitrypsin coding
 sequence based on Homo sapiens sequence

<400> 14
 gaggacccgc agggcgacgc cgcccagaag accgacacca gccaccacga ccaggaccac 60
 ccgacgttca acaagatcac cccgaatttg gccgaattcg ccttcagcct gtaccgccag 120
 ctgcgcaccc agtccaactc caccaacatc ttcttcagcc cgggtgagcat cgccaccgcc 180
 ttccgcatgc tgtccctggg taccaaggcg gacacccacg acgagatcct cgaagggctg 240
 aacttcaacc tgacggagat cccggaggcg cagatccacg agggcttcca ggagctgctc 300
 aggacgtca accagccgga ctcccagctc cagctcacca ccggcaacgg gctcttctctg 360
 tccgagggcc tcaagctcgt cgataagttc ctggaggacg tgaagaagct ctaccactcc 420
 gaggcggttca ccgtcaactt cggggacacc gaggaggcca agaagcagat caacgactac 480
 gtcgagaagg ggacccaggg caagatcgtg gacctggtca aggaattgga cagggaacac 540
 gtcttcgcgc tcgtcaacta catcttcttc aagggaaggt gggagcgccc gttcgagggtg 600
 aaggacaccg aggaggagga cttccacgtc gaccaggtca ccaccgtcaa ggtcccgatg 660
 atgaagaggc tggcgtggtt caacatccag cactgcaaga agctctccag ctgggtgctc 720
 ctcatgaagt acctggggaa cgccaccgcc atcttcttcc tgccggacga gggcaagctc 780
 cagcacctgg agaacgagct gacgcacgac atcatcacga agttcctgga gaacgaggac 840
 aggcgtctcc ctagcctcca cctcccgaag ctgagcatca ccggcacgta cgacctgaag 900
 agcgtgctgg gccagctggg catcacgaag gtcttcagca acggcgcgga cctctccggc 960
 gtgacggagg agggccccct gaagctctcc aaggccgtgc acaaggcggg gctcacgac 1020
 gacgagaagg ggacgggaagc tgccggggcc atgttctctgg aggccatccc cgtgtccatc 1080
 ccgcccaggg tcaagttcaa caagcccttc gtcttctctga tgatcgagca gaacacgaag 1140
 agccccctct tcatggggaa ggtcgtcaac ccacgcaga agtga 1185

<210> 15
 <211> 786
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Rice Gt1 promoter and Gt1 leader coding sequence

<400> 15
 catgagtaat gtgtgagcat tatgggacca cgaaataaaa agaacatttt gatgagtcgt 60
 gtatcctcga tgagcctcaa aagttctctc accccggata agaaaccctt aagcaatgtg 120
 caaagtttgc attctccact gacataatgc aaaataagat atcatcgatg acatagcaac 180
 tcatgcatca tatcatgcct ctctcaacct attcatctct actcatctac ataagtatct 240
 tcagctaaat gttagaacat aaaccataaa gtcacgtttg atgagtatta ggcgtgacac 300
 atgacaaatc acagactcaa gcaagataaa gcaaaatgat gtgtacataa aactccagag 360
 ctatatgtca tattgcaaaa agaggagagc ttataagaca aggcattgact caaaaaatt 420
 cacttgccct tcgtgtcaaa aagaggaggg ctttacatta tccatgtcat attgcaaaag 480
 aaagagagaa agaacaacac aatgctgctg caattataca tatctgtatg tccatcatta 540
 ttcattccacc tttcgtgtac cacacttcat atatcataag agtcacttca cgtctggaca 600
 ttaacaaact ctatcttaac atttagatgc aagagccttt atctcactat aaatgcacga 660
 tgatttctca ttgtttctca caaaaagcgg ccgcttcatt agtcctacaa caacatggca 720
 tccataaata gccccatagt tttcttcaca gtttgcttgt tctcttctgt cgatggctcc 780
 ctagcc 786

<210> 16
 <211> 1055
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Rice Glb promoter and Gt1 leader coding sequence

<400> 16

ctgcaggggag	gagagggggag	agatggtgag	agaggaggaa	gaagaggagg	ggtgacaatg	60
atatgtgggg	catgtgggca	cccaattttt	taattcattc	ttttgttgaa	actgacatgt	120
gggtcccatg	agattttatta	tttttcggat	cgaatcgcca	cgtaagcgct	acgtcaatgc	180
tacgtcagat	gaagaccgag	tcaaatttagc	cacgtaagcg	ccacgtcagc	caaaaccacc	240
atccaaaccg	ccgaggggacc	tcatctgcac	tggttttgat	agttgaggga	cccgttgat	300
ctggtttttc	gattgaagga	cgaaaatcaa	atgtgttgac	aagttaaggg	accttaaatg	360
aacttattcc	atctcaaaaat	attctgtgag	ccatatatac	cgtgggcttc	caatcctcct	420
caaattaaag	ggccttttta	aaatagataa	ttgccttctt	tcagtcaccc	ataaaagtac	480
aaaactacta	ccaacaagca	acatgcgag	ttacacacat	tttctgcaca	tttcgcccac	540
gtcacaaaga	gctaagagtt	atccctagga	caatctcatt	agtgtagata	catccattaa	600
tcttttatca	gaggcaaacg	taaagccgct	ctttatgaca	aaaatagggtg	acacaaaagt	660
gttatctgcc	acatacataa	cttcagaaat	tacccaacac	caagagaaaa	ataaaaaaaa	720
atctttttgc	aagctccaaa	tcttggaaac	cttttttact	ctttgcagca	ttgtactctt	780
gctctttttc	caaccgatcc	atgtcacccct	caagcttcta	cttgatctac	acgaagctca	840
ccgtgcacac	aaccttgcc	acaaaaaccc	tataaaaccc	catccgatcg	ccatcatctc	900
atcatcagtt	cattaccaac	aaacaaaaga	ggaaaaaaa	catatacact	tctagtatt	960
gtctgattga	tcatcaatct	agaggcgcc	gcattgctag	caaggctcgtc	ttcttcgcgg	1020
cggcgctcat	ggcggccatg	gtggccatct	cgggc			1055

<210> 17

<211> 976

<212> DNA

<213> Artificial Sequence

<220>

<223> Bx7 promoter

<400> 17

ctgcaggcca	gggaaagaca	atggacatgc	aaagaggtag	gggcaggga	gaaacacttg	60
gagatcatag	aagaacataa	gaggttaaac	ataggagggc	ataatggaca	attaaatcta	120
cattaattga	actcatttg	gaagtaaaca	aaatccatat	tctgggtgtaa	atcaaaactat	180
ttgacgcgga	tttactaaga	tcctatgtta	attttagaca	tgactggcca	aaggtttcag	240
ttagttcatt	tgtcacggaa	aggtgttttc	ataagtccaa	aactctacca	acttttttgc	300
acgtcatagc	atagatagat	gttgtgagtc	attggataga	tattgtgagt	cagcatggat	360
ttgtgttgcc	tggaaatcca	actaaatgac	aagcaacaaa	acctgaaatg	ggcttttagga	420
gagatggttt	atcaattttac	atgttccatg	caggctacct	tccactactc	gacatggtta	480
gaagttttga	gtgccgcata	tttgcggaag	caatggcact	actcgacatg	gttagaagtt	540
ttgagtgccg	catattttgcg	gaagcaatgg	ctaacagata	catattctgc	caaaccceaa	600
gaaggataat	cactcctctt	agataaaaaag	aacagacca	tgtacaaaca	tccacacttc	660
tgcaaacat	acaccagaac	taggattaag	cccattacgt	ggcttttagca	gaccgtccaa	720
aaatctgttt	tgcaagcacc	aattgtctct	tacttatcca	gcttcttttg	tgttggcaaa	780
ctgccctttt	ccaaccgatt	ttgtttcttc	tcacgctttc	ttcataaggct	aaactaacct	840
cggcgatcac	acaaccatgt	cctgaacctt	cacctcgctc	ctataaaagc	ccatccaacc	900
ttacaatctc	atcatcaccc	acaacaccga	gcaccccaat	ctacagatca	attcactgac	960
agttcactga	tctaga					976

<210> 18

<211> 1009

<212> DNA

<213> Artificial Sequence

<220>

<223> Glub-2 promoter

<400> 18

ctgcagtaat	ggatacctag	tagcaagcta	gcttaaacaa	atctaaattc	caatctgttc	60
gtaaacgttt	tctcgatcgc	aattttgatc	aaaactattg	aaaacctcaa	ttaaaccatt	120
caaaattttt	aataataccca	acaagagcgt	ccaaacccaa	tatgtaaata	tggatgtcat	180
gataattgac	ttatgacaat	gtgattatct	catcaagctc	ttaaatcatt	aattctagtt	240
gaaggtttat	gtttttcttat	gctaaagggt	tatgtttata	taagaatatt	aaagagcaaa	300
ttgcaataga	tcaacacaa	aaatttgaat	gtttccagat	gtgtaaaaat	atccaaatta	360
attgttttaa	aatagtttta	agaaggatct	gatatgcaag	tttgatagtt	agtaaactgc	420
aaaagggtct	attacatgga	aaattcctta	ttgaatatgt	ttcattgact	ggtttatttt	480
acatgacaac	aaagttacta	gtatgtcaat	aaaaaaat	aaggttactt	gtcaattgta	540
ttgtgccaa	taaagatgac	aacaaacata	caaatttatt	tgttctttta	tagaaacacc	600

taacttatca	aggatagttg	gccacgcaaa	aatgacaaca	tactttacaa	ttgtatcatc	660
ataaagatct	tatcaagtat	aagaacttta	tggtgacata	aaaaataatc	acaagggcaa	720
gacacatact	aaaagtattg	acagaaatth	cttaacaaac	tccatttggt	ttgtatccaa	780
aagcataaga	aatgagtcac	ggctgagtc	tgatatgtag	ttcaatcttg	caaaattgcc	840
tttttgttaa	gtattgtttt	aacactacaa	gtcacatatt	gtctatactt	gcaacaaaca	900
ctattaccgt	gtatcccaag	tggccttttc	attgctatat	aaactagctt	gatcggctct	960
tcaactcaca	tcaattagct	taagttttcca	ttagcaactg	ctaataagct		1009

<210> 19

<211> 839

<212> DNA

<213> Artificial Sequence

<220>

<223> Gt3 promoter

<400> 19

ctgcagtgtg	agtgtagctt	cttatagctt	agtgcctttac	tatcttcaca	agcacatgct	60
atagtattgt	tccaagatga	aagaataaatt	catccttgct	accaacttgc	atgatattat	120
atttgtgaat	atcctatctc	ttggcttata	atgaaatgtg	ctgctgggtt	attctgacca	180
tggtatttga	gagcctttgt	atagctgaaa	ccaacgtata	tcgagcatgg	aacagagaac	240
aaaaatgcaag	gattttttta	ttctgggtca	tgccctggat	gggttaatat	cgtgatcatc	300
aaaaaagata	tgcataaaat	taaagtaata	aatttgctca	taagaaacca	aaacaaaag	360
cacatatgtc	ctaaacaaac	tgcattttgt	ttgtcatgta	gcaatacaag	agataatata	420
tgacgtgggt	atgacttatt	cactttttgt	gactccaaaa	tgtagtaggt	ctaactgatt	480
gtttaaagt	atgtcttact	gtagaagttt	catcccaaaa	gcaatcacta	aagcaacaca	540
cacgtatagt	ccaccttcac	gtaattcctt	gtggaagata	acaagaaggc	tcactgaaaa	600
ataaaagcaa	agaaaaggat	atcaaacaga	ccattgtgca	tccatttgat	ccttgatgt	660
ctattttact	atcttccttt	tgtgtacctt	acttctatct	agtgagtcac	ttcatatgtg	720
gacattaaca	aactctatct	taacatctag	tcgatcacta	ctttacttca	ctataaaagg	780
accaacatat	atcatccatt	tctcacaaaa	gcattgagtt	cagtcccaca	aaatctaga	839

<210> 20

<211> 1302

<212> DNA

<213> Artificial Sequence

<220>

<223> Glub-1 promoter

<400> 20

ctgcagagat	atggattttc	taagattaat	tgattctctg	tctaaagaaa	aaaagtatta	60
ttgaattaaa	tggaaaaaga	aaaaggaaaa	aggggatggc	ttctgctttt	tgggctgaag	120
gcggcgtgtg	gccagcgtgc	tgctgctgga	cagcgagcga	acacacgacg	gagcagctac	180
gacgaacggg	ggaccgagtg	gaccggacga	ggatgtggcc	taggacgagt	gcacaaggct	240
agtggactcg	gtccccgcgc	ggtatcccga	gtggctccact	gtctgcaaac	acgattcaca	300
tagagcgggc	agacgcggga	gccgtcctag	gtgcaccgga	agcaaattccg	tcgcctgggt	360
ggatttgagt	gacacggccc	acgtgtagcc	tcacagctct	ccgtggctcag	atgtgtaaaa	420
ttatcataat	atgtgttttt	caaatagtta	aataatatat	ataggcaagt	tatatgggtc	480
aataagcagt	aaaaaggctt	atgacatggg	aaaattactt	acaccaatat	gccttactgt	540
ctgatataat	ttacatgaca	acaaagttac	aagtacgtca	tttaaaaaata	caagttactt	600
atcaatttga	gtgtatcaag	taaatgacaa	caaacctaca	aatttgctat	tttgaaggaa	660
cacttaaaaa	aatcaatagg	caagttatat	agtcaataaaa	ctgcaagaag	gcttatgaca	720
tggaaaaatt	acatacacca	atatgcttta	ttgtccggta	tatttttaca	gacaacaaag	780
ttataagtat	gtcattttaa	aatacaagtt	acttatcaat	tgtcaagtaa	atgaaaacaa	840
acctacaaat	ttgttatttt	gaaggaacac	ctaaattatc	aaatatagct	tgctacgcaa	900
aatgacaaca	tgcttacaa	ttattatcat	cttaaagtta	gactcatctt	ctcaagcata	960
agagctttat	gggtgcaaaaa	caaataataa	gacaaggcaa	agatacatat	atattaagag	1020
tatggacaga	catcttcttta	acaaactcca	tttgtattac	tccaaaagca	ccagaagttt	1080
gtcatggctg	agtcattgaa	tgtatagttc	aatcttgcaa	agttgccttt	ccttttgtac	1140
tgtgttttaa	cactacaagc	catatatatt	ctgtacgtgc	aacaaactat	atcaccatgt	1200
atcccaagat	gcttttttat	tgctatataa	actagcttgg	tctgtctttg	aactcacatc	1260
aattagctta	agttttccata	agcaagtaca	aatagctcta	ga		1302

<210> 21

<211> 675
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Rice proalmin promoter

<400> 21
 ctgcagcattc ggcttaggtg tagcaacacg actttattat tattattatt attattatta 60
 ttattttaca aaaatataaa atagatcagt ccctcaccac aagtagagca agttgggtgag 120
 ttattgtaaa gttctacaaa gctaatttaa aagttattgc attaacttat ttcattattac 180
 aaacaagagt gtcaatggaa caatgaaaac catatgacat actataattt tgtttttatt 240
 attgaaatta tataattcaa agagaataaa tccacatagc cgtaaaagttc tacatgtggt 300
 gcattaccaa aatatatata gcttacaaaa catgacaagc ttagtttgaa aaattgcaat 360
 ccttatcaca ttgacacata aagtgaagtga tgagtcataa tattattttt cttgctaccc 420
 atcatgtata tatgatagcc acaaagttac tttgatgatg atatcaaaga acattttttag 480
 gtgcacctaa cagaatatcc aaataatatg actcacttag atcataatag agcatcaagt 540
 aaaactaaca ctctaaagca accgatggga aagcatctat aaatagacaa gcacaatgaa 600
 aatcctcatc atccttcacc acaattcaaa tattatagtt gaagcatagt agtagaatcc 660
 aacaacaatc tagag 675

<210> 22
 <211> 1098
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Rice cysteine peptidase promoter

<400> 22
 ccaggcttca tcctaaccat tacaggcaag atgttgatg aagaaggggcg aacatgcaga 60
 ttgttaaact gacacgtgat ggacaagaat gaccgattgg tgaccggtct gacaatgggtc 120
 atgtcgtcag cagacagcca tctcccacgt cgcgcctgct tccggtgaaa gtggaggtag 180
 gtatgggccc tcccgtcaga aggtgattcg gatggcagcg atacaaatct ccgtccatta 240
 atgaagagaa gtcaagttga aagaaaggga gggagagatg gtgcatgtgg gatccccttg 300
 ggatataaaa ggaggacctt gccacttag aaaggagagg agaaagcaat cccagaagaa 360
 tcgggggctg actggcactt ttagcttct tcatcgcga atccaccaa acacaggagt 420
 agggatttac gcttctcagc ggcccgaacc tgtatacatc gcccggtgct tgtgtgtttc 480
 cgctcttgcg aaccttccac agattgggag cttagaacct caccagggc ccccgggcga 540
 actggcaaa gggggcctgc gcggtctccc ggtgaggagc ccacgctcc gtcagttcta 600
 aattaccgga tgagaaaggg aggggggggg gggaaatctg ccttgtttat ttacgatcca 660
 acggatttgg tcgacaccga tgaggtgtct taccagttac cacgagctag attatagtac 720
 taattacttg aggattcggg tcctaatttt ttaccgacg gacttcgcca tggaaaattt 780
 tttattcggg ggagaatatc caccctgttt cgctccta ataatagtag attgttacga 840
 ttagcaacct aattcagatc agaattgtta gttagcggcg ttggatccct cacctcatcc 900
 catcccaatt cccaaacca aactcctctt ccagtcgcgg acccaaacac gcattcgcgg 960
 cctataaatc ccacccgcat cgagcctatc aagcccaaaa aaccacaaac caaacgaaga 1020
 aggaaaaaaa aaggaggaaa agaaaagagg aggaaagcga agagggttga gagagacgct 1080
 cgtctccacg tcgcccgc 1098

<210> 23
 <211> 432
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Barley D-Hordian promoter

<400> 23
 cttcgagtgc ccgcccattt gccagcaatg gctaacagac acatattctg ccaaaacccc 60
 agaacaataa tcaattctcg tagatgaaga gaacagacca agatacaaac gtccacgctt 120
 cagcaaacag taccacagaa ctaggattaa gccgattacg cggcttttagc agaccgtcca 180
 aaaaaactgt tttgcaaaag tccaattcct ccttgcttat ccaatttctt ttgtgttggc 240
 aaactgcact tgtccaaccg attttgttct tcccgtgttt cttcttaggc taactaacac 300
 agccgtgcac atagccatgg tccggaatct tcacctcgtc cctataaaaag cccagccaat 360

ctccacaatc tcatcatcac cgagaacacc gagaaccaca aaactagaga tcaatttcatt 420
gacagtccac cg 432

<210> 24

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> bx7 signal peptide sequence

<400> 24

atggctaage gcctgggtcct ctttgccgga gtagtcgtcg ccctcgtggc tctcaccgcc 60

<210> 25

<211> 72

<212> DNA

<213> Artificial Sequence

<220>

<223> Glub-2 signal peptide sequence

<400> 25

atggcaacta ccatttttctc tcgtttttct atatactttt gtgctatgct attatgccag 60
ggttctatgg cc 72

<210> 26

<211> 85

<212> DNA

<213> Artificial Sequence

<220>

<223> Gt3 signal peptide sequence

<400> 26

atgtggacat taacaaactc tatcttaaca tctagtcgat cactacttta cttcactata 60
aaaggaccaa catatatcat ccatt 85

<210> 27

<211> 72

<212> DNA

<213> Artificial Sequence

<220>

<223> Glub-1 signal peptide sequence

<400> 27

atggcgagtt ccgtttttctc tcgtttttct atatactttt gtgttcttct attatgccat 60
ggttctatgg cc 72

<210> 28

<211> 69

<212> DNA

<213> Artificial Sequence

<220>

<223> proalmin signal peptide sequence

<400> 28

atgaagatca ttttcgtatt tgctctcctt gctattgttg catgcaacgc ttctgcacgg 60
tttgatgct 69

<210> 29

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Rice cysteine peptidase signal peptide sequence

<400> 29

atggccgccc gcgcgcgcgc cgcgcggttc ctgctgctgc tcatcgctcg tggtcaccgc 60
gcc 63

<210> 30

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> D-Hodian signal peptide sequence

<400> 30

atggctaagc ggctggctct ctttgtggcg gtaatcgctg ccctcgctgg tctcaccacc 60
gcc 63

<210> 31

<211> 1314

<212> DNA

<213> Artificial Sequence

<220>

<223> O2 transcription factor sequence

<400> 31

atggagcacg	tcatctcaat	ggaggagatc	ctcgggcccct	tctgggagct	gctaccaccg	60
ccagcgccag	agccagagcg	agagcagcct	ccggtaaccg	gcacgctcgt	cggcagtgct	120
atagacgttg	ctgctgctgg	tcatggtgac	ggggacatga	tggatcagca	gcacgccaca	180
gagtggaact	ttgagagggt	actagaagag	gaggctctga	cgacaagcac	accgcccgcg	240
gtggtggtgg	tgccgaactc	ttgttgctca	ggcgccctaa	atgctgaccg	gccgcccgtg	300
atggaagagg	cggtaactat	ggcgccctgc	gcggtgagta	gtgccgtagt	aggtgacccc	360
atggagtaca	atgccatact	gaggagggaag	ctggaggagg	acctcgaggc	cttcaaaatg	420
tggaggggcg	cctccagtg	tgtgacctca	gatcaacgtt	ctcaaggctc	aaacaatcac	480
actggaggta	gcagcatcag	gaataatcca	gtgcagaaca	agctgatgaa	cggcgaagat	540
ccaatcaaca	ataaccacgc	tcaaactgca	ggccttggcg	tgaggcttgc	tactagctct	600
tcctcgagag	atccttcacc	atcagacgaa	gacatggacg	gagaagtaga	gattctgggg	660
ttcaagatgc	ctaccgagga	aagagtggag	aaaagaaaagg	aatccaatag	agaatcagcc	720
agacgctcga	gatacaggaa	agccgctcac	ctgaaagaac	tggaagacca	ggtagcacag	780
ctaaaagccg	agaattcttg	cctgctgagg	cgcatcgccg	ctctgaacca	gaagtacaac	840
gacgctaacg	tcgacaacag	ggtgctgaga	gcggacatgg	agaccctaag	agctaagggtg	900
aagatggggg	aggactctct	gaagcgggtg	atagagatga	gctcatcagt	gccgtcgtcc	960
atgcccactc	cggcgccgac	ccccagctcc	gacgctccag	tgccgcccgc	gcctatccga	1020
gacagcatcg	tcggctactt	ctccgccaca	gccgcagacg	acgatgcttc	ggtcggcaac	1080
ggtttcttgc	gactgcaagc	tcatcaagag	cctgcatcca	tggtcgctcg	tggaactctg	1140
agcgccacag	agatgaaccg	agtagcagca	gccacgcatt	gcgcgggggc	catggagcac	1200
atccagacgg	cgatgggatc	catgccgcgc	acctccgect	ccggatctac	accgcccgcg	1260
caggattatg	agctgctggg	tccaaatggg	gccatacaca	tgacatgta	ttag	1314

<210> 32

<211> 987

<212> DNA

<213> Artificial Sequence

<220>

<223> PBF transcription factor sequence

<400> 32

atggacatga	tctccggcag	cactgcagca	acatcaacac	cccacaacaa	ccaacaggcg	60
gtgatgttgt	catcccccat	tataaaggag	gaagctaggg	acccaaagca	gacacgagcc	120
atgccccaaa	taggtggcag	tggggagcgt	aagccgaggc	cgcaactacc	tgaggcgctc	180

aagtgtccac	gctgcgactc	caacaacacc	aagttttgct	actacaacaa	ttatagcatg	240
tcacaaccac	gctacttttg	caaggcttgc	cgccgctatt	ggacacatgg	tgggtaccctc	300
cgcaatgtcc	ccattgggtg	tgggtgtcgc	aagaacaaac	atgcctctag	atttgtcttg	360
ggctctcaca	cctcatcgtc	ctcatctgct	acctatgcac	cattatcccc	tagcaccaac	420
gctagctcta	gcaatatgag	catcaacaaa	catatgatga	tgggtgcctaa	catgacgatg	480
cctaccccaa	cgacaatggg	cttattccct	aatgtgctcc	caacacttat	gccgacaggt	540
ggaggcgggg	gctttgactt	cactatggac	aaccaacata	gatcattgtc	cttcacacca	600
atgtctctac	ctagccaggg	gccagtgcct	atgctggctg	caggaggagg	tgaggcaaca	660
ccgtctttcc	tagagatgct	gagaggaggg	atttttcatg	gtagtagtag	ctataacaca	720
agtctcacga	tgagtgggtg	caacaatgga	atggacaagc	cattttcgct	gccatcatat	780
ggtgcaatgt	gcacaaatgg	gttgagtggc	tcaaccacta	atgatgccag	acaactggtg	840
gggcctcagc	aggataacaa	ggccatcatg	aagagcagta	ataacaacaa	tgggtgatca	900
ttgttgaaac	tctactggaa	caagcacaac	aacaacaaca	acaacaacaa	caacaacaac	960
aacaacaaca	acaacaaggg	acaataa				987

<210> 33

<211> 3902

<212> DNA

<213> Artificial Sequence

<220>

<223> Reb transcription factor sequence

<400> 33

atggagcggg	tgttctccgt	ggaggagatc	tccgacccat	tctgggtccc	gcctccgcgcg	60
ccgcagtcgg	cggcgggcggc	ccagcagcag	ggcgggcggcg	gcgtggcttc	gggaggtggt	120
ggtggtgtag	cggggggcgg	cggcgggcggg	aacgcgatga	accggtgccc	gtcggagtgg	180
tacttccaga	agtttcttga	ggaggcgggtg	ctcgatagcc	ccgtcccga	ccctagcccg	240
agggtccgaag	cgggagggat	cagggggcgca	ggagggtggg	tgccgggtcga	tgttaagcag	300
ccgcagctct	cggcgggcggc	gacgacgagc	gcgggtgggtg	accccggtgga	gtacaacgcg	360
atgctgaagc	agaagctgga	gaaggacctc	gccgcggtcg	ccatgtggag	ggtacagcca	420
ttctcccccc	ctctagtact	cgagagctta	ctgagatcgg	caatgctagc	tactgtttgc	480
atcgaatggt	tataggtatt	tagatcgggc	atttctatag	accaatggcg	tccatggtct	540
tgcaatgcgc	ctgtgtgagt	gtcgggtggtt	ggttcgactc	atagtatgta	gggttgtgcg	600
tatgtacaaa	cgggaagcttc	atagacctcg	gtattgagat	tgcgatatcg	atgcaacctg	660
cgaattggcg	atgtaatcag	tcatattctt	actaaactgc	gagacagtgg	tttgtttgca	720
attgcaatat	ttttgtatgg	ggctgcttaa	actgtcattg	cctttttaga	ttggcaatat	780
gtgactttat	gcaagtattt	gattgggcgg	atccaggaac	aaaaagttgg	ggggattcaa	840
cataccgagt	acactggcat	aaacacatca	tctcagttat	aaactatgct	aaaatgctat	900
taagagacct	ttagcacctc	ttatcttata	aacctgggtg	aaaaaattga	aggggggact	960
caggggggta	tccatgggtc	cgatgggtgc	aggggggact	gagtccccc	tgccccacg	1020
ttgaatccgc	cctggcatgc	gtataagctg	tcacagccat	ttctaggtgc	ttgtgcttag	1080
ttgggtgatg	tcagcttaat	ttgtcttttc	tatgtcgtca	tcgattttct	aagaaacgaa	1140
aaatagccta	tttatgtgct	ccagaatttg	atgatccctg	gcccttcatt	tgctgaaatt	1200
agcctatttg	ttggttgccc	ttcagttttt	tcccagctta	tgttggttga	atgtgtggct	1260
atgcctcggt	ttgtgccccta	taatttatta	tttgcaattc	atttttgtac	atgacttaaa	1320
atgacactag	agcaacatgc	actgattggt	tatcctataa	tcatttatgt	agttctgttc	1380
attttatcat	gctagctcat	gtcattttca	tcttcaggcc	tctggcacag	ttccacctga	1440
gcgtcctgga	gctggttcat	ccttgctgaa	tcagatggtt	tcacacatag	gcgctcctaa	1500
ttccatcgga	ggtacttatc	ttatctggtt	acattttcag	attgttatga	aactacccaa	1560
atatacctgca	caattgcatg	ggattaaatt	ttagtttctt	tgaatataga	gtagagttgt	1620
attgctgtca	cgatcatcaa	tagttctgaa	gctatgaata	aataagttcc	gcatttgtta	1680
gtgattcttt	gaacattaga	attgttatgc	ttaagtagat	agggttatgt	ttgtttggag	1740
ttcccttaaa	tcatttcatt	gctgactgcc	agctggcagg	agcatttggt	gttgccctga	1800
ccatgaatga	agaccttcct	gttctgagtg	ctcacaagaa	aacatatttt	gattaatgca	1860
ccttgaatcc	ttaggatctt	gcaaagatgg	gcacttagct	ttagaattga	gtagtactta	1920
aatagctggt	gttatcatga	tttgtcctgt	agtgaatgt	cgacaaaaca	ggaatgctac	1980
ttttgacttc	tgatatttca	tgcctggctt	tacttatgct	ctgtttggaa	catgggcaca	2040
tatcaggcaa	tgctactcca	gttcaaaaca	tgctaagtg	cccaagtggg	ggatcgggct	2100
cacagttggg	acagaatggt	gatgtccttg	taaagcagcc	caccagctct	tcatacaagg	2160
agcagtcaga	tgatgatgac	atgaaggag	aagctgagac	cactggaact	gcaagacctg	2220
ctgatcaaag	attacaacga	aggtgatcat	tcattgcttc	cttgtaatat	agattctgta	2280
cataattaac	ctacctcgtc	atgcatgcat	gtgtcctatt	ttcaccttag	ccctttcagt	2340
tggatttcca	ctttcatccg	gtagcctttc	agtttccctat	tgcacgcgat	atatgatctt	2400
ttacctacca	tattagttct	ctgtgtgcca	tactcagtgc	ttagtgtctc	gagcaagaga	2460

ggaatattgta	tggctattac	acgtagcact	ttgctctcta	cttgtttatt	gacataagca	2520
atttgggatg	aattaaatct	gagttcacat	catattcett	atgtcacaa	tttctgaaac	2580
cgattgtatc	tagtatctgg	ttgatgcacc	cccatcttgg	atttgcaaat	caaagttata	2640
ctccctagag	agctttacct	ttcataaagc	aattacccca	ataaaccacg	gatttgatag	2700
ctattgacta	tgattaccag	aattcatttg	gcagctattt	tctcaattta	agtttggtat	2760
tagtctcagt	tggctgtaaa	ataatgtcac	ggtagggtac	atgtatgtgc	agcatacaag	2820
gtatgggtga	gttatgatat	ggacagtgtg	tacacccac	atttgctcac	taaaatcaaa	2880
atattcaaac	gtcacgtgat	gatatgggtg	attgcattat	accttgtatt	gtttattatg	2940
ttacttgtgc	tagacaataa	tataggctgt	tcttttgggt	gattttgtat	gaagatgttg	3000
agcaagcact	tctcgatata	atgctagttt	tgttgacctg	ttccaggaag	caatccaatc	3060
gggagtcagc	caggcgctca	agaagcagaa	aggcagctca	cttgaatgag	ctggaggcac	3120
aggtgtgata	gttcacatag	ttattttcga	taagacataa	aatcctaaat	tactggctac	3180
tgacttcagt	tatggattta	cttggttacag	gtatcgcaat	taagagtcga	gaactcctcg	3240
ctgttaaggc	gtcttgctga	tgtaaccag	aagtacaatg	atgctgctgt	tgacaataga	3300
gtgctaaaag	cagatgttga	gaccttgaga	gcaaaggtat	gctatatatg	ccttttgcaa	3360
tatgcatccc	atggattgct	actttggctt	gtttcaaaact	ttcaacgtga	cttgtgtacc	3420
ctgttattag	aagaataatc	ccgcctacca	ttatactcta	taaatcacca	tttgccagct	3480
ccaaacatga	ttattaaatc	aggtcaatct	gaacattgaa	atgtatcaaa	aattcgcagg	3540
tgaagatggc	agaggactcg	gtgaagcggg	tgacaggcat	gaacgcgttg	tttccgcgcg	3600
cttctgatat	gtcatccctc	agcatgccat	tcaacagctc	cccatctgaa	gcaacgtcag	3660
acgctgctgt	tcccatccaa	gatgaccgga	acaattactt	cgctactaac	aacgacatcg	3720
gaggtaacaa	caactacatg	cccgacatac	cttcttcggc	tcaggaggac	gaggacttcg	3780
tcaatggcgc	tctggctgcc	ggcaagattg	gccggccagc	ctcgctgcag	cgggtggcga	3840
gcctggagca	tctccagaag	aggatgtgcg	gtgggcccgc	ttcgtctggg	tcgacgtcct	3900
ga						3902

<210> 34

<211> 361

<212> DNA

<213> Bombyx mori

<400> 34

atcagtcgct	tagttcccac	agcaacagct	atatcgacag	caaaactcct	ttttaaaatc	60
caggaattaa	atctaaaat	ttcaaaatga	acttcaacaa	gatcttcgtg	ttcgttgccc	120
tcctcctggc	tattagtttg	ggacaatctg	aagccgggtg	gctgaagaaa	ttgggcaaaa	180
gaattgtaag	tattgctaaa	aacaaatatt	aaaatgtttt	aatatctcta	aattaaatat	240
taattatttt	taaaggaacg	cattggccag	cacactcgag	atgccaccat	tcagggtctg	300
ggaattgcgc	aacaagcggc	caatgtggca	gccacggcca	gaggatgaaa	gttccatcaa	360
a						361

<210> 35

<211> 1098

<212> DNA

<213> Xenopus laevis

<400> 35

ttgaaataca	actatatattg	gaaaagatgt	tcaaaggatt	atztatctgt	tcactaattg	60
ctgtgatctg	tgcaaatgca	ctgccacaac	cagaggcctc	tgcggatgaa	gatatggatg	120
aaagagaggt	ccggggaatt	ggtaaatttt	tgcatctcagc	gggcaaattt	ggaaaagctt	180
ttgtgggaga	gataatgaag	tcaaaacgag	atgcagaagc	agtaggacca	gaggcctttg	240
cagatgaaga	tttagatgaa	agagaggtcc	ggggaatttg	taaaattttg	cactcagcaa	300
aaaaattttg	aaaagctttt	gtgggagaga	taatgaattc	aaaacgagat	gcagaagcag	360
taggaccaga	ggcctttgca	gatgaagatt	tagatgaaag	agaggtccgg	ggaattggta	420
aattttttgca	ctcagcaaaa	aaattttgga	aagcttttgt	gggagaaata	atgaattcaa	480
aacgagatgc	agaagcagta	ggaccagagg	cctttgcaga	tgaagattta	gatgaaagag	540
aggtccgggg	aatttgtaaa	tttttgcact	cagcaaaaaa	atttggaaaa	gcttttgttg	600
gagaaataat	gaattcaaaa	cgagatgcag	aagcagtagg	accagaggcc	tttgcagatg	660
aagattttcg	tgaaagagag	gtccggggaa	ttggtaaatt	tttgactca	gcaaaaaaat	720
ttggaaaagc	ttttgtggga	gaaataatga	attcaaaacg	agatgcagaa	gcagtaggac	780
cagagggcct	tgcagatgaa	gatttagatg	aaagagaggt	ccggggaatt	gttaaatttt	840
tgacttcagc	aaaaaaattt	ggaaaagctt	ttgtgggaga	aataatgaat	tcaaaacgag	900
atgcagaagc	agtagatgac	aggagatggg	ttgaatagtt	tcagcaacag	aaaaacatca	960
ctgaagaaca	atgattcccc	tgaaaatgtg	taaacggctg	aaatgattct	gaacatacta	1020
gtgctgaata	aaaagaagat	cccttagata	acctatacct	acagtatata	tacctacaaa	1080
taaactgagc	atcacaaat					1098

<210> 36
 <211> 297
 <212> DNA
 <213> Mamestra brassicae

<400> 36
 atgctgtgcc tcgctgacat tcgtatcgtg gcttcgtgtt ccgccgccat taagagtgga 60
 tacggacagc aaccgtggct ggcccacggt gcaggccctt atgccaaactc tctattcgat 120
 gatgtgcccc cggatagcta tcacgcggcc gtcgagtact tgcgcctgat acccgccagt 180
 tgttacctgc tagacggata cgccgccggt cgtgatgacg gtagggctca ttgcatagcc 240
 ccacgcaacc gccgactata ctgtgcctcg tatcaggctc gcgtctgtcg atattga 297

<210> 37
 <211> 524
 <212> DNA
 <213> Tachypleus tridentatus

<400> 37
 gcctaacatc gcctcagcaa cttggccaat agtttcgaac ttcacgtca ctatgaagaa 60
 acttgtaatt gctttgtgcc tgatgatggt attggctgtg atgggtggaag aggccgaggc 120
 taaatgggtgc ttcagagtat gctacagggg aatttggtat cgcagatgtc gaggggaagag 180
 gaatgaagta cgtcagtacc gtgaccgtgg gtatgatgta agagccatac cagaagaaac 240
 gttctttaca cgacaagacg aagatgaaga tgacgatgaa gagtgaggcg tttacagctg 300
 aaggaagtat ccagctgatg aatattggcg aagacttgga acgaatgact tttacgattt 360
 ttaaacctcg ttaaagggtt aatatttttg tatattgggtg ttcattaaaa cacattcttt 420
 tatcactatt tcttctttgt gcacaaaagta aacttcatat actttctcta ctatttggat 480
 tgttggtaaa gtttgatatt ataataaata tataacaagc attc 524

<210> 38
 <211> 19
 <212> PRT
 <213> Parasilurus asotus

<400> 38
 Lys Gly Arg Gly Lys Gln Gly Gly Lys Val Arg Ala Lys Ala Lys Thr
 1 5 10 15
 Arg Ser Ser

<210> 39
 <211> 129
 <212> PRT
 <213> Bufo bufo gagarizans

<400> 39
 Met Ser Gly Arg Gly Lys Gln Gly Gly Lys Val Arg Ala Lys Ala Lys
 1 5 10 15
 Thr Arg Ser Ser Arg Ala Gly Leu Gln Phe Pro Val Gly Arg Val His
 20 25 30
 Arg Leu Leu Arg Lys Gly Asn Tyr Ala Gln Arg Val Gly Ala Gly Ala
 35 40 45
 Pro Val Tyr Leu Ala Ala Val Leu Glu Tyr Leu Thr Ala Glu Ile Leu
 50 55 60
 Glu Leu Ala Gly Asn Ala Ala Arg Asp Asn Lys Lys Thr Arg Ile Ile
 65 70 75 80
 Pro Arg His Leu Gln Leu Ala Val Arg Asn Asp Glu Glu Leu Asn Lys
 85 90 95
 Leu Leu Gly Gly Val Thr Ile Ala Gln Gly Gly Val Leu Pro Asn Ile
 100 105 110
 Gln Ala Val Leu Leu Pro Lys Thr Glu Ser Ser Lys Pro Ala Lys Ser
 115 120 125
 Lys

<210> 40
 <211> 466
 <212> DNA
 <213> Bufo bufo gagarizans

<400> 40
 gtctgtacag agaagagaac gatgtctgga cgcggcaaac aaggaggcaa agtgcgggct 60
 aaggccaaga cccgctcatc ccgggcaggc ctccagttcc cggtcggccg tgtgcacagg 120
 ctccctccgca agggcaacta cgcccagagg gtgggcgcgc gcgctcccgt ctacttgccc 180
 gctgtgctcg agtatctcac cgccgagatc ctggagctgg ccggcaatgc cgcccgcgac 240
 aacaagaaga ctgcgcatcat cccccgtcac ctgcagctgg ccgtgcgcaa cgacgaggag 300
 ctcaacaagc tgctgggcgg agtcaccatc gccagggggg gggtcctgcc caacatccag 360
 gccgtgctgc tgcccaagac cgagagcagc aagccggcca agagcaagtg agccccgctc 420
 tgccccgaac caaaggctct tctaagaaaa aaaaaaaaaa aaaaaa 466

<210> 41
 <211> 581
 <212> DNA
 <213> Sus scrofa

<400> 41
 tcacctgggc accatggaga cccagagggc cagcctgtgc ctggggcgct ggctactgtg 60
 gcttctgctg ctgggactcg tgggtgccctc ggccagcgcc caggccctca gctacaggga 120
 ggccgtgctt cgtgctgtgg atcgctcaa cgagcagtc tgggaagcta atctctaccg 180
 cctcctggag ctggaccagc cgcccaaggc cgacgaggac ccgggcaccc caaaacctgt 240
 gagcttcacg gtgaaggaga ctgtgtgtcc caggccgacc cggcagcccc cggagctgtg 300
 tgacttcaag gagaacgggc ggggtgaagca gtgtgtgggg acagtcacct tgaaagagat 360
 cagaggcaac tttgacataa cctgtaatca gcttcagagt gtcaggatta tagacctgct 420
 gtggagagta cgtcggccac agaaacccaa atttgtgact gtatgggtca gataattagt 480
 gcatgaagga atttcattct tcaagacaca gggaagagtt cttaaagttct gcttattcta 540
 ccacaagctt caggactgga aaaataaatt cttgtgaaag c 581

<210> 42
 <211> 23
 <212> PRT
 <213> Morone sp.

<400> 42
 Phe Phe His His Ile Phe Arg Gly Ile Val His Val Gly Lys Thr Ile
 1 5 10 15
 His Asp Leu Val Thr Gly Thr
 20

<210> 43
 <211> 484
 <212> DNA
 <213> Morone sp.

<400> 43
 agagagattt acaccacctg tacatctcaa accgcttctt aagctcactt tgataacagc 60
 tttttgactt tcagtcacag tcggtggaaa gaccgaagag atgaagtgcg ccacgctctc 120
 tcttgctgctg tcgatggctg tcctcatggc tgaaocctggg gatgccttct ttcaccacat 180
 tttccgtgga attgttcacg tcggcaagac gatccacaaa cttgtgaccg ggggaaaagc 240
 ggagcaagat cagcaagatc agcaatatca gcaagatcag caagatcagc aagcgcagca 300
 atatcagcgc ttttaaccgcg agcgtgcggc ttttgactag actttgtgaa tctatggcgg 360
 ttcattctgaa agagcagctc tgaaactttg ttttaatta cattcttggt gctctttaac 420
 tgatatttga acatatatca gttatttggg ataaatatgc atatctaaaa aaaaaaaaaa 480
 aaaa 484

<210> 44
 <211> 10
 <212> PRT
 <213> Anoplius samariensis

<400> 44

Gly Leu Leu Lys Arg Ile Lys Thr Leu Leu
 1 5 10

<210> 45
 <211> 2879
 <212> DNA
 <213> Homo sapiens

<400> 45
 ttttttcata ctgagtctca ctctgttacc caggctggag tgcagtggca tgatctcagc 60
 taactgcaac ttctgcttcc cgggttcaat ggggttcaagt gattctcatg cctcagcttg 120
 tagctgggaa ctactggtgt gagccatcat gcgtggctaa ttttcatatt tttagtagag 180
 atggggtttc accatgttgg ccaagcttgt ctggaactcc ttatctcagg tgatccgccc 240
 accttggcct cccaaagtgc tgggattata ggctgacaga ccgtgccctg cctcattcat 300
 caattcttaa tcgatgccta cagggtgcca ggcaatgcta gagctggaga tttagcagtc 360
 catcatactg actcctgagg agtagaagga tgtagaatag gcacctggct ctcttccctc 420
 ctggagggat ttaacgctct tgagcaccctc tggctatgac aatctccggt cagggtctggg 480
 aggttgtcag agatgaagaa accacttccct catcttgcac acaaggaagg cctcactcac 540
 tgcccagcaa gtccctgtgaa gcaatagcca gggctaaagc aaaccccagc ccacaccctg 600
 gcaggcagcc agggatgggt ggatcaggaa ggctcctggt tgggcttttg catcaggctc 660
 aggtctggga taaaggaggc tctgtgtggc tagaggaggc cagacatggg gaccatgaag 720
 acccaaaggg atggccactc cctggggcgg tggctactgc tgctcctgct cctgggcctg 780
 gtgatgcctc tggccatcat tggccaggtc ctcagctaca aggaagctgt gcttcgtgct 840
 atagatggca tcaaccagcg gtccctcggat gctaacctct accgcctcct ggacctggac 900
 cccaggccca cgatggtgag ctttggggga cattctgctc tgctctggct gggcttggca 960
 cgtgttgttc cttctgctcc tgctgcactg cctgccagga gggcatctcc ccctttaaat 1020
 gtggctcccg gttttccagg gaaccttcta gagctcgtgt ctctcccag ctcgagagct 1080
 tctgccttat aattcctgct gtggcagaga tccctcacc cgaacccacg caggttttgg 1140
 gacttctgcg agctccaggc actagaatgg ggtcattggc tctgggcagt gacctcctc 1200
 gctttaagtc tcttctgtac cacgttacc cecataggga agaactctat ccagacttta 1260
 ggttccagtg ggcattgtct gtccccagg aagccccctg acttcccttg cccccacccc 1320
 agagtgggag gggctccttg ttagagctca tctgagggtc gctcctactc actgttcacc 1380
 taggagggta ggaattggctc agtccctcct cctcaatgc cccagtgcga agcccagcac 1440
 ccagtgcctg tcgcacatca ggtactgtgg aaagcctgcc ctcttgggtg ggaggctcatg 1500
 gacacaaatc agaaaataca agaattgggc tccccatttc ctctctgac taggatgggg 1560
 acccagacac gccaaagcct gtgagcttca cagtgaagga gacagtgtgc cccaggacga 1620
 cacagcagtc accagaggat tgtgacttca agaaggacgg ggtgaggctg ggggctgggg 1680
 gtgttgggtg gtgcctccca aggagctgaa cagggggcac ctgggggaata tttcccactg 1740
 ggatgtggct gggagggtcat ggcaaatggg ttcaagtttg accttgagct tctcctttcc 1800
 agctggtgaa gcggtgtatg gggacagtga cctcaacca ggcagggggc tccattgaca 1860
 tcagttgtga taagggtgagt gggctgttct gggatgcagg ggctgatggg gccatagagt 1920
 gtggaccatc caatgggtca attactact ccccaacccc aggcagaga aagccccctc 1980
 taccaggggc tcttccccaa acctgagttc catctccagg gccggctctg gaatccctta 2040
 gagcggtaga tctccaagtg tagcccttcc tggggactcg ttagatatgc aaattctcag 2100
 gccctactca gacctactca gacagactct gggtagggcc cagaaattcg tattttgata 2160
 agcttccagg gagattccgg cttctgtaaa gtttgagagc cactgtctaa gagtactcag 2220
 ctctcagccc tgtgttccca tctcagttgt gctgggctgg gctgtgtgac cctgcagagc 2280
 cctcactat ctccgggact ctgttttctc atctttttat tgggtgtagg gattcaatca 2340
 catgcttcaa aggtcacagc cagaggttga actggggccc caaagctctg cggggggcca 2400
 cgaagagggg cgtctaggtg gggaggggtc ttggattgac cctgggtaca tccccgacaa 2460
 ggaacctgtt tcttctctgta cacaacccca ggataacaag agatttgccc tgctgggtga 2520
 tttcttccgg aatatctaaag agaagattgg caaagagttt aaaagaattg tccagagaat 2580
 caaggatttt ttgcggaatc ttgtaccag gacagagtc tagtgtgtgc cctaccctgg 2640
 ctcaggcttc tgggctctga gaaataaact atgagagcaa tttcctcagg cttcagttc 2700
 acttgttttg cctcctctct ctcaccacaa ctgagccctt agctcaggga gtccacgtgt 2760
 gagtgtgagt gtgtgtgagt gtgacacaga ggtggcgagg gcagtgttcc atccaggagg 2820
 acacagggta aggcagtagg gccaaagagat ccaagatggc attcccatte tcagtggaa 2879

<210> 46
 <211> 6836
 <212> DNA
 <213> Homo sapiens

<400> 46
 gttggcctta gatgtaacac agggagagct catccctttt aatagaggaa gacagaaagg 60

ctggtacatt	gggttgtagg	aggatatata	aggatatcag	ttaggtgtta	agaacagaaa	120
gactgagtta	acagtgattt	aacaaaagag	gagagtattt	ctcctttaag	taaattttca	180
ggtaggcagc	ctagagctga	tgcaagtggc	ccatacccaa	ttatccaggc	tccatctatc	240
ttgttgccat	gccaacctca	acatgcagct	tacacttgct	ggcctgaaag	tgaactattc	300
catctcctgc	tatcatgact	gaatactagt	caaagggaaa	aagggggaag	tgggagggga	360
aagcacatac	ctttcttaaa	gagcacaacc	tagagatggt	ccctattctg	ttcatatctc	420
gttggcctta	atttaatcac	atcatcactg	caagctacag	tgggggctgg	gaaatttggt	480
gtttgagggg	atgaccaagc	gcccactga	aaatgttatt	cttctcttat	gagaagttaga	540
gaatggatgt	tagggaatgg	ctagaagact	ctactacaaa	ggatacgggtg	gttcttttct	600
gattgcttct	aacttctcaa	tgaaggtga	gttcatcagc	tgaatgaaag	aataaatgag	660
attttgggga	cttgaggagc	aaggagaagg	agtataatag	tcattttgaa	gagtgagtta	720
agtagggaac	tgtagtaaga	ttgacagaca	gtgtggaggg	attacttgaa	tcttgtgaat	780
agaggaaaga	gtagaatcag	attatcctga	ctcctgcctg	aagctttaca	tattcagaga	840
aaaatgttgg	aagaaacttt	gatataatgc	tatgtctgtg	atcaggcaca	cattttactg	900
tacttttact	gtcaggggccg	tcatttagtg	ccaagatgtc	tagagagttc	gttcttttct	960
tactcaattg	gctgagaaaa	tgtgtccatg	caaaaaacca	aacaccgcat	gttctcactc	1020
atagatggga	attgaacaat	gagaatactt	ggacacagga	aggggaacat	cacactctgg	1080
ggactgttgt	ggggtggggg	gaggggggag	ggatagcatt	agaagatata	cctaattgcta	1140
aatgatgagt	taatgggtgc	agcacaccag	catggcacat	gtatacatat	gtaactaacc	1200
tgcacattgt	gcacatgtac	cctaaaactt	aaagtataat	aataataaaa	aatgtgtgcc	1260
atggctcttg	gaggagcatg	tttgttttcc	tcattttcca	gtctgtaaat	aagcaaatgt	1320
aaaggggtta	gtgataatgt	ccatctccag	aagctgtcag	atttcctttg	tcaaatctta	1380
tgatttgggc	tgaagtaggt	gttgagaggg	cagctaccac	gtgcacccag	atggccactc	1440
gtttaatatg	ttaccatttc	ccattatatt	cgcaggatag	atagccaaag	tggagccctg	1500
agagatttct	tcatttttcc	tgtcataaag	aatttgtaat	tcagtagtca	taggagtttg	1560
taataaataa	ctcacattga	tttctctgtt	ctgaaataat	tttgcttccc	ctcttcccga	1620
agctctgaca	cctgccccaa	caagcaatgt	tggaaaatta	tttacaatgt	ggcgcaaat	1680
cccttactgc	tttggatata	aatccaggca	ggaggaggtg	gctctaaggc	aagagactta	1740
ggacttctag	cccctgaact	ttcagccgaa	tacatctttt	ccaaaggagt	gaattcaggc	1800
ccttgatca	ctggcagcag	gacgtgacca	tggagaagct	gttgtgtttc	ttggtcttga	1860
ccagcctctc	tcattgctttt	ggccagacag	gtaagggccca	ccccaggcta	tgggagagat	1920
ttgatctgag	gtatgggggt	gggtgtctaa	actgcatgaa	cagtctcaaa	aaaaaaaaaa	1980
aaagactgta	tgaacagaac	agtggagcat	ccttcatggt	gtgtgtgtgt	gtgtgtgtgt	2040
gtgtgtgtgt	gtgtgtgtgt	tgtactgtga	gaaggggtca	gtctgtttct	caatcttaaa	2100
ttctatacgt	aagttagggg	atagatctgt	gtgatctgag	aaacctctca	catttgcttg	2160
tttttctggc	tcacagacat	gtcgaggaag	gcttttgtgt	ttcccaaaga	gtcggatact	2220
tcctatgtat	ccctcaaagc	accgttaacg	aagcctctca	aagccttcac	tgtgtgcctc	2280
cacttctaca	cggaaactgtc	ctcgacccgt	gggtacagta	ttttctcgta	tgccaccaag	2340
agacaagaca	atgagattct	catatttttg	tctaaggata	taggatacag	ttttacagtg	2400
ggtgggtctg	aaatattatt	cgaggttctc	gaagtccag	tagctccagt	acacatttgt	2460
acaagctggg	agtccgcctc	agggatcgtg	gagttctggg	tagatgggaa	gcccagggtg	2520
aggaagagtc	tgaagaaggg	atacactgtg	ggggcagaag	caagcatcat	cctggggcag	2580
gagcaggatt	ccttcggtgg	gaactttgaa	ggaagccagt	ccctgggtgg	agacattgga	2640
aatgtgaaca	tgtgggactt	tgtgctgtca	ccagatgaga	ttaacaccat	ctatcttggc	2700
gggcccttca	gtcctaattgt	cctgaactgg	cgggcactga	agtatgaagt	gcaaggcgaa	2760
gtgtttacca	aaccacagct	gtggccctga	ggccagctg	tgggtcctga	aggtacctcc	2820
cgttttttta	caccgcatgg	gccccacgtc	tctgtctctg	gtacctcccg	cttttttaca	2880
ctgcatgggt	cccacgtctc	tgtctctggg	cctttgttcc	cctatatgca	ttgcaggcct	2940
gctccaccct	cctcagcgcc	tgagaatgga	ggtaaagtgt	ctgggtctggg	agctcgtaa	3000
ctatgctggg	aaacggtcca	aaagaatcag	aatttgaggt	gttttgtttt	catttttatt	3060
tcaagttgga	cagatcttgg	agataatttc	ttacctcaca	tagatgagaa	aactaacacc	3120
cagaaaggag	aaatgatgtt	ataaaaaact	cataaggcaa	gagctgagaa	ggaagcgctg	3180
atcttctatt	taattcccca	cccatgaccc	ccagaaaagca	ggagggcatt	gcccacattc	3240
acagggtctc	tcagtctcag	aatcaggaca	ctggccagggt	gtctgggttg	ggtccagagt	3300
gctcatcatc	atgtcataga	actgctgggc	ccaggtctcc	tgaatggga	agcccagcaa	3360
taccacgcag	tccctccact	ttctcaaagc	acactggaaa	ggccattaga	attgccccag	3420
cagagcagat	ctgctttttt	tccagagcaa	aatgaagcac	taggtataaa	tatgttggtta	3480
ctgccaaaga	cttaaatgac	tggtttttgt	ttgcttgtag	tgttttctta	attttatggc	3540
tcttctggga	aactcctccc	cttttccaca	cgaacctgtg	ggggctgtga	attctttctt	3600
catccccgca	ttcccaatat	accagggcca	caagagtggg	cgtgaaccac	aggggtgctt	3660
gtcagaggag	cccatctccc	atctccccag	ctccctatct	ggaggatagt	tggatagtta	3720
cgtgttccta	gcaggaccaa	ctacagtctt	cccaaggatt	gagttatgga	ctttgggagt	3780
gagacatctt	cttgctgctg	gatttccaag	ctgagaggac	gtgaacctgg	gaccaccagt	3840
agccatcttg	tttgccacat	ggagagagac	tgtgaggaca	gaagccaaac	tgggaagtga	3900
ggagccaagg	gattgacaaa	caacagagcc	ttgaccacgt	ggagtctctg	aatcagcctt	3960

gtctggaacc	agatctacac	ctggactgcc	caggtctata	agccaataaa	gcccctgttt	4020
acttgagtga	gtccaagctg	ttttctgata	gttgcttttag	aagttgtgac	taacttctct	4080
aatgaccttt	gaaatcaagt	ccctgtaagt	ttacacttcc	cattacacaa	catatccatg	4140
gcactttaac	ctgtccaaga	gtcccctgaa	gggattccct	ctgacaagca	cagtcctctt	4200
tttgagatag	ctggccattc	atgtagggcc	agttgtgggg	gttttagtac	tctcattcta	4260
tcattattcc	aatttgagga	gcaaaaacta	aagatcattg	agatccatgg	atctgctgcc	4320
aggttcagtg	actggaatgc	acttgcccca	gcgggttcaa	gttctttgta	gaaacagacc	4380
cctcaaagat	gacagcagct	ggcttccctc	agaagtgtcc	ttcaacactg	tctttcagta	4440
ctaagtgacc	agtagaagtt	actccataaa	gaccatagcc	aggaatgaat	tatggggctt	4500
cctgcatatc	aaatggttac	ctaggggaagg	catatggtaa	actagttaag	cacagaggtt	4560
ctggaatcag	actgcctagg	tccaaatcct	ggctttacaa	cttactagct	atgtgacctt	4620
aggatgata	acctctttac	gcctcagttt	tcttacctac	aaagcaggag	aaataatagt	4680
aattagctca	aagggttgct	ataagaatca	aagaaaatca	tgcaagcaaa	gcacttacta	4740
cagtgcctgg	cacatagtaa	gctcttataa	cagttttctg	gtattattct	tattaagcca	4800
ttttgtatta	tatccataaa	tgtagattca	taaaatacat	aaaaatacaa	cttggtagaa	4860
tttgtcttgg	ttggcagcag	ttgaagcttc	tcatttgcta	agaattaggt	ttagaatggt	4920
agaaggggtga	acaacaagtc	tttgaagcca	ttatttatga	atattgactt	ctctaattga	4980
cttttgataa	ccatttgtaa	gcatagaagc	taaggataca	ttctcagaca	aaaatgagag	5040
tgcaaatact	ctctgttgct	caattagggc	agatccatcc	atcctcacat	tcagctaagc	5100
agaagagagg	attgacactc	ttttctagtt	aaataatcca	gatacgtctt	aattatagaa	5160
ggccatccca	actccccagg	gatgtaggtt	gagctaatat	tttcttcaga	attcagttgc	5220
ttgcatctta	ctatacatag	tctggtagac	agcctctggc	cttaataatt	tgagccagga	5280
aaatcttaac	agctggtatt	tgcatccaga	gaacattaaa	aaatcttaac	aaagcattcg	5340
taggatgttt	agattttaagc	aatttagcct	actggtgggt	acatcaatgt	aaacttggtg	5400
gaaaatgtat	ttccaggtaa	ctggtgaatt	gtcataccat	atagagctat	tagatgtaat	5460
ccaattaaat	gttagatgac	atctgccaaa	gaagatgatt	ctgatttggg	catgttggtt	5520
ctaagcaatg	cttttttttt	tttgtagtaa	gtggagagta	agatcaaaca	tccaccgtgg	5580
ccactcttag	ctgatgctaa	taaaaccatg	ttgtcaggac	tggttttctg	ttagagttag	5640
gaccattggt	ccaatttttc	cagaagtaga	atcaggcttc	aaggaatgag	ttaaaagagc	5700
atagggacaa	atggctctata	aaacctataa	tagaaaggag	gtgaaagcgg	ggtggtggcc	5760
cagtgcctga	agagatactg	tagtgcatct	acaggccaaa	taagagaaaag	tagaacaata	5820
gtatgaatca	gagatgatgg	aggctgctac	ttgatagaac	actttctttt	tggaactgaa	5880
aatagccatg	ttggtataag	ggttagggcc	aggcctgacc	ctaattgtaag	cgccttagca	5940
tgggatgtga	tatggagatg	tgcatagcag	aaacctaaaa	tctccctgtg	tcagaaaaaa	6000
atgttagctc	actggccagg	caacaagact	tgcacacttt	ccaaaaagag	gtggtcatgg	6060
agagagcatt	gttaagaatg	ggctgcatgt	caggccaggt	gagtggatga	gaagccacct	6120
ctgtgatgag	acttggtcta	tccaggccct	ggaggagaga	gtacaggctg	cctcaaaagt	6180
gaggctggga	cctgatggaa	atggatttgg	agtcccagct	gacaaagtaa	gtagaatggt	6240
aaatctgtga	tattctcccc	catgctttca	ctatcctcaa	gtttgatgtg	ccccaggcaa	6300
gcaaagattg	acaaaggaag	caaatgactc	tagatgattg	agcaccacca	ccacgtagat	6360
caaatttcca	aagtatagat	ttctgatcac	tagaatccca	gggaaaagag	aagcactgct	6420
ttgatttcta	ggtagcttat	cctaggacaa	ctgcccaacta	gtctcagatg	ggttgctcgt	6480
cttgccctgga	ttccttctct	tcagctccct	gtctctatct	caaatatgca	tcttttctatt	6540
tcctctgcac	tagaacgtag	gttaagcggg	cttggttagcc	ttttccttct	gaaaatgcag	6600
actcctagct	caaatttctt	ggatcacctg	gcccagtgat	cacttatagt	ctgctgatta	6660
caggcatcat	tttctctctg	ggtgttttag	tatagattca	ttgaccccag	tcctcctaga	6720
gtagattctc	tgggttagt	gctcaggaat	atgcaagatc	acaagctctt	ttgtaccatg	6780
tgctttttaga	gctatcactg	tagtgttttc	ctgacagctc	agattttctta	tctcag	6836

<210> 47

<211> 603

<212> DNA

<213> *Cyprinus carpio*

<400> 47

gcgagctctg	taatcatgaa	gcttattctt	gctgtgctgg	tgctggcgct	ggttttggtg	60
gtcgagactc	aggctcaatg	gcaccgctac	ccaggacaag	ccattggagg	tgcaaaaggac	120
atgtggcggtg	cttaccggga	tatgaggaag	gccaaactgga	agggcgcgaga	caagtatttc	180
catgcacgtg	ggaactatga	tgctgcaaga	aggggccccg	gaggcagatg	ggcagccaaa	240
gtaatcagta	atggaagaga	ggctctgcag	ggactctctc	gccgcggtaa	ttcagatgct	300
gcagcagacc	aggccgccaa	tcgctgggga	cgtaacgggtg	gtaaccccaa	ccgctacaga	360
ccgagggggcc	ttcccaaaaa	gtactgaaaa	gaagagcatc	actgcaacag	aaaatgactg	420
aaatcatttt	gctctagaac	tgcttgatct	aattttgcat	aaaaccattt	cacgcttggt	480
ttttttcagt	cttgacaatt	atcttgcat	aaatctgcac	ttatgacaaa	aatgcaatct	540
tttaagtttt	agcatcattt	agcctatgta	acacactctg	gtttgaataa	aggaagtaac	600

ttc

603

<210> 48

<211> 701

<212> DNA

<213> *Dermacentor variabilis*

<400> 48

acagtttgag	atcccggtccg	caaaaagaaa	ccgacgaaga	tggccgctac	tcagccccgt	60
cagaactacc	acgtcgaatg	cgaagcccg	atcaacaagc	agatcaacat	ggaactctac	120
gcgagctacg	tctacacatc	aatggcctac	tactttgacc	gcgatgatgt	ggcactgccg	180
ggattccaca	agttcttcaa	gaaaagcagc	gaggaggagc	gtgaacatgc	tgagaagctg	240
atgaagtatc	agaatatgcg	tggaggggcg	gtgggtgctgc	agccaatcca	gaagccagca	300
caggacgaat	ggggtgcagg	cttggaacgc	atgcaggctt	cactggagct	ggagaagact	360
gttaaccagt	cactgcttga	cctgcacaag	ctggcgactg	atcacaatga	cgctcagctg	420
tgtgacttcc	tggaaagtga	gtacctggca	gagcagggtga	aggccatcaa	ggagctgtcc	480
gattatgtga	ccaacttgaa	gcgtgtttgg	cctggtcttg	gagagtacat	gtttgacaag	540
gagaccctgt	cagactaggc	tctacatgca	gacacaagat	gtgggtctgtc	tgttggtgaa	600
gggcttgca	tcattgtgcc	actgtgtacc	atttggcccc	ttctaccatg	caagaataaa	660
atagttcaaa	gcaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	a		701

<210> 49

<211> 1234

<212> DNA

<213> *Mus sp.*

<400> 49

ggagtcagca	cagcccagcc	cttcagagaga	gaggcaagag	aggccacga	tgagagccct	60
gggagctggt	gtcactctcc	tgctctgggg	tcagcttttt	gctgtggagt	tgggcaatga	120
tgccatggac	tttgaagatg	acagctgccc	aaagccccc	gagattgcaa	acggctatgt	180
ggagcacttg	gttcgctatc	gctgccgaca	gttctacaga	ctacggggccg	aaggagatgg	240
ggtgtacacc	ttaaaccgacg	agaagcaatg	ggtgaacaca	gtcgtggac	acaaactccc	300
cgaatgtgag	gcagtgtgtg	ggaagcccaa	gcaccctgtg	gaccaggtgc	agcgcacatc	360
cggtggctct	atggatgcca	aaggcagctt	tccttggcag	gccaagatga	tctcccgcca	420
cggactcacc	accggggcca	cggtgatcag	tgaccagtgg	ctgctgacca	cggccaaaaa	480
cctcttctctg	aaccacagcg	agacggcgtc	agccaaggac	atcaccccca	ccctaaccgt	540
ctacgtaggg	aaaaaccagc	tggtggagat	tgagaaggtc	gttctccacc	ccaaccactc	600
cgtgggtgat	atcgggctaa	tcaaactcaa	gcagaggggtg	cttgtaaccg	agagagtcac	660
gcctatctgc	ctgccttcca	aagactacat	agcaccaggc	cgtgtgggct	acgtgtctgtg	720
ctggggggcg	aacgccaact	ttagatttac	cgatcgtctc	aagtatgtca	tgtgctctgt	780
ggccgaccag	gacaagtgtg	tggtgcacta	tgagaatagt	acagtggccg	agaagaaaaa	840
cttgacgagt	cccgttgggg	tccagcctat	cttgaacgag	cacaccttct	gtgctggcct	900
caccaagtac	caggaagaca	cctgctacgg	tgacgccggc	agtgcctttg	ccattcatga	960
catggaggag	gacacctggt	acgcagctgg	gatcctgagc	tttgacaaga	cgtgcgctgt	1020
cgtgagtat	ggtgtgtacg	tgagggcgac	cgacctgaag	gactgggttc	aggaaaccat	1080
ggccaagaac	tagttcaggg	ctcactagaa	ggctgcacat	ggcagggcag	gctgggagcc	1140
atggaagagg	agggttgggc	tatactctga	tgggttctag	ccctgcactg		1200
ctcagtcaac	aataaaaaaa	tgtgctttgg	accc			1234

<210> 50

<211> 3321

<212> DNA

<213> *Homo sapiens*

<400> 50

atgaagattt	tgatacttgg	tatttttctg	tttttatgta	gtaccccgagc	ctgggcgaaa	60
gaaaagcatt	attacattgg	aattattgaa	acgacttggg	attatgcctc	tgaccatggg	120
gaaaagaaac	ttattttctgt	tgacacggaa	cattccaata	tctatcttca	aaatggccca	180
gatagaattg	ggagactata	taagaaggcc	ctttatcttc	agtacacaga	tgaacctttt	240
aggacaacta	tagaaaaaac	ggtctggcct	gggttttttag	gccctattat	caaagctgaa	300
actggagata	aagtttatgt	acacttaaaa	aaccttgctt	ctaggcccta	cacctttcat	360
tcacatggaa	taacttacta	taaggaacat	gagggggcca	tctaccctga	taacaccaca	420
gattttcaaa	gagcagatga	caaagtatat	ccaggagagc	agtatacata	catgttgctt	480
gccactgaag	aacaaagtcc	tgggggaagga	gatggcaatt	gtgtgactag	gatttaccat	540
tcccacattg	atgctccaaa	agatattgcc	tcaggactca	tcggaccttt	aataatctgt	600

aaaaaagatt	ctctagataa	agaaaaagaa	aaacatattg	accgagaatt	tgtgggtgatg	660
ttttctgttg	tggtatgaaa	tttcagctgg	tacctagaag	acaacattaa	aacctactgc	720
tcagaaccag	agaaagtga	caaagacaac	gaagacttcc	aggagagtaa	cagaatgtat	780
tctgtgaatg	gatacacttt	tggaagtctc	ccaggactct	ccatgtgtgc	tgaagacaga	840
gtaaaatggt	accttttttg	tatgggtaat	gaagttgatg	tgcacgcagc	tttctttcac	900
gggcaagcac	tgactaacia	gaactaccgt	attgacacaa	tcaacctctt	tcctgctacc	960
ctgtttgatg	cttatatggt	ggcccagaac	cctggagaat	ggatgctcag	ctgtcagaat	1020
ctaaaccatc	tgaagccgg	tttgcaagcc	tttttccagg	tccaggagt	taacaagtct	1080
tcatcaaaag	ataatatccg	tggaagcat	gttagacact	actacattgc	cgctgaggaa	1140
atcatctgga	actatgctcc	ctctggtata	gacatcttca	ctaaagaaaa	cttaacagca	1200
cctggaagt	actcagcgg	gttttttgaa	caaggtagca	caagaattgg	aggctcttat	1260
aaaaagctgg	tttatcgtga	gtacacagat	gcctccttca	caaatcgaaa	ggagagaggc	1320
cctgaagaag	agcatcttgg	catcctgggt	cctgtcattt	gggcagaggt	gggagacacc	1380
atcagagtaa	ccttccataa	caaaggagca	tatccccca	gtattgagcc	gattgggggtg	1440
agattcaata	agaacaacga	gggcacatac	tattccccaa	attacaaccc	ccagagcaga	1500
agtgtgcctc	cttcagcctc	ccatgtggca	cccacagaaa	cattcaccta	tgaatggact	1560
gtccccaaag	aagtaggacc	cactaatgca	gatcctgtgt	gtctagctaa	gatgtattat	1620
tctgtctgtg	atcccactaa	agatatattc	actgggctta	ttggggcaat	gaaaatatgc	1680
aagaaaggaa	gtttacatgc	aaatgggaga	cagaaagatg	tagacaagga	attctatttg	1740
tttctacag	tatttgatga	gaatgagagt	ttactcctgg	aagataatat	tagaatgttt	1800
acaactgcac	ctgatcaggt	ggataaggaa	gatgaagact	ttcagggaatc	taataaaatg	1860
cactccatga	atggattcat	gtatgggaat	cagccgggtc	tcactatgtg	caaaggagat	1920
tcggctcgtg	ggtacttatt	cagcgccgga	aatgaggccg	atgtacatgg	aatatacttt	1980
tcaggaaaca	catactctgt	gagaggagaa	cggagagaca	cagcaaacct	cttccctcaa	2040
acaagtctta	cgctccacat	gtggcctgac	acagagggga	cttttaatgt	tgaatgcctt	2100
acaactgatc	attacacagg	cggcatgaag	caaaaatata	ctgtgaacca	atgcaggcgg	2160
cagtctgagg	attccacctt	ctacctggga	gagaggacat	actatatcgc	agcagtggag	2220
gtggaatggg	attattcccc	acaaaggag	tggaagaaag	agctgcatca	tttacaagag	2280
cagaatgttt	caaatgcatt	tttagataag	ggagagtgtt	acataggctc	aaagtacaag	2340
aaagttgtgt	atcggcagta	tactgatagc	acattccgtg	ttccagtggg	gagaaaagct	2400
gaagaagaac	atctgggaat	tctaggtcca	caacttcatg	cagatgttgg	agacaaagtc	2460
aaaattatct	ttaaaaacat	ggccacaagg	ccctactcaa	tacatgcccc	tgggggtacaa	2520
acagagagtt	ctacagttac	tccaacatta	ccagggtgaa	ctctcactta	cgtatggaaa	2580
atcccagaaa	gatctggagc	tggaacagag	gattctgctt	gtattccatg	ggcttattat	2640
tcaactgtgg	atcaagttaa	ggacctctac	agtggtattaa	ttggccccct	tgtgttttgt	2700
cgaagacctt	acttgaaagt	attcaatccc	agaagggaagc	tggaaatttgc	ccttctgttt	2760
ctagtttttg	atgagaatga	atcttggtac	ttagatgaca	acatcaaaac	atactctgat	2820
caccccgaga	aagtaaacia	agatgatgag	gaattcatag	aaagcaataa	aatgcatgct	2880
attaatggaa	gaatgttttg	aaacctacaa	ggcctcacia	tgcacgtggg	agatgaagtc	2940
aactgggtatc	tgatgggaat	gggcaatgaa	atagacttac	acactgtaca	ttttcacggc	3000
catagcttcc	aatacaagca	caggggagtt	tatagttctg	atgtctttga	cattttccct	3060
ggaacatacc	aaacctaga	aatgtttcca	agaacacctg	gaatttggtt	actccactgc	3120
catgtgaccg	accacattca	tgctggaatg	gaaaccactt	acaccgttct	acaaaatgaa	3180
gacaccaa	ctggctgaat	gaaataaatt	ggtgataagt	ggaaaaaaga	gaaaaaccaa	3240
tgattcataa	caatgtatgt	gaaagtgtaa	aatagaatgt	tactttggaa	tgactataaa	3300
cattaaaaga	gactggagca	t				3321

<210> 51

<211> 8878

<212> DNA

<213> Homo sapiens

<400> 51

gaattcatgc	cccttttgaa	atagacttat	gtcattgtca	gaaaacataa	gcatttatgg	60
tatatcatta	atgagtcacg	atttttagtg	ttgccttggt	agtaggtcaa	atttactaag	120
cttagatttg	ttttctcaca	tattctttcg	gagcttgtgt	agtttccaca	ttaatattacc	180
agaaacaaga	tacacactct	ctttgaggag	tgccctaact	tcccatcatt	ttgtccaatt	240
aaatgaattg	aagaaattta	atgtttctaa	actagaccaa	caaagaataa	tagttgtatg	300
acaagtaaat	aagctttgct	gggaagatgt	tgcttaaatg	ataaaaatgg	tcagccaaca	360
agtgaaccaa	aaattaaata	ttaactaagg	aaaggtaacc	atttctgaag	tcattcctag	420
cagaggactc	agatatatat	aggattgaag	atctctcagt	taagtctaca	tgaaaaggat	480
ggtttcttgg	agcttccaca	aacttaaaac	catgaaacat	ctattattgc	tactatttgt	540
tgtttttcta	gttaagtccc	aagggtgtcaa	cgacaatgag	gaggtgaatt	ttttaaagca	600
ttattatatt	attagtagta	ttattaatat	aagatgtaac	ataatcatat	tatgtgctta	660
ttttaatgaa	attagcattg	cttatagtta	tgaaatggaa	ttgttaacct	ctgacttatt	720

gtattttaaag	aatgttttcat	agtattttctt	atataaaaaac	aaagtaatttt	cttggttttct	780
agttttatcac	ctttgttttct	ttaagatgag	gatggcttag	ctaagtgaag	atgtgttttct	840
ctcacttgct	attctgagta	ctgtgattttt	cattttacttc	tagcaataca	ggattacaat	900
taagaggaca	agatctgaaa	atctcacaaa	ctataaaaata	ataaaaagagc	agaatttttaa	960
gataaaagaa	actggttgga	ggtagattgt	tcttttggtga	aggaaggtta	tatatattgt	1020
tactgagatt	actattttata	aaaattataa	ctaagcctaa	aagcaaaaata	catcaagtgt	1080
aatgatagaa	aatgaaatat	tgcttttttct	agatgaaaag	ttcaaattag	agttagtggtg	1140
tattgtttatt	attaatagtt	atgaaacacg	gttcagttcta	atttattttat	ttgtagaaca	1200
gtttgtcctc	aactattatt	tttgctgact	tattgtgtgt	aattttgcagt	tactaaaaat	1260
acagaaatgc	atttaggaca	atggatatttt	aagaaatttta	aatttttatca	tcaaacgtat	1320
catggccaaa	tttctttacat	atagcatagt	atcattaaac	tagaaataag	aatacacaaat	1380
aatattttaaa	tgaagtgtt	catttcggat	cattatttgag	tttcaaggga	acttgagtgt	1440
tgtactttatc	agactctaca	tgtaagaaca	tatagttaat	ctggttggtgt	gtgtaaaaaac	1500
atatggttaa	tctggttaag	tctggttaat	catatttaggt	aagaaaaatg	taaagaatgt	1560
gtaagacgaa	atttttgtaa	agtactctgc	aaagcactttt	cacattttctg	cttatcaact	1620
aaacctcaca	gagatagttt	aatagttttag	gctttaaaaat	ggatttttgat	tattcaacaa	1680
gtggccttca	taattttcttt	aagtgtttttt	ctttaagtat	atacttttctt	taaatatttt	1740
ttaaaatttct	cttttctcta	gtaaagccag	accatccatg	ctacctctct	agtggcactc	1800
tgaaataaaaa	agaaaatagt	tttctctggt	ataattgtat	ttgtaataag	cagatgaatc	1860
acattttctta	aaattttgttt	tagagagggt	aagctctgac	taggaccatg	acttcaatgt	1920
gaaatatgta	tatatcctcc	gaatctttac	atattaaaga	tgtatatagt	caactgggtta	1980
aacaggaaaa	cttggaacag	cctggctggg	ttttaactctt	agcaccatcc	tactaaatgt	2040
taaaataatat	tataatctaa	tgaataaattg	acaatgcaat	tccaaataga	gttcatctga	2100
tgactttctag	actcacaaaa	ttgcaagaga	gctcagttgt	tgctcagttg	ttccaaatca	2160
tgctggtttgt	taattttgtaa	ttaagctcca	aaggatgtat	agctactgac	aaaaaaaaaaa	2220
atgagaatgt	agttaatcca	aatcaaaaact	ttcctatttgc	aatgcgtatt	ttctgcttca	2280
ttatcctttta	atataatatt	ttaagtttagc	aagtaattttt	aattacaatg	cacaagcctt	2340
gagaatttatt	ttaaataataa	gaaaatcata	atgttttgata	aagaaatcat	gtaagaaatt	2400
tcaagataat	ggtttaacaa	ataatttttgt	tgtatagaaga	taagactaaa	agtgaatttc	2460
gaagtggaga	ggacacttaa	actgtagtac	ttgttatgtg	tgattccagt	aaaaatagta	2520
atgagcactt	attattgcca	agtactgttc	tgagggtacc	atatgcaata	agttattttaa	2580
tcctttacaat	aatcttgtaa	ggcagattca	aactatcatt	acactttattt	tacagatgag	2640
aaaactgggg	cacagataaa	gcaacttgcc	caagggtctca	tagctgtaag	tcaaccctac	2700
ggctcaagac	tacaagtagc	cgagctccag	agtaacttat	gagggtcaaa	gattgtctta	2760
ttacaaaataa	attccaagta	gaatcaacct	tttaataagtc	tttaatgtct	cttaaatatg	2820
tttatatagg	agtctaataca	ccaattcaca	aaaatgaaaag	tagggaaatg	attaacaata	2880
atcataggaa	tctaacaatc	caagtggctt	gagaatatctc	attcttcttg	acagtataga	2940
ttcttttaca	tttcgttaagt	tccaatgtat	gttttaggaa	tatgagggtca	ttactatttca	3000
taatctgata	cagcttttatc	ctaaggcctc	tcttttaaaaa	ctacactgca	tcatagcttt	3060
tttggtgcagt	tggtcttttct	actgttactg	aacagtaagc	aacctacaga	ttcactatca	3120
ccaaggaacc	agttgatgga	tcttaagcaa	attatcaagc	ttgtgataac	ctaaattata	3180
aaatgagggt	gtttgaatag	ttacattcca	aatcttctat	aacactctgt	atttatatttc	3240
tgctctcatc	cttgtaggggt	ttcttcagtg	cccggtgtca	tcgaccctt	gacaagaaga	3300
gagaagaggc	tcccagcctg	aggcctgccc	caccgcccac	cagtggagggt	ggctatcggg	3360
ctcgtccagc	caaagcagct	gccactcaaa	agaaagtaga	aagaaaagcc	cctgatgctg	3420
gaggctgtct	tcacgctgac	ccagaccttg	tggtgtcact	gatgttttctt	gcagtggtgg	3480
ctctctcatg	cagagaaagc	ctgtagtcat	ggcagttctgc	taatgttttca	ctgaccaca	3540
ttaccatcac	tgttatttttg	tttgttttatt	ttggaaataa	aattcaaaaac	ataaacatat	3600
tgggcctttg	gttttaggctt	tcttttcttgt	tttcttttggt	ctgggcccac	aattttcaaat	3660
taggatattgt	gggtgcccac	tttccatttg	tatttttgcca	ctgccttttgt	ttagttggta	3720
aaatttttcat	agcccaatta	tatttttttct	ggggtaaagta	atatttttaaa	tctctatgag	3780
agtatgatga	tgacttttcca	atttctggct	ttacagaaaa	ccaaataata	aattttttatg	3840
ttggctaatc	gtatcgctga	atttttctat	gtgctattttt	aacaaatgtc	catgacccaa	3900
atcctttcatc	taatgcctgc	tatttttcttt	gttttttaggg	ggtgttggtgt	cctacaggat	3960
gtcagttgca	agaggctttg	ctacaacagg	aaaggccaat	cagaaatagt	gttgatgagt	4020
taaataacaa	tgtggaagct	gtttcccaga	cctcctcttc	ttccttttcag	tacatgtatt	4080
tgctgaaaga	cctgtggcaa	aagaggcaga	agcaagtaaa	aggtagatat	ccttggtgctt	4140
tccattcgat	tttcagctat	aaaattggaa	ccgttagact	gccacgagaa	tgcatgggtg	4200
tgagaagatt	aacattttctg	ggttagttaa	tagcattcat	acgctttttgg	gcaccttccc	4260
ctgcaacttg	ccagataagc	actatttcagc	tcttatcccc	agtctgacat	cagcaagtgt	4320
gatttttctat	gaaaaattct	actatgactc	cttatttttaa	gtatacaaga	aacttgtgac	4380
tcagaagata	atattttacag	agtggaaaaa	aacccttagc	atttatagtt	ttaacatttg	4440
aggtttttgaa	tgagagagtt	atccataata	tattcaattg	tgttgtggat	aatgacacct	4500
aacctgtgaa	tcttgagggtc	agaatgttga	gtgctgttga	cttggtgggtc	aggaaacagc	4560
tagtgcgtag	gcctggcaca	ggcatctcag	tgagtagcat	accacagtt	ggaaattttt	4620

caaagaaatc	aaaggaatca	tgacatctta	taaatttcaa	ggttctgcta	tacttatgtg	4680
aaatggataa	ataaatcaag	catatccact	ctgtaagatt	gaacttctca	gatggaagac	4740
cccaatactg	ctttctcctc	ttttccctca	ccaaagaaat	aaacaaccta	tttcatttat	4800
tactggacac	aatcttttagc	gtatacctat	ggtaaattac	tagtatgggtg	gtaggatttt	4860
atgttaattt	gtatatgtca	tgcgccaaat	catttccact	aaatatgact	atatatcata	4920
actgcttggt	gatagctcag	tgtttaatag	tttattctca	gaaaatcaaa	attgtatagt	4980
taaatacatt	agttttatga	ggcaaaaatg	ctaactattt	ctacataatt	tcatttttcc	5040
agataatgaa	aatgtagtca	atgagtactc	ctcagaactg	gaaaagcacc	aattatatat	5100
agatgagact	gtgaatagca	atatcccaac	taaccttcgt	gtgcttcggt	caatcctgga	5160
aaacctgaga	agcaaaaatc	aaaagttaga	atctgatgtc	tcagctcaaa	tgggaatattg	5220
tcgcacccca	tgcactgtca	gttgcaatat	tcctgtgggtg	tctggcaaaag	gtaactgatt	5280
cataaacata	tttttagaga	gttccagaag	aactcacaca	ccaaaaataa	gagaacaaca	5340
acaacaacaa	aatgctaag	tggattttcc	caacagatca	taatgacatt	acagtacatc	5400
ataaaaaatat	ccttagccag	ttgtgttttg	gactggcctg	gtgcatttgc	tggttttgat	5460
gagcaggatg	gggcacaggt	agtcccaggg	gtggctgatg	tgtgcatctg	cgtactggct	5520
tgaacagatg	gcagaaccac	agatagatgt	agaagtttct	ccattttgtg	tgttctggga	5580
gctcatggat	attccaggac	acaaaagggtg	gagaagagct	ttgttcatcc	tcttagcaga	5640
taaacgtcct	caaaactggg	ttggacttac	taaagtaaaa	tgaaaatcta	atatttgtta	5700
tattattttc	aaaggtctat	aataacacac	tccttagtaa	cttatgtaat	gttattttaa	5760
agaattgggtg	actaaataca	aagtaattat	gtcataaacc	cctgaacata	atgttgtctt	5820
acattttgcag	aatgtgagga	aattatcagg	aaaggagggtg	aaacatctga	aatgtatctc	5880
attcaacctg	acagttctgt	caaaccgtat	agagtatact	gtgacatgaa	tacagaaaat	5940
ggaggtaagc	tttcgacagt	tgttgacctg	ttgatctgta	attatttggga	taccgtaaaa	6000
tgccaggaaa	caaggccagg	tgtgggtggct	catacctgta	attccagcac	cttggggaggc	6060
caaagtgggc	tgatagcttg	agcctaggag	tttgaaacta	gcctgggcaa	cataatgaga	6120
ccctaactct	acaaaaaaaa	aaaaaatacc	aaaaaaaaaa	aaaaaatcag	ctgtgttgggt	6180
agtatgtgct	tgtagtccca	gctatccagg	aggctgagat	gggagatcac	ctgagcccac	6240
aacctggagt	cttgatcatg	ctactgaact	gtagcctggg	caacagagga	tagtgagatc	6300
ctgtctcaaa	aaaaaaaaaatt	aattaaaaaag	ccaggaaaca	agacttagct	ctaacatcta	6360
acatagctga	caaaggagta	atttgatgtg	gaattcaacc	tgatatttaa	aagttataaa	6420
atatctataa	ttcacaaattt	ggggtaagat	aaagcacttg	cagtttccaa	agattttaca	6480
agtttacctc	tcatattttat	ttccttattg	tgtctatttt	agagcaccaa	atatatacta	6540
aatggaatgg	acaggggatt	cagatattat	tttcaaagtg	acattatttg	ctgttgggta	6600
atatatgtct	ttttgttttc	tgtcaaccaa	aggatggaca	gtgattcaga	accgtgaaag	6660
cggtagtggt	gactttggca	ggaaatggga	tccatataaa	cagggttttg	gaaatgttgc	6720
aaccaacaca	gatgggaaga	attactgtgg	cctaccagggt	aacgaacagg	catgcaaaat	6780
aaaatcattc	tatttgaaat	gggatttttt	ttaattaaaa	aacattcatt	gttgggaagcc	6840
tgttttaggc	agttaagagg	agtttcctga	caaaaatgtg	gaagctaaag	ataaggggaag	6900
aaaggcagtt	tttagtttcc	caaaaatttta	tttttgggtga	gagattttat	tttgtttttc	6960
ttttagggtga	atataggctt	ggaaatgata	aaattagcca	gcttaccagg	atgggaccca	7020
cagaactttt	gatagaaatg	gaggactgga	aaggagacaa	agtaaaggct	cactatggag	7080
gattcactgt	acagaatgaa	gccaacaaat	accagatctc	agtgaacaaa	tacagaggaa	7140
cagccggtaa	tgcctcatg	gatggagcat	ctcagctgat	gggagaaaac	aggaccatga	7200
ccattcacaa	cggcatgttc	ttcagcacgt	atgacagaga	caatgacggc	tggatatgtgt	7260
ggcactcttt	gctcctgctt	taaaaatcac	actaatatca	ttactcagaa	tcattaacaa	7320
tatttttaat	agctaccact	tcctgggcac	ttactgtcag	ccactgtcct	aagctcttta	7380
tgcactactc	gaagcatttt	caactataag	gtagacattc	ttattctcat	ttatcacagt	7440
agatttagag	agattacgtg	atttgtccaa	tgtcacacaa	ctaccagag	ataaaaactag	7500
aatttgagca	cagttacttt	ctgaataatg	agcattttaga	taaaaccta	tatctctata	7560
ttctaaagtg	tgtgtgaaaa	ctttcatattt	catttccagg	gttctctgat	actaaggggtt	7620
gtaaaagcta	ttattccagt	ataaagtaac	aaacacagtc	cctagatgga	ttgccacaaa	7680
ggcccagtta	tctctctttc	ttgctatagg	gcacaggagg	tctttgggtg	attagtgtga	7740
ctctatgtat	agcaccacaa	ggaaagacta	ctgtgcacac	gagtgtagca	gtcttttatg	7800
ggtaatctgc	aaaacgtaac	ttgaccaccg	tagttctggt	tctaataacg	ccaaacacat	7860
tttcttttcag	gttaacatca	gatcccagaa	aacagtgttc	taaagaagac	ggtgggtggat	7920
ggtgggtataa	tagatgtcat	gcagccaatc	caaacggcag	atactactgg	ggtggacagt	7980
acacctggga	catggcaaaag	catggcacag	atgatgggtg	agtatggatg	aattggaagg	8040
ggtcatggta	ctcaatgagg	aagatgagta	tgaagatcag	gcccttcttc	ccacagcaat	8100
attgcccacat	acgtagattt	ttgctcttct	gtatgtgaca	acatttttgt	acattatgtt	8160
agtggaattt	tctttcatatc	atttatattcc	tctaaaactc	tcaagcagac	gtgagtgtga	8220
ctttttgaaa	aaagtatagg	ataaattaca	ttaaaatagc	acatgatttt	cttttgtttt	8280
cttcattttct	cttgctcacc	caagaagtaa	caaaagtata	gttttgacag	agttgggtgtt	8340
cataattttca	gttctagtgt	attgcgagaa	ttttcaaata	aggaagaggg	gtcttttatc	8400
cttgctgtag	gaaaaccatg	acggaaagga	aaaactgatg	tttaaaagtc	cactttttaa	8460
actatattta	tttatgtagg	atctgtcaaa	gaaaacttcc	aaaaagattt	attaattaaa	8520

ccagactctg	ttgcaataag	ttaatgtttt	cttgtttttgt	aatccacaca	ttcaatgagt	8580
taggctttgc	acttgtaagg	aaggagaagc	gttcacaacc	tcaaatagct	aataaacagg	8640
tcttgaatat	ttgaagattt	aaaatctgac	tctaggacgg	gcacgggtggc	tcacgactat	8700
aatcccaaca	ctttggggagg	ctgaggcggg	cggtcacaag	gtcaggaggtt	caagaccagc	8760
ctgaccaata	tggtgaaacc	ccatctctac	taaaaatata	aaaatttagcc	aggcgtggtg	8820
gcaggtgcct	gtaggtccca	gctagcctgt	gaggtggaga	ttgcattgag	ccaagatc	8878

<210> 52

<211> 2182

<212> DNA

<213> Homo sapiens

<400> 52

gtctaggagc	cagccccacc	cttagaaaag	atgtttttcca	tgaggatcgt	ctgcctagtt	60
ctaagtgttg	tgggcacagc	atggactgca	gatatgtgtg	aaggtgactt	tctagctgaa	120
ggaggaggcg	tgcgtggccc	aagggttgtg	gaaagacatc	aatctgcctg	caaagattca	180
gactggccct	tctgctctga	tgaagactgg	aactacaaat	gcccttctgg	ctgcaggatg	240
aaagggttga	ttgatgaagt	caatcaagat	tttacaacaa	gaataaataa	gctcaaaaat	300
tcactatttg	aatatcagaa	gaacaataag	gattctcatt	cgttgaccac	taatataatg	360
gaaattttga	gaggcgatth	ttcctcagcc	aataaccgtg	ataataccta	caaccgagtg	420
tcagagatc	tgagaagcag	aattgaagtc	ctgaagcgca	aagtcataga	aaaagtacag	480
catatccagc	ttctgcagaa	aaatgttaga	gctcagttgg	ttgatatgaa	acgactggag	540
gtggacattg	atattaagat	ccgatcttgt	cgagggtcat	ggagtagggc	tttagctcgt	600
gaagtagatc	tgaaggacta	tgaagatcag	cagaagcaac	ttgaacaggt	cattgccaaa	660
gacttacttc	cctctagaga	taggcaacac	ttaccactga	taaaaatgaa	accagttcca	720
gacttggttc	ccggaaatth	taagagccag	cttcagaagg	tacccccaga	gtggaaggca	780
ttaacagaca	tgccgcagat	gagaatggag	ttagagagac	ctggtggaaa	tgagattact	840
cgaggaggct	ccacctctta	tggaaaccgga	tcagagacgg	aaagccccag	gaaccctagc	900
agtgtctggaa	gctggaactc	tgggagctct	ggacctggaa	gtactggaaa	cgaaaccctt	960
gggagctctg	ggactggagg	gactgcaacc	tggaaacctg	ggagctctgg	acctggaagt	1020
gctggaagct	ggaactctgg	gagctctgga	actggaagta	ctggaaacca	aaaccttgga	1080
agtcctagac	ctggtagtac	cggaaacctg	aatcctggca	gctctgaacg	cggaagtgtt	1140
gggcactgga	cctctgagag	ctctgtatct	ggtagtactg	gacaatggca	ctctgaatct	1200
ggaagtthta	ggccagatag	cccaggctct	gggaacgcga	ggcctaacaa	cccagactgg	1260
ggcacatttg	aagagggtgtc	aggaaatgta	agtcacggga	caaggagaga	gtaccacaca	1320
gaaaaactgg	tcacttctaa	aggagataaa	gagctcagga	ctggtaaaga	gaaggtcacc	1380
tctggtagca	caaccaccac	gcgtcggtca	tgctctaaaa	ccgttactaa	gactgttatt	1440
ggtcctgatg	gtcacaacaa	agttaccaa	gaagtgtgtg	cctccgaaga	tggttctgac	1500
tgtcccgagg	caatggatth	aggcacattg	tctggcatag	gtactctgga	tgggttccgt	1560
cataggcacc	ctgatgaagc	tgccttcttc	gacactgcct	caactggaaa	aacattccca	1620
ggtttcttct	cacctatgth	aggagagtht	gtcagtgaga	ctgagtctag	gggtcagaa	1680
tctggcatct	tcacaaatac	aaaggaaatcc	agttctcatc	acctgggat	agctgaatth	1740
ccttcccgtg	gtaaatcttc	aagttacagc	aaacaattta	ctagtagcac	gagttacaac	1800
agaggagact	ccacatttga	aagcaagagc	tataaaatgg	cagatgaggc	cggaagtga	1860
gccgatcatg	aaggaaacaa	tagcaccaag	agagggcag	ctaaatctcg	cctgttcaga	1920
ggtatccaca	cttctcctth	ggggaagcct	tccctgtccc	cctagactaa	gttaaatatt	1980
tctgcacagt	gttcccattgg	ccccttgcat	ttccttctta	actctctgtt	acacgtcatt	2040
gaaactacac	tttttttggtc	tgttttttgtg	ctagactgta	agttccttgg	gggcaggggc	2100
tttgtctgtc	tcactctctgt	attcccaaat	gcctaacagt	acagagccat	gactcaataa	2160
atacatgtta	aatggatgaa	tg				2182

<210> 53

<211> 737

<212> DNA

<213> Homo sapiens

<400> 53

cctcctggte	tcagtatggc	gctgtcctgg	gttcttacag	tcctgagcct	cctacctctg	60
ctggaagccc	agatcccatt	gtgtgccaac	ctagtagcgg	tgcccatcac	caacgccacc	120
ctggaccaga	tcactggcaa	gtggttttat	atcgcatcgg	cctttcgaaa	cgaggagtac	180
aataagtcgg	ttcaggagat	ccaagcaacc	ttcttttact	tcacccccaa	caagacagag	240
gacacgatct	ttctcagaga	gtaccagacc	cgacaggacc	agtgcattct	taacaccacc	300
tacctgaatg	tccagcgggg	aaatgggacc	atctccagat	acgtgggagg	ccaagagcat	360
ttcgctcact	tgctgatcct	cagggacacc	aagacctaca	tgcttgctth	tgacgtgaac	420
gatgagaaga	actgggggct	gtctgtctat	gctgacaagc	cagagacgac	caaggagcaa	480

ctgggagagt	tctacgaagc	tctcgactgc	ttgcgcattc	ccaagtcaga	tgtcgtgtac	540
accgattgga	aaaaggataa	gtgtgagcca	ctggagaagc	agcacgagaa	ggagaggaaa	600
caggaggagg	gggaatccta	gcaggacaca	gccttggtac	aggacagaga	cttggggggcc	660
atcctgcccc	tccaacccga	catgtgtacc	tcagcttttt	ccctcacttg	catcaataaa	720
gcttctgtgt	ttggaac					737

<210> 54

<211> 1409

<212> DNA

<213> Bos taurus

<400> 54

gccctggtga	ggatcatgtc	gctgtttaca	tcacttcctt	ttcttctcct	gactgcggtg	60
acagcatctt	gtgcagacac	agaaacagag	aactgtgaga	acatccggaa	gacctgcccc	120
gtgattgcct	gtggtcctcc	gggcatcaat	ggcatcccag	gcaaagatgg	gcgtgatggt	180
gccaaaggag	aaaagggaga	accaggtcaa	ggactcagag	gctcgcaggg	ccccctgga	240
aagatggggc	ctcaagggaac	gccagggatc	cctgggatac	caggaccaat	aggccaaaaa	300
ggagaccctg	gagaaaatat	gggtgactat	attcgcttgg	ctacttcaga	aagagcaact	360
ctacaatctg	aattgaacca	gatcaaaaaa	tggctaattc	tctctctggg	caaaagagtt	420
gggaagaagg	catttttttac	caatggtaaa	aagatgcctt	ttaatgaagt	gaagactctg	480
tgtgcacagt	tccaggggcg	tgtggccacc	cctatgaatg	ctgaagaaaa	caggggccctc	540
aaggatttag	tcactgaaga	ggccttcctg	ggcatcacag	atcaggagac	tgaaggcaaa	600
tttgtggatc	tgacaggaaa	gggggtgacc	tacaaaaact	ggaatgatgg	cgagcctaac	660
aacgcttctc	ctggggagca	ctgtgtgaca	cttctgtcgg	acggcacatg	gaatgacatc	720
gcttggtccg	cctccttttt	gaccgtctgt	gaattctctc	tctgagggag	aatgagccta	780
aagtcctcct	gttccttttac	tcactctcatg	ggccccacaac	ctgggtttgga	ggataaatct	840
atgtcaattt	cacacaccca	gtactgagtt	gctcttttgt	ggggaaacag	agacaataag	900
aatggattga	gaatgatgag	atgtggaaact	gaaaagcgtg	aagagactta	tagtgggtga	960
agagtttctg	gttcagagccc	agtcactaat	aataatcatt	tcaggaacaa	caaaaataatg	1020
gtagtaatat	tagcaacagc	agtaatatgtg	ggagtactaa	taacatatatt	taaaatgttt	1080
actatgagtc	agacattaca	tgtaaagatta	tacattaagt	atctaattta	actaggctgt	1140
tcattttttgt	ttgagactat	aaagaaaagct	gagcactgaa	gaattgatgg	ttttgaactg	1200
tgggtgttga	gaagactcct	aagagtcctt	tggactgcaa	ggagatccaa	ccagtcctac	1260
ttaaaggaga	tcagtcctag	gtgttcattg	gaaggactga	agttgaagct	gaaactccaa	1320
tactttggcc	acctgatgca	aagagctgac	tcatttgaaa	agacctgaa	gctgggaaag	1380
attgagggca	ggagaaaaag	ccggaattc				1409

<210> 55

<211> 1463

<212> DNA

<213> Homo sapiens

<400> 55

tgcactggga	atctaggatg	ggggccttgg	ccagagccct	gccgtccata	ctgctggcat	60
tgtgtcttac	gtccacccca	gaggctctgg	gtgccaaacc	cggcttggtc	gccaggatca	120
ccgacaagg	actgcagtat	gcggcccagg	aggggctatt	agctctgcag	agtgagctgc	180
tcaggatcac	ctgcctgac	ttcacccggg	acttgaggat	ccccacgtc	ggcgtgggc	240
gctatgagtt	ccacagcctg	aacatccaca	gctgtgagct	gcttcactct	gcgctgaggc	300
ctgtccctgg	ccaggggcctg	agtctcagca	tctccgactc	ctccatccgg	gtccagggca	360
ggtggaagg	gcgcaagtca	ttcttcaaac	tacagggtct	ctttgatgtc	agtgtcaagg	420
gcacacagcat	ttcgggtcaac	ctcctgttgg	gcagcgagtc	ctccgggagg	cccacagtta	480
ctgcctccag	ctgcagcagt	gacatcgctg	acgtggagg	ggacatgtcg	ggagacttgg	540
ggtggctgtt	gaacctcttc	cacaaccaga	ttgagtccaa	gttccagaaa	gtactggaga	600
gcaggatttg	cgaaatgatc	cagaaatcgg	tgtcctccga	tctacagcct	tatctccaaa	660
ctctgccagt	tacaacagag	attgacagtt	tcgccgacat	tgattatagc	ttagtgggaag	720
cccctcgggc	aacagcccag	atgctggagg	tgatgtttaa	gggtgaaatc	tttcatcgta	780
accaccgttc	tccagttacc	ctccttgctg	cagtcatgag	ccttcctgag	gaacacaaca	840
aaatggtcta	ctttgccatc	tcggattatg	tcttcaacac	ggccagcctg	gtttatcatg	900
aggaaggata	cttgaacttc	tccatcacag	atgacatgat	accgcctgac	tctaataacc	960
gactgaccac	caagtccttc	cgacctctcg	tcccacgggt	agccaggctc	taccccaaca	1020
tgaacctgga	actccaggga	tcagtgcctt	ctgctccgct	cctgaacttc	agccctggga	1080
atctgtctgt	ggacccctat	atggagatag	atgcctttgt	gctcctgccc	agctccagca	1140
aggagcctgt	cttccggctc	agtgtggcca	ctaattgtgt	cgccaccttg	accttcaata	1200
ccagcaagat	cactgggttc	ctgaagccag	gaaaggtaaa	agtggaaactg	aaagaatcca	1260
aagttggact	attcaatgca	gagctgttgg	aagcgtcctt	caactattac	atccttaaca	1320

ccctctaccc	caagttcaat	gataagttgg	ccgaaggctt	cccccttctt	ctgctgaagc	1380
gtgttcagct	ctacgacctt	gggctgcaga	tccataagga	cttcctgttc	ttgggtgcca	1440
atgtccaata	catgagagtt	tga				1463

<210> 56

<211> 4577

<212> DNA

<213> Homo sapiens

<400> 56

gctacaatcc	atctgggtctc	ctccagctcc	ttctttctgc	aacatgggga	agaacaaact	60
ccttcaccca	agtctgggtc	ttctcctctt	ggctcctctg	cccacagacg	cctcagtctc	120
tggaaaaccg	cagtatatgg	ttctgggtccc	ctccctgctc	cacactgaga	ccactgagaa	180
gggctgtgtc	cttctgagct	acctgaatga	gacagtgact	gtaagtgtt	ccttggagtc	240
tgtcagggga	aacaggagcc	tcttcaactga	cctggaggcg	gagaatgacg	tactccactg	300
tgctgccttc	gctgtcccaa	agtcttcate	caatgaggag	gtaatgttcc	tcactgtcca	360
agtgaaggga	ccaaccaag	aatttaagaa	gcggaccaca	gtgatgggta	agaacgagga	420
cagtctgggtc	tttgtccaga	cagacaaatc	aatctacaaa	ccagggcaga	cagtgaatt	480
tcgtgttggtc	tccatggatg	aaaactttca	ccccctgaat	gagttgattc	cactagtata	540
cattcaggat	cccaaaggaa	atcgcatcgc	acaatggcag	agtttccagt	tagaggggtg	600
cctcaagcaa	ttttcttttc	ccctctcttc	agagcccttc	cagggtcctc	acaagggtgt	660
ggtagagaag	aaatcaggtg	gaaggacaga	gcaccctttc	accgtggagg	aatttgttct	720
tcccaagttt	gaagtacaag	taacagtgcc	aaagataatc	accatcttgg	aagaagagat	780
gaatgtatca	gtgtgtggcc	tatacacata	tgggaagcct	gtccctggac	atgtgactgt	840
gagcattttgc	agaaagtata	gtgacgcttc	cgactgccac	ggtgaagatt	cacaggcttt	900
ctgtgagaaa	ttcagtggac	agctaaacag	ccatggctgc	ttctatcagc	aagtaaaaaac	960
caaggtcttc	cagctgaaga	ggaaggagta	tgaaatgaaa	cttcacactg	aggcccagat	1020
ccaagaagaa	ggaacagtgg	tggaattgac	tggaaggcag	tccagtgaat	tcacaagaac	1080
cataaccaaa	ctctcatttg	tgaaagtggg	ctcacacttt	cgacagggaa	ttcctctctt	1140
tgggcagggtg	cgcttagtag	atgggaaagg	cgctccctata	ccaaataaag	tcatattcat	1200
cagaggaaat	gaagcaaact	attactccaa	tgctaccacg	gatgagcatg	gccttgtaca	1260
gttctctatc	aacaccacca	acgttatggg	tacctctctt	actgttaggg	tcaattacaa	1320
ggatcgtagt	ccctgttacg	gctaccagt	gggtgcagaa	gaacacgaag	aggcacatca	1380
cactgcttat	cttgtgttct	ccccaaagca	gagctttgtc	caccttgagc	ccatgtctca	1440
tgaactaccc	tgtggccata	ctcagacagt	ccaggctcat	tatatctctga	atgggagcac	1500
cctgctgggg	ctgaagaagc	tctcctttta	ttatctgata	atggcaaagg	gaggcattgt	1560
ccgaactggg	actcatggac	tgcttgtgaa	gcaggaagac	atgaagggcc	atttttccat	1620
ctcaatccct	gtgaagtcag	acattgctcc	tgctgctcgg	ttgctcatct	atgctgtttt	1680
acctaccggg	gacgtgattg	gggattctgc	aaaatatgat	gttgaaaatt	gtctggccaa	1740
caaggtggat	ttgagcttca	gcccacaca	aagtcctcca	gcctcacacg	cccacctgcg	1800
agtcacagcg	gtcctcagtc	cgtctgcgc	cctccgtgct	gtggaccaaa	gcgtgctgct	1860
catgaagcct	gatgctgagc	tctcggcgtc	ctcgggtttac	aacctgctac	cagaaaagga	1920
cctcactggc	ttccctgggc	ctttgaatga	ccaggacgat	gaagactgca	tcaatcgtca	1980
taatgtctat	attaatggaa	tcacatatac	tccagtatca	agtacaaatg	aaaaggatat	2040
gtacagcttc	ctagaggaca	tgggcttaaa	ggcattcacc	aactcaaaga	ttcgtaaacc	2100
caaaatgtgt	ccacagcttc	aacagtatga	aatgcatgga	cctgaaggtc	tacgtgtagg	2160
tttttatgag	tcagatgtaa	tgggaagagg	ccatgcacgc	ctggtgcatg	ttgaagagcc	2220
tcacacggag	accgtacgaa	agtacttccc	tgagacatgg	atctgggatt	tgggtggtgt	2280
aaactcagca	ggggtggctg	aggtaggagt	aacagtccct	gacaccatca	ccgagtggaa	2340
ggcagggggc	ttctgcctgt	ctgaagatgc	tggacttggt	atctcttcca	ctgcctctct	2400
ccgagccttc	cagcccttct	ttgtggagct	tacaatgcct	tactctgtga	ttcgtggaga	2460
ggccttcaca	ctcaaggcca	cggtccttaa	ctaccttccc	aaatgcatcc	gggtcagtgt	2520
gcagctggaa	gctcctcccg	ccttctcttg	tgtcccagtc	gagaaggaa	aagcgcctca	2580
ctgcatctgt	gcacaacggc	ggcaactgt	gtcctgggca	gtaaccccaa	agtcattagg	2640
aaatgtgaat	ttcactgtga	gcgcagaggc	actagagtct	caagagctgt	gtgggactga	2700
ggtgccttca	gttctctgaac	acggaaggaa	agacacagtc	atcaagcctc	tgttgggtga	2760
acctgaagga	ctagagaagg	aaacaacatt	caactcccta	ctttgtccat	caggtggtga	2820
ggtttctgaa	gaattatccc	tgaactgcc	accaaagtgt	gtagaagaat	ctgcccagagc	2880
ttctgtctca	gttttgggag	acatatagg	ctctgccatg	caaaacacac	aaaatcttct	2940
ccagatggcc	tatggtgtg	gagagcagaa	tatggtcctc	tttgctccta	acaatctatgt	3000
actggattat	ctaaatgaaa	cacagcagct	tactccagag	gtcaagtcca	aggccattgg	3060
ctatctcaac	actggttacc	agagacagtt	gaactacaaa	cactatgatg	gctcctacag	3120
cacctttggg	gagcgatatg	gcaggaacca	gggcaacacc	tggctcacag	cctttgttct	3180
gaagactttt	gcccagctc	gagcctacat	cttcatcgat	gaagcacaca	ttacccaagc	3240
cctcatatgg	ctctcccaaga	ggcagaagga	caatggctgt	ttcaggagct	ctgggtcact	3300

gctcaacaat	gccataaagg	gaggagtaga	agatgaagtg	accctctccg	cctatatcac	3360
catcgccctt	ctggagattc	ctctcacagt	cactcaccct	gttgtccgca	atgcccctgtt	3420
ttgcctggag	tcagcctgga	agacagcaca	agaaggggac	catggcagcc	atgtatatac	3480
caaagcactg	ctggcctatg	cttttgcctt	ggcaggtaac	caggacaaga	ggaagggaagt	3540
actcaagtca	cttaatgagg	aagctgtgaa	gaaagacaac	tctgtccatt	gggagcgccc	3600
tcagaaaccc	aaggcaccag	tggggcattt	ttacgaaccc	caggctccct	ctgctgaggt	3660
ggagatgaca	tcctatgtgc	tcctcgctta	tctcacggcc	cagccagccc	caacctcgga	3720
ggacctgacc	tctgcaacca	acatcgtgaa	gtggatcacg	aagcagcaga	atgcccaggg	3780
cggtttcttc	tccaccaggg	acacagtggg	ggctctccat	gctctgtcca	aatatggagc	3840
cggcacattt	accaggactg	ggaaggctgc	acaggtgact	atccagtctt	cagggacatt	3900
ttccagcaaa	ttccaagtgg	acaacaacaa	tcgcctgtta	ctgcagcagg	tctcattgcc	3960
agagctgcct	ggggaatata	gcatgaaagt	gacaggagaa	ggatgtgtct	acctccagac	4020
ctccttgaaa	tacaatatct	tcccagaaaa	ggaagagttc	ccctttgctt	taggagtgca	4080
gactctgect	caaacttgtg	atgaacccaa	agccccacac	agcttccaaa	tctccctaag	4140
tgtcagttac	acagggagcc	gctctgcctc	caacatggcg	atcgttgatg	tgaagatggg	4200
ctctggcttc	attccccctga	agccaacagt	gaaaatgctt	gaaagatcta	accatgtgag	4260
ccggacagaa	gtcagcagca	accatgtctt	gattttacct	gataagggtg	caaatacagc	4320
actgagcttg	ttcttcacgg	ttctgcaaga	tgtcccagta	agagatctca	aaccagccat	4380
agtgaagtc	tatgattact	acgagacgga	tgagtttgca	atcgctgagt	acaatgctcc	4440
ttgcagcaaa	gatcttgtaa	atgcttgaag	accacaaggc	tgaaaagtgc	tttgctggag	4500
tcctgttctc	tgagctccac	agaagacacg	tgtttttgta	tctttaaaga	cttgatgaat	4560
aaacactttt	tctggtgc					4577

<210> 57

<211> 4090

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<222> (1)...(4090)

<223> n = A,T,C or G

<400> 57

tgagcagcac	taacctgact	cagtaggttg	tatcagtata	tctattcata	tacacgcaga	60
acagtaataa	ttaaagaaa	agaggccaag	agaggccgag	aacttgagaa	ggaggagggt	120
ggggggtgag	attggagggg	gtagacagag	gaggcaaagt	gtataaatgc	acaaatatgg	180
aatctacaga	gatggtgcc	cgatgctgac	gtaaggctca	ggatttgag	gaaggctcgg	240
gatttgtagg	aagggaatct	tgggttgaa	tctacagaga	acgatatctc	ttctgatcat	300
tgatgcaat	caacacacac	caggagcat	ccatagtggg	cggaatgcat	ggcagccata	360
gtttttggct	gacatctgaa	gcctgtggag	gatttagggg	taggaagaat	gaattccctt	420
aacaagtcaa	aggacaccac	agctccacgc	tgattgcagc	gatgagtaag	aggccctgtc	480
tctttcctcc	agtcgcttc	agtgatctct	attagccatt	gatgcaattc	attcattact	540
ttaagggcca	ttttaagccc	ttaatcttaa	aaccagtccc	aagttgtttc	acgactaaga	600
gactgagctc	caacaagggt	ggctgaaggc	aaggccttct	gtgtgtaact	ttcattttgt	660
tagggattgc	tttccaggtt	ccagtcattg	aatgggcacc	gggcaaacag	cggctcaaagt	720
atttggtcga	aatcaacatc	agttgggtga	gccatgcgtc	atcttcatct	cgagaaggaa	780
tttgatttga	gcaataggtt	agagtttcaa	aatcccagaa	agtccaggaa	cattaaaaaa	840
aaaaaaaaaa	atagaggtgt	atggcagagt	ttcagccgag	agtccacgtg	tgatgtctca	900
gcttcagggt	agggttaaccc	ttttgtgact	ggttcttctc	tactagttag	agcaatagtc	960
ttgggatttc	acaatgctca	gggctcaggt	gatcaactgc	aatgtccctc	tgctgataac	1020
ttccaagaca	gcttctgtca	ttttcagctc	tcaggtgggt	actctgatgg	cctcatgggc	1080
ctgcctgaaa	cagaaagtct	gccacctact	tctgtagcag	caagactcct	gttctgtggc	1140
taagcttcct	gcctgtgcaa	gagccacagg	gaggggccaa	atgcatgcca	ctggggccac	1200
gctcctggta	aagacataaa	tagtgatcct	cgggactggg	catcagagct	ccccttgctc	1260
tcaccatgaa	gtcctgcggc	cttttacctt	tcacgggtgct	ccttgctctg	gggatacctg	1320
cacctgggac	tgtggaagga	ggcaaaaatg	atgggtgagtt	ggagttgctt	tggtccaaca	1380
ttgcctggag	tatcagagtg	ggggctcctg	atgggtttggg	ctgtttttca	gtttactctg	1440
acctcagggc	gacatctgtg	tctcgtctgg	gtctgagctc	tgatgaattt	caaggcatgt	1500
agagtgtctc	tcttatgcag	ccatcctgtc	agctcaggtc	tcacacagct	ctgtccctgc	1560
cctttacctt	tctgtttgtt	tctgtctctg	gctttagttc	tcagcatctc	acatattatt	1620
cccttcctat	ttatttctct	tgggtctccat	tattcctgct	cccaccttct	gtctttgtga	1680
gatcttgctt	agccctgagg	tagaaaacct	tgactgggag	cccggtaacct	tttatcctca	1740
gtctgatttt	tcctggtcac	atgtcttggg	tgaccaagtc	tgacttttcc	tccttaggtc	1800
agccccgtat	ttctctttga	agttgtgggc	cactggaatg	agggctctata	ttctgagatg	1860

tggaccttcg	tctaggaatg	atgatagtc	gtaccctcct	aacatcttga	gcttctcttt	1920
agctatcaaa	atcggagcct	gccctgctaa	aaagcctgcc	cagtgcctta	agcttgagaa	1980
gccacaatgc	cgtactgact	gggagtgc	gggaaagcag	aggtgctgcc	aagatgcttg	2040
cggttccaag	tgcgtgaatc	ctgttcccat	tcgcaaacca	gtgagccagc	agagaaaaca	2100
gagaggggac	aatcaacgtc	aagagccacc	cagagtgaat	gaagagtc	tcccaggc	2160
ccttgcttct	aagagctgtt	gggtccaaca	tgtcagttag	tcaaggcctc	ttgctgccat	2220
tggcaggaag	agccatttgc	ctgcagttgc	cactctgtga	ggatgatggc	tttacaacca	2280
ttcggccctt	tccgggctgg	agagctggtc	ccctgcattg	gctggaggct	agatggtagc	2340
gaggaagcag	gtcctctttc	aggaggcgtg	tttgggagga	ggtgacaaag	gtgatgggtg	2400
gatggggacg	aacttcctcg	ctcaggaagg	agatgttggg	ttaaagacat	ggagatgtgt	2460
tgcctgagcc	ctaagacatc	ggccctgtga	atcctttctc	aacagggaga	tgtgttgctt	2520
gagccctaag	acatcggccc	tgtgaatcct	ttctcaacag	gtgtggagga	agcctgggag	2580
gtgcgtcaaa	actcaggcaa	gatgtatgat	gcttaaccct	cccaatgtct	gccagaggga	2640
cgggcagtg	gacggcaaat	acaagtgtctg	tgagggtata	tgtgggaaag	tctgcctgcc	2700
cccgatgtga	gtgaggctaa	ggagtgggtc	ggggtaggga	gagctcattt	ctctccagcc	2760
cttttctgct	tcaaggctct	ggagctcgaa	gcacatgcgc	caggagattc	tattctaata	2820
aggagccctt	gtctgtttgc	cgtttaaggg	aactgtcccc	tgtttcttag	ggcaaagcct	2880
agcattgcct	gaggctttga	tcctgtggca	gcctgtcagg	gtaaggccct	cgggtgaatg	2940
gctctcagag	tctctgcctg	cacaatagaa	ataccgaagt	tgaagttttg	cgaggcacag	3000
tatacgttat	ttgcttccca	tgttatttac	ttcatggtgt	ctctctggct	agagcacaaa	3060
ggataggtat	aatgtgagga	atgtctagct	cagtgtctctg	cgttttgagt	tttacaacga	3120
caacgagatt	tgaacctgtt	ctcctgaaac	cccagatcct	ctcaagtcca	gccccacttt	3180
accctttgac	cgttctcagc	atgctccctt	acaagggtctg	gataaggaatg	atctagctgg	3240
ttcctatgtt	gaaggctctt	ccttactttc	tcttgtaggc	ctgatccctg	acattggcgc	3300
cggctctgga	ctcgtgctcg	gtgtgctctg	gaaactactt	ccctgctccc	aggcgtccct	3360
gctccgggtt	ccatggctcc	cggctccctg	tatcccaggc	ttggatcctg	tggaccaggg	3420
ttactgtttt	accactaaca	tctccttttg	gctcagcatt	caccgatcct	tagggaaatg	3480
ctgttgagga	gcaaataaat	aaacgcattc	atttctctat	gcacaccgtg	agtctgattc	3540
tctgagggga	tgagatgggt	gtggtaggaa	ccaataaagg	aaggatggga	acaggccctc	3600
cctcttacat	tccgggagga	gaaggatgag	aattccaggc	ttgggcatgc	ttgaacctnc	3660
caaatatcat	gtttttatct	gcaaagtggc	aacagggacc	gctttcttac	aacgtgtagt	3720
gcaccagagg	ttgcagaatg	gtgcaaagtg	tgtgtgtgct	atacagctgt	gctttaccaa	3780
agtgccttca	cacaccacgg	ccacttattg	ctaccggcg	ttatgaaaat	gcgggtgcag	3840
tgtgtgcacg	cacacacatg	tgcagtgtgc	ttacttgtat	gtgtttttga	aggacagagt	3900
angacacagg	gtgtcttctc	ctgtcacttt	cttggttatta	ccttgagtca	ngatctctgt	3960
gctgaaccaa	aagccgacca	atcagggtctg	ggttcgtctg	tctgcccctc	cccttccccg	4020
gttactgggg	ttacangaaa	atgcacacac	cngctttnta	tgtgggttcc	cgggattgac	4080
tgattctgcc						4090

<210> 58

<211> 565

<212> DNA

<213> *Oryctolagus cuniculus*

<400> 58

atggagaccc	ataagcacgg	accttccctg	gcctgggtgt	cactgttgc	gctgctgctg	60
ggcctgctga	tgccccagc	catcgcccag	gacctcacct	accgggaggc	tgtgctccgc	120
gctgtggatg	ccttcaacca	gcagtccctc	gaggccaacc	tctaccgcct	cctgagcatg	180
gacccccagc	agctggagga	tgcgaagcca	tacaccccg	agcctgtgag	ctttacgggtg	240
aaggagacgg	agtgcctccg	gacaacatgg	aagctaccag	agcagtgtga	cttcaaggaa	300
gatgggctgg	tgaagcgggtg	tgtggggact	gtgacacgg	accaggcctg	ggactccttt	360
gacatccgct	gcaacagggc	ccaagagtcc	ccagaacctc	ctgggctg	caagcgctta	420
cgaaaattta	gaaacaagat	taaagaaaag	cttaaaaaaa	ttggtcagaa	aatccagggt	480
ttcgtgccga	aacttgaccc	caggacagat	tactagggtc	tgccctgccc	tggactctga	540
aaaataaact	gtgtgaaagc	aacaa				565

<210> 59

<211> 686

<212> DNA

<213> *Allium cepa*

<400> 59

aacgaaaatt	acgaaattac	atcaatatct	cgagccatgg	ttcgcgttgt	atctttactt	60
gcagcatcga	ccttcatact	gttgattatg	ataatcagca	gtccgtatgc	aaatagtcag	120
aacatatgcc	caagggttaa	tcgaatttgt	acaccctgtg	tggcctacgg	actcgggaag	180

gcaccaatcg	ccccatgctg	cagagccctg	aacgatctac	ggtttgtgaa	tactagaaac	240
ctacgacgtg	ctgcatgccg	ctgcctcgta	ggggtagtga	accggaaccc	cggtctgaga	300
cgaaacccta	gatttcagaa	cattcctcgt	gattgtcgca	acacctttgt	tcgtcccttc	360
tgggtggcgtc	caagaattca	atgcggcagg	attaacctta	cggataagct	tatatacttg	420
gacgctgagg	aatgaagact	aggctctact	gttatgcact	atagtttata	gtatatatac	480
taaataaaaac	agtatgtgct	gtataatttg	caatatggac	ttatttatag	caagtcctaa	540
tgggtgtctgc	tacttgggtc	cagcattgag	cactatatag	gcactatata	gggtactatg	600
ggctgattat	gatgtcaacg	gcggtacttt	atcttacata	aataaataat	gggtttatct	660
tgcttgaaaa	aaaaaaaaaa	aaaaaa				686

<210> 60

<211> 550

<212> DNA

<213> Bos taurus

<400> 60

agactgggga	ccatgcagac	ccagagggcc	agcctctcac	tggggcggtg	gtcgtgtggtg	60
ctactgctgc	tgggactagt	ggtgccctcg	gccagcgccc	aagccctcag	ctacagggag	120
gccgtgcttc	gtgctgtgga	tcagctcaat	gagctgtcct	cagaagctaa	tctctaccgc	180
ctcctggagc	tagaccacc	tcccaaggat	aatgaagatc	tgggcactcg	aaagcctgtg	240
agcttcacgg	tgaaggagac	tgtgtgcccc	aggacgattc	agcagcccg	ggagcagtgt	300
gacttcaagg	agaaagggcg	ggtgaaacag	tgtgtgggga	cagtcaccct	ggacccatcc	360
aatgaccagt	ttgacctaaa	ctgtaatgag	ctccagagtg	tcatectacc	ctggaaatgg	420
ccatgggtggc	cttggcgag	aggttgatgg	agaagagctg	tcagatcctg	agcctcgga	480
agagtcttaa	gtgtctgatt	tgttcagatt	cgggcttctg	gacagtgaaa	ataaattctt	540
gtgaaaacgg						550

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☒ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.